

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56c67e45f0106c67230e73cb

Fecha de creación: undefined

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

0

isbn: 0963699636

keywords: Anas clypeata,Anas platyrhynchos,Ardea herodias,Astragalinus psaltria,Bartramia longicauda,Buteo swainsoni,Cairina moschata,Calidris melanotos,Cardellina canadensis,Cathartes aura,Catharus minimus,Coccyzus americanus,Columba livia,Columbina talpacoti,Contopus cooperi,Contopus virens,Coragyps atratus,Crotophaga ani,Dendrocygna autumnalis,Egretta caerulea,Egretta thula,Empidonax alnorum,Empidonax traillii,Falco columbarius,Fulica americana,Geothlypis philadelphia,Hirundo rustica,Ixobrychus exilis,Jacana jacana,Mniotilta varia,Nycticorax nycticorax,Oxyura jamaicensis,Pandion haliaetus,Pardirallus maculatus,Parkesia noveboracensis,Phalacrocorax brasilianus,Piranga olivacea,Pitangus sulphuratus,Podilymbus podiceps,Porzana carolina,Pyrocephalus rubinus,Rupornis magnirostris,Setophaga fusca,Sturnella magna,Tringa flavipes,Tringa melanoleuca,Tringa solitaria,Troglodytes aedon

0: Alderfer, Jonathan

institution:

address:

publisher: Washington D.C. National Geographic Society

accessed:

websites:

chapter:

series:

pages: 664

issue:

year: 2006

0: Alderfer, J

1: Zickefoose, J

2: Littlehales, B

title: National Geographic Complete Birds of North America

source:

type: book

tags: Anas clypeata,Anas platyrhynchos,Ardea herodias,Astragalinus psaltria,Bartramia longicauda,Buteo swainsoni,Cairina moschata,Calidris melanotos,Cardellina canadensis,Cathartes aura,Catharus minimus,Coccyzus americanus,Columba livia,Columbina talpacoti,Contopus cooperi,Contopus virens,Coragyps atratus,Crotophaga ani,Dendrocygna autumnalis,Egretta caerulea,Egretta thula,Empidonax alnorum,Empidonax traillii,Falco columbarius,Fulica americana,Geothlypis philadelphia,Hirundo rustica,Ixobrychus exilis,Jacana jacana,Mniotilta varia,Nycticorax nycticorax,Oxyura jamaicensis,Pandion haliaetus,Pardirallus maculatus,Parkesia noveboracensis,Phalacrocorax brasilianus,Piranga olivacea,Pitangus sulphuratus,Podilymbus podiceps,Porzana carolina,Pyrocephalus rubinus,Rupornis magnirostris,Setophaga fusca,Sturnella magna,Tringa flavipes,Tringa melanoleuca,Tringa solitaria,Troglodytes aedon

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

keywords: *Accipiter striatus*, *Actitis macularius*, *Adelomyia melanogenys*, *Agelaiocercus kingii*, *Amblycercus holosericeus*, *Ampelion rubrocristatus*, *Anas andium*, *Anas clypeata*, *Anas discors*, *Andigena nigristrois*, *Anisognathus igniventris*, *Ardea alba*, *Ardea herodias*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Asthenes flammulata*, *Asthenes fuliginosa*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Butorides virescens*, *Cairina moschata*, *Calidris bairdii*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Cathartes aura*, *Catharus minimus*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chaetura pelagica*, *Chalcostigma heteropogon*, *Chordeiles minor*, *Chrysomus icterocephalus*, *Cinclodes albidiventris*, *Cinclus leucocephalus*, *Cinnycerthia unirufa*, *Cistothorus apolinari*, *Cistothorus platensis*, *Coccyzua pumila*, *Coccyzus americanus*, *Coccyzus melacoryphus*, *Coeligena bonapartei*, *Colibri coruscans*, *Colibri thalassinus*, *Columba livia*, *Columbina talpacoti*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cinereus*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Dendrocygna autumnalis*, *Dendrocygna viduata*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dolichonyx oryzivorus*, *Dubusia taeniata*, *Egretta caerulea*, *Egretta thula*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Geothlypis philadelphia*, *Geranoaetus melanoleucus*, *Glaucidium jardinii*, *Grallaria quitesis*, *Grallaria ruficapilla*, *Grallaria squamigera*, *Haplospiza rustica*, *Heliangelus amethysticollis*, *Heliangelus exortis*, *Hellmayrea gularis*, *Hemispingus superciliaris*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus icterus*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Lafresnaya lafresnayi*, *Leiothlypis peregrina*, *Leptasthenura andicola*, *Lesbia nuna*, *Lesbia victoriae*, *Machetornis rixosa*, *Margarornis squamiger*, *Mecocerculus leucophrys*, *Megascops choliba*, *Melanerpes rubricapillus*, *Merganetta armata*, *Metallura tyrianthina*, *Milvago chimachima*, *Mimus gilvus*, *Mniotilta varia*, *Molothrus bonariensis*, *Muscisaxicola alpinus*, *Myiarchus crinitus*, *Myiodynastes luteiventris*, *Myiotheretes fumigatus*, *Myiotheretes striaticollis*, *Myiothlypis nigrocristata*, *Nomonyx dominicus*, *Nycticorax nycticorax*, *Ochthoeca cinnamomeiventris*, *Ochthoeca diadema*, *Ochthoeca fumicolor*, *Ochthoeca rufipectoralis*, *Orochelidon murina*, *Oxypogon guerinii*, *Oxyura jamaicensis*, *Pachyramphus versicolor*, *Pandion haliaetus*, *Pardirallus maculatus*, *Parkesia noveboracensis*, *Patagioenas fasciata*, *Petrochelidon pyrrhonota*, *Phaetusa simplex*, *Phalacrocorax brasilianus*, *Pheucticus aureoventris*, *Pheucticus ludovicianus*, *Phimosus infuscatus*, *Phrygilus unicolor*, *Phyllomyias uropygialis*, *Piaya cayana*, *Picoides fumigatus*, *Pipraeidea melanonota*, *Pipreola riefferii*, *Piranga olivacea*, *Piranga rubra*, *Pitangus sulphuratus*, *Pluvialis dominica*, *Podilymbus podiceps*, *Porphyrio martinicus*, *Porzana carolina*, *Progne tapera*, *Protonotaria citrea*, *Pseudocolopteryx acutipennis*, *Pseudoscops clamator*, *Pterophanes cyanopterus*, *Pygochelidon cyanoleuca*, *Pyrocephalus rubinus*, *Pyrrhomyias cinnamomeus*, *Quiscalus lugubris*, *Rallus semiplumbeus*, *Ramphocelus dimidiatus*, *Ramphomicron microrhynchum*, *Riparia riparia*, *Rupornis magnirostris*, *Sayornis nigricans*, *Serpophaga cinerea*, *Setophaga castanea*, *Setophaga cerulea*, *Setophaga petechia*, *Setophaga ruticilla*, *Setophaga striata*, *Sicalis flaveola*, *Sicalis luteola*, *Spiza americana*, *Sporagra spinescens*, *Sturnella magna*, *Synallaxis subpudica*, *Systellura longirostris*, *Tangara vitriolina*, *Thraupis episcopus*, *Thraupis palmarum*, *Tringa flavipes*, *Tringa melanoleuca*, *Tringa solitaria*, *Troglodytes aedon*, *Troglodytes solstitialis*, *Turdus fuscater*, *Turdus ignobilis*, *Tyrannus melancholicus*, *Tyrannus savana*, *Tyrannus tyrannus*, *Tyto alba*, *Vanellus chilensis*, *Vireo olivaceus*, *Vultur gryphus*, *Zenaida auriculata*, *Zonotrichia capensis*

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

pages: 836

issue:

year: 1986

0: Hilty, Steven

1: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

tags: Accipiter striatus,Actitis macularius,Adelomyia melanogenys,Aglaiocercus kingii,Amblycercus holosericeus,Ampelion rubrocristatus,Anas andium,Anas clypeata,Anas discors,Andigena nigrirostris,Anisognathus igniventris,Ardea alba,Ardea herodias,Arremon assimilis,Asio flammeus,Asio stygius,Asthenes flammulata,Asthenes fuliginosa,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Butorides virescens,Cairina moschata,Calidris bairdii,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Cathartes aura,Catharus minimus,Catharus ustulatus,Chaetocercus mulsant,Chaetura pelagica,Chalcostigma heteropogon,Chordeiles minor,Chrysomus icterocephalus,Cinclodes albidiventris,Cinclus leucocephalus,Cinnycerthia unirufa,Cistothorus apolinari,Cistothorus platensis,Coccyzua pumila,Coccyzus americanus,Coccyzus melacoryphus,Coeligena bonapartei,Colibri coruscans,Colibri thalassinus,Columba livia,Columbina talpacoti,Conirostrum rufum,Conirostrum sitticolor,Contopus cinereus,Contopus fumigatus,Contopus sordidulus,Contopus virens,Coragyps atratus,Crotophaga ani,Crotophaga major,Dendrocygna autumnalis,Dendrocygna viduata,Diglossa albilatera,Diglossa cyanea,Diglossa humeralis,Diglossa lafresnayii,Diglossa sittoides,Dolichonyx oryzivorus,Dubusia taeniata,Egretta caerulea,Egretta thula,Elaenia flavogaster,Elaenia frantzii,Elanus leucurus,Empidonax alnorum,Empidonax traillii,Ensifera ensifera,Eriocnemis cupreiventris,Eriocnemis vestita,Eupsittula pertinax,Falco columbarius,Falco peregrinus,Falco sparverius,Forpus conspicillatus,Fulica americana,Gallinago nobilis,Gallinula galeata,Geothlypis philadelphia,Geranoaetus melanoleucus,Glaucidium jardinii,Grallaria quitesis,Grallaria ruficapilla,Grallaria squamigera,Haplospiza rustica,Heliangelus amethysticollis,Heliangelus exortis,Hellmayrea gularis,Hemispingus supercilialis,Hemispingus verticalis,Henicorhina leucophrys,Hirundo rustica,Icterus chrysater,Icterus icterus,Icterus nigrogularis,Ixobrychus exilis,Jacana jacana,Lafresnaya lafresnayi,Leiothlypis peregrina,Leptasthenura andicola,Lesbia nuna,Lesbia victoriae,Machetornis rixosa,Margarornis squamiger,Mecocerculus leucophrys,Megascops choliba,Melanerpes rubricapillus,Merganetta armata,Metallura tyrianthina,Milvago chimachima,Mimus gilvus,Mniotilta varia,Molothrus bonariensis,Muscisaxicola alpinus,Myiarchus crinitus,Myiodynastes luteiventris,Myiotheretes fumigatus,Myiotheretes striaticollis,Myiothlypis nigrocristata,Nomonyx dominicus,Nycticorax nycticorax,Ochthoeca cinnamomeiventris,Ochthoeca diadema,Ochthoeca fumicolor,Ochthoeca rufipectoralis,Orochelidon murina,Oxypogon guerinii,Oxyura jamaicensis,Pachyramphus versicolor,Pandion haliaetus,Pardirallus maculatus,Parkesia noveboracensis,Patagioenas fasciata,Petrochelidon pyrrhonota,Phaetusa simplex,Phalacrocorax brasilianus,Pheucticus aureoventris,Pheucticus ludovicianus,Phimosus infuscatus,Phrygilus unicolor,Phyllomyias uropygialis,Piaya cayana,Picoides fumigatus,Pipraeidea melanonota,Pipreola riefferii,Piranga olivacea,Piranga rubra,Pitangus sulphuratus,Pluvialis dominica,Podilymbus podiceps,Porphyrion martinicus,Porzana carolina,Progne tapera,Protonotaria citrea,Pseudocolopteryx acutipennis,Pseudoscops clamator,Pterophanes cyanopterus,Pygochelidon cyanoleuca,Pyrocephalus rubinus,Pyrrhomyias cinnamomeus,Quiscalus lugubris,Rallus semiplumbeus,Ramphocelus dimidiatus,Ramphomicron microrhynchum,Riparia riparia,Rupornis magnirostris,Sayornis nigricans,Serpophaga cinerea,Setophaga castanea,Setophaga cerulea,Setophaga petechia,Setophaga ruticilla,Setophaga striata,Sicalis flaveola,Sicalis luteola,Spiza americana,Sporagra spinescens,Sturnella magna,Synallaxis subpudica,Systellura longirostris,Tangara vitriolina,Thraupis episcopus,Thraupis palmarum,Tringa flavipes,Tringa melanoleuca,Tringa solitaria,Troglodytes aedon,Troglodytes solstitialis,Turdus fuscater,Turdus ignobilis,Tyrannus melancholicus,Tyrannus savana,Tyrannus tyrannus,Tyto alba,Vanellus chilensis,Vireo olivaceus,Vultur gryphus,Zenaidura macroura,Zonotrichia capensis

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:  
created:  
mimeType:  
dataType:  
identifier:  
habitatUnstructured: La Garza Patiamarilla habita en manglares, estuarios y playones costeros de lodo; menos numerosa y local en el interior en pantanos y lagunas de agua dulce; especialmente numero en el bajo valle del Magdalena y Ciénaga Grande (Hilty & Brown, 1986); es bastante común en humedales y ríos de poca profundidad, bahías, marismas, zonas costeras e islas (Alderfer et al., 2006).

commonNamesAtomized

ancillaryData

usedBy:  
usedIn  
distributionUnstructured:  
endDate:  
startDate:

ancillaryData

type:  
language: Francés  
name: Aigrette neigeuse  
usesManagementAndConservation

ancillaryData

managementAndConservationUnstructured:

managementAndConservationAtomized

identificationKeys

fullDescription

0

dataObject:  
additionalInformation:

0

isbn: 0-618-57424-7  
keywords: Anas clypeata,Anas platyrnchos,Ardea herodias,Bartramia longicauda,Buteo swainsoni,Cairina moschata,Calidris melanotos,Columbina talpacoti,Contopus cooperi,Contopus virens,Crotophaga ani,Dendrocygna autumnalis,Dolichonyx oryzivorus,Egretta caerulea,Egretta thula,Empidonax alnorum,Empidonax traillii,Falco columbarius,Falco sparverius,Geothlypis philadelphia,Hirundo rustica,Leiothlypis peregrina,Myiarchus crinitus,Myiodynastes luteiventris,Nomonyx dominicus,Nycticorax nycticorax,Pandion haliaetus,Parkesia noveboracensis,Petrochelidon pyrrhonota,Podilymbus podiceps,Protonotaria citrea,Sayornis nigricans  
institution:  
edition: Hillstar E  
address:  
publisher:

accessed:

websites:

chapter:

series:

pages: 391

issue:

year: 2005

0: Kaufman, K

title: Guía de campo a las aves de Norteamérica

source:

type: book

tags: *Anas clypeata*, *Anas platyrhynchos*, *Ardea herodias*, *Bartramia longicauda*, *Buteo swainsoni*, *Cairina moschata*, *Calidris melanotos*, *Columbina talpacoti*, *Contopus cooperi*, *Contopus virens*, *Crotophaga ani*, *Dendrocygna autumnalis*, *Dolichonyx oryzivorus*, *Egretta caerulea*, *Egretta thula*, *Empidonax alnorum*, *Empidonax traillii*, *Falco columbarius*, *Falco sparverius*, *Geothlypis philadelphia*, *Hirundo rustica*, *Leiothlypis peregrina*, *Myiarchus crinitus*, *Myiodynastes luteiventris*, *Nomonyx dominicus*, *Nycticorax nycticorax*, *Pandion haliaetus*, *Parkesia noveboracensis*, *Petrochelidon pyrrhonota*, *Podilymbus podiceps*, *Protonotaria citrea*, *Sayornis nigricans*

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

1

keywords: *Accipiter striatus*, *Actitis macularius*, *Adelomyia melanogenys*, *Agelaiocercus kingii*, *Amblycercus holosericeus*, *Ampelion rubrocristatus*, *Anas andium*, *Anas clypeata*, *Anas discors*, *Andigena nigrirostris*, *Anisognathus igniventris*, *Ardea alba*, *Ardea herodias*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Asthenes flammulata*, *Asthenes fuliginosa*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Butorides virescens*, *Cairina moschata*, *Calidris bairdii*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Cathartes aura*, *Catharus minimus*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chaetura pelagica*, *Chalcostigma heteropogon*, *Chordeiles minor*, *Chrysomus icterocephalus*, *Cinclodes albidiventris*, *Cinclus leucocephalus*, *Cinnycerthia unirufa*, *Cistothorus apolinari*, *Cistothorus platensis*, *Coccyzus pumila*, *Coccyzus americanus*, *Coccyzus melacoryphus*, *Coeligena bonapartei*, *Colibri coruscans*, *Colibri thalassinus*, *Columba livia*, *Columbina talpacoti*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cinereus*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Dendrocygna autumnalis*, *Dendrocygna viduata*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dolichonyx oryzivorus*, *Dubusia taeniata*, *Egretta caerulea*, *Egretta thula*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Geothlypis philadelphia*, *Geranoaetus melanoleucus*, *Glaucidium jardinii*, *Grallaria quitesis*, *Grallaria ruficapilla*, *Grallaria squamigera*, *Haplospiza rustica*, *Heliangelus amethysticollis*, *Heliangelus exortis*, *Hellmayrea gularis*, *Hemispingus superciliosus*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus icterus*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Lafresnaya lafresnayi*, *Leiothlypis peregrina*, *Leptasthenura andicola*, *Lesbia nuna*, *Lesbia victoriae*, *Machetornis rixosa*, *Margarornis squamiger*, *Mecocerculus leucophrys*, *Megascops choliba*, *Melanerpes rubricapillus*, *Merganetta armata*, *Metallura tyrianthina*, *Milvago chimachima*, *Mimus gilvus*, *Mniotilta varia*, *Molothrus bonariensis*, *Muscisaxicola alpinus*, *Myiarchus crinitus*, *Myiodynastes luteiventris*, *Myiotheretes fumigatus*, *Myiotheretes striaticollis*, *Myiothlypis nigrocristata*, *Nomonyx dominicus*, *Nycticorax nycticorax*, *Ochthoeca cinnamomeiventris*, *Ochthoeca diadema*, *Ochthoeca fumicolor*, *Ochthoeca rufipectoralis*, *Orochelidon murina*, *Oxygogon guerini*, *Oxyura jamaicensis*, *Pachyramphus versicolor*, *Pandion haliaetus*, *Pardirallus maculatus*, *Parkesia noveboracensis*, *Patagioenas fasciata*, *Petrochelidon pyrrhonota*, *Phaetusa simplex*, *Phalacrocorax brasilianus*, *Pheucticus aureoventris*, *Pheucticus ludovicianus*, *Phimosus infuscatus*, *Phrygilus unicolor*, *Phyllomyias uropygialis*, *Piaya cayana*, *Picoides fumigatus*, *Pipraeidea melanonota*, *Pipreola riefferii*, *Piranga olivacea*, *Piranga rubra*, *Pitangus sulphuratus*, *Pluvialis dominica*, *Podilymbus*

podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

pages: 836

issue:

year: 1986

0: Hilty, Steven

1: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccyzus pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta caerulea, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliosus, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnaya, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Mactornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striatocollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerini, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx

acutipennis,Pseudoscops clamator,Pterophanes cyanopterus,Pygochelidon cyanoleuca,Pyrocephalus rubinus,Pyrrhomyias cinnamomeus,Quiscalus lugubris,Rallus semiplumbeus,Ramphocelus dimidiatus,Ramphomicron microrhynchum,Riparia riparia,Rupornis magnirostris,Sayornis nigricans,Serpophaga cinerea,Setophaga castanea,Setophaga cerulea,Setophaga petechia,Setophaga ruticilla,Setophaga striata,Sicalis flaveola,Sicalis luteola,Spiza americana,Sporagra spinescens,Sturnella magna,Synallaxis subpudica,Systellura longirostris,Tangara vitriolina,Thraupis episcopus,Thraupis palmarum,Tringa flavipes,Tringa melanoleuca,Tringa solitaria,Troglodytes aedon,Troglodytes solstitialis,Turdus fuscater,Turdus ignobilis,Tyrannus melancholicus,Tyrannus savana,Tyrannus tyrannus,Tyto alba,Vanellus chilensis,Vireo olivaceus,Vultur gryphus,Zenaida auriculata,Zonotrichia capensis

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

data\_type:

identifier:

fullDescriptionUnstructured: La Garza Patiámarilla es completamente blanca mide aproximadamente 61 cm y pesa 375 g; sus patas y pico son negros; dedos amarillo brillante, bridas amarillas. En plumaje nupcial tiene plumas largas, recurvadas (airones) en coronilla, pecho y espalda (Hilty & Brown, 1986). Los juveniles pueden tener patas verdosas en un principio, base del pico gris (Kaufman, 2005).

briefDescription:

## references

link: <http://maps.iucnredlist.org/map.html?id=22696974>

keywords: Egretta thula

## editors

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2014

## authors

authors: BirdLife International

authors: NatureServe

title: Bird Species Distribution Maps of the World. 2012. Egretta thula. The IUCN Red List of Threatened Species.

Version 2014.3  
source:  
type: miscellany  
tags: Egretta thula  
abstract:

## identifiers

last\_modified:  
created:  
group\_id:  
profile\_id:  
id:

taxonRecordName

## ancillaryData

providerSpecificData

anyTwo:  
providerLink:

publicationStatus

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

replacementNameFor

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

sanctioned

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:



laterHomonymOf

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

[conservedAgainst](#)

basedOn

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

basionym

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

[spellingCorrentionOf](#)

typeNameEntity

source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:

typificacion

typeOfType:

[lectotypeMicroReferenceVoucher](#)

lectotypePublicationVoucher

voucherReference

simple:  
microReference:  
year:  
  
publishedIn  
  
simple:  
source:  
datatype:  
identifier:  
  
specialAuthorship

authors

year

simple:  
canonicalAuthorship

authors

year

simple:  
simple: (Molina, 1782)  
  
canonicalName  
  
infraspecificEpithet:  
specificEpithet:  
infragenericEpithet:  
linkType:  
ref:  
uninomial:  
simple: Egretta thula  
rank: SPECIES  
simple: Egretta thula (Molina, 1782)  
  
attributes  
  
nomenclaturalCode:  
isAnamorphic:  
id:

synonymsAtomized

ancillaryData

synonymStatus:  
synonymName  
  
anyTwo:

anyOne

providerLink:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:

ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:

typeNameEntity

source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:

typeVoucherEntity

typeOfType:  
simple:  
microReference:  
year:  
source:  
datatype:  
identifier:

combinationAuthorship

basionymAuthorship

simple:  
authorship

simple:  
simple: Molina, 1782

epithet

infraspecificEpithet:  
specificEpithet:  
infragenericEpithet:

genus

linkType:  
ref:  
uninomial:  
simple: Ardea thula  
rank: SPECIES  
simple: Ardea thula Molina, 1782  
nomenclaturalCode:  
isAnamorphic:  
id:

[hierarchy](#)

[ancillaryData](#)

dataObject:  
additionalInformation:

[reference](#)

geoPoint:  
location:  
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:  
rightsHolder:  
rights:  
license:  
title:  
modified: 2016-02-19T02:26:12.342Z  
created: 2016-02-19T02:26:12.342Z

[agent](#)

homepage:  
role:  
email:  
phone:  
address:  
position:

organisation:  
lastName:  
firstName:  
mimeType:  
dataType:  
identifier:  
parentTaxon: Egretta  
higherClassification:  
infraspecificEpithet:  
specificEpithet:  
taxonRank: SPECIES  
subGenus:  
genus: Egretta  
family: Ardeidae  
order: Pelecaniformes  
classHierarchy: Aves  
phylum: Chordata  
kingdom: Animalia  
recommended:  
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:  
additionalInformation:

0

link: <http://eol.org/pages/1048667/details>

keywords: Egretta thula

0: Shapiro, Leo

1: Hole, Robert

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

0: EOL

title: Egretta thula

source: Encyclopedia of life

type: miscellany

tags: Egretta thula

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Asociación Bogotana de Ornitología  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:

reproductionUnstructured: La temporada reproductiva inicia a finales de marzo e inicios de abril. El nido es construido por la hembra a menudo en hábitats aislados, estuarinos y pueden estar ubicados en el suelo o muy alto en los árboles; los nidos se componen de ramas de tejidas y pequeños palos, también reutilizan nidos; anidan en colonias mixtas (EOL, n.d.). La hembra pone entre 4 a 5 huevos azul verdoso pálido, la incubación es realizada por ambos sexos y dura 24 días en promedio, los polluelos abandonan el nido por lo general a los 14 días después de la eclosión; los juveniles alcanzan la madurez reproductiva después de los 2 años (EOL, n.d.).

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:  
additionalInformation:

0

link: <http://eol.org/pages/1048667/details>

keywords: Egretta thula

0: Shapiro, Leo

1: Hole, Robert

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

0: EOL  
title: Egretta thula  
source: Encyclopedia of life  
type: miscellany  
tags: Egretta thula  
abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Asociación Bogotana de Ornitología  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
feedingUnstructured: Es una especie carnívora; consume lombrices, gusanos, anélidos, insectos acuáticos y terrestres, cangrejos, camarones, ranas, sapos, lagartijas y serpientes (EOL, n.d.).  
  
environmentalEnvelope  
environmentalEnvelopeUnstructured:  
  
ecologicalSignificance  
ecologicalSignificanceUnstructured:  
  
dispersal  
dispersalUnstructured:  
distance:  
structureDispersed:  
type:  
purpose:  
  
behavior  
  
0  
dataObject:  
additionalInformation:  
  
0  
isbn: 0963699636  
keywords: Anas clypeata, Anas platyrhynchos, Ardea herodias, Astragalinus psaltria, Bartramia longicauda, Buteo swainsoni, Cairina moschata, Calidris melanotos, Cardellina canadensis, Cathartes aura, Catharus minimus, Coccyzus americanus, Columba livia, Columbina talpacoti, Contopus cooperi, Contopus virens, Coragyps atratus, Crotophaga ani, Dendrocygna autumnalis, Egretta caerulea, Egretta thula, Empidonax alnorum, Empidonax traillii, Falco columbarius, Fulica americana, Geothlypis philadelphia, Hirundo rustica, Ixobrychus exilis, Jacana jacana, Mnioilta



varia, Nycticorax nycticorax, Oxyura jamaicensis, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Phalacrocorax brasilianus, Piranga olivacea, Pitangus sulphuratus, Podilymbus podiceps, Porzana carolina, Pyrocephalus rubinus, Rupornis magnirostris, Setophaga fusca, Sturnella magna, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon

0: Alderfer, Jonathan

institution:

address:

publisher: Washington D.C. National Geographic Society

accessed:

websites:

chapter:

series:

pages: 664

issue:

year: 2006

0: Alderfer, J

1: Zickefoose, J

2: Littlehales, B

title: National Geographic Complete Birds of North America

source:

type: book

tags: Anas clypeata, Anas platyrhynchos, Ardea herodias, Astragalinus psaltria, Bartramia longicauda, Buteo swainsoni, Cairina moschata, Calidris melanotos, Cardellina canadensis, Cathartes aura, Catharus minimus, Coccyzus americanus, Columba livia, Columbina talpacoti, Contopus cooperi, Contopus virens, Coragyps atratus, Crotophaga ani, Dendrocygna autumnalis, Egretta caerulea, Egretta thula, Empidonax alnorum, Empidonax traillii, Falco columbarius, Fulica americana, Geothlypis philadelphia, Hirundo rustica, Ixobrychus exilis, Jacana jacana, Mniotilta varia, Nycticorax nycticorax, Oxyura jamaicensis, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Phalacrocorax brasilianus, Piranga olivacea, Pitangus sulphuratus, Podilymbus podiceps, Porzana carolina, Pyrocephalus rubinus, Rupornis magnirostris, Setophaga fusca, Sturnella magna, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

1

keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Agelaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccyzus pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta caerulea, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria

squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerinii, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicrodon microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

pages: 836

issue:

year: 1986

0: Hilty, Steven

1: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccyzus pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta caerulea, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus

superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerinii, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

2

link: <http://eol.org/pages/1048667/details>

keywords: Egretta thula

0: Shapiro, Leo

1: Hole, Robert

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

0: EOL

title: Egretta thula

source: Encyclopedia of life

type: miscellany

tags: Egretta thula

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

3

isbn: 978-0-679-45122-8

keywords: *Anas clypeata*, *Ardea herodias*, *Buteo swainsoni*, *Cairina moschata*, *Crotophaga ani*, *Dendrocygna autumnalis*, *Dendrocygna viduata*, *Egretta caerulea*, *Egretta thula*, *Empidonax alnorum*, *Geothlypis philadelphia*, *Ixobrychus exilis*, *Leiothlypis peregrina*, *Myiodynastes luteiventris*, *Nomonyx dominicus*, *Parkesia noveboracensis*, *Petrochelidon pyrrhonota*, *Pheucticus ludovicianus*, *Protonotaria citrea*, *Sayornis nigricans*

institution:

edition: 1 st ed.

address:

publisher: National Audubon Society

accessed:

websites:

chapter:

series:

pages: 545

issue:

year: 2000

0: Sibley, D A

title: The SIBLEY Guide to Birds

source:

type: book

tags: *Anas clypeata*, *Ardea herodias*, *Buteo swainsoni*, *Cairina moschata*, *Crotophaga ani*, *Dendrocygna autumnalis*, *Dendrocygna viduata*, *Egretta caerulea*, *Egretta thula*, *Empidonax alnorum*, *Geothlypis philadelphia*, *Ixobrychus exilis*, *Leiothlypis peregrina*, *Myiodynastes luteiventris*, *Nomonyx dominicus*, *Parkesia noveboracensis*, *Petrochelidon pyrrhonota*, *Pheucticus ludovicianus*, *Protonotaria citrea*, *Sayornis nigricans*

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

behaviorUnstructured: La Garza Patiamarilla tiene una amplia gama de comportamientos de forrajeo en comparación con las otras especies de garzas; captura el alimento buscando activamente en el agua, removiendo con las patas a medida que camina, también lo hace saltando, corriendo, flotando y persiguiendo a su presa. A menudo se alimentan en horas de la mañana y de la tarde (Alderfer et al., 2006; Hilty & Brown, 1986). Se alimentan en grupos grandes ya que les permite un mayor éxito en la búsqueda de fuentes de alimento y a protegerse de los depredadores (EOL, n.d.). En la época reproductiva el macho realiza exhibiciones, sobrevolando y vocalizando para atraer a las hembras; el macho corteja a la hembra realizando movimientos con su cuerpo de arriba hacia abajo y proyectando su pico hacia el cielo y vocaliza para atraerla; la coloración de las patas cambia de amarillo a naranja rojizo indicando el comienzo de la reproducción; una vez el macho encuentre su pareja viene la cópula y la construcción del nido que es realizado por la hembra y el macho lo protege (Alderfer et al., 2006; EOL, n.d.).

Es una especie generalmente silenciosa, pero cuando está en temporada reproductiva vocaliza un áspero ahhhh o

un uah uah (Hilty & Brown, 1986). Un reclamo similar al de la Garza azul (*Egretta caerulea*) un nasal raaaaaaa pero más áspero y alto; en vuelo ocasionalmente hace un tosido carrrrf (Sibley, 2000).

## ancillaryData

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights:  
license:  
modified:  
created:  
mimeType:  
dataType:  
identifier:

## measurementOrFact

relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

## endemicAtomized

## distribution

## ancillaryData

dataObject:  
additionalInformation:

## reference

link: [http://awsassets.panda.org/downloads/migratorias{\\\_}aves{\\\_}42{\\\_}final.pdf](http://awsassets.panda.org/downloads/migratorias{\_}aves{\_}42{\_}final.pdf)  
keywords: *Egretta thula*

## editors

editors: Naranjo, L G  
editors: Amaya, J D  
editors: Eusse-González, D  
editors: Cifuentes-Sarmiento, Y  
institution:  
address:  
publisher: Ministerio de Ambiente y Desarrollo Sostenible / WWF Colombia  
accessed:

websites:  
chapter:  
series:  
pages: 708  
issue:  
volume: 1  
year: 2012

## authors

authors: Ruíz-Guerra, Carlos  
title: Egretta thula  
source: Guía de las Especies Migratorias de la Biodiversidad en Colombia Aves  
type: book\_section  
tags: Egretta thula  
abstract:

## identifiers

last\_modified:  
created:  
group\_id:  
profile\_id:  
id:  
geoPoint:  
location:  
thumbnailURL:

## mediaURL

description:

## subject

source:

## audience

bibliographicCitation:  
rightsHolder:  
rights: Asociación Bogotana de Ornitología  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

## agent

mimeType:  
dataType:  
identifier:  
distributionUnstructured: La Garza Patiámarilla se distribuye en Canadá, Estados Unidos, Antillas y Sudamérica. Las poblaciones de Norteamérica migran hasta Sudamérica, transeúntes desde Norteamérica han sido registrados en Guyana, Venezuela y Colombia; pero la distribución de los migratorios en Sudamérica es aún desconocida (Ruíz-Guerra, 2012).  
Hasta unos 1000 msnm; ocasionalmente 2600 msnm (Sabana de Bogotá) (Hilty & Brown, 1986).

## distributionAtomized

stateProvince: Córdoba

## ancillaryData

locality:  
municipality:  
country: Colombia  
temporalCoverage  
endDate:  
startDate:  
distributionScope  
type:

## threatStatus

territory  
territoryUnstructured:  
areaOfOccupancy:  
extentOfOccurrence:  
populationBiology

0

dataObject:  
additionalInformation:

0

link: <http://maps.iucnredlist.org/map.html?id=22696974>

keywords: Egretta thula

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2014

0: BirdLife International

1: NatureServe

title: Bird Species Distribution Maps of the World. 2012. Egretta thula. The IUCN Red List of Threatened Species.

Version 2014.3

source:

type: miscellany

tags: Egretta thula

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Asociación Bogotana de Ornitología  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
populationBiologyUnstructured: Es una especie con un amplio rango de distribución y se considera como de Preocupación Menor según la IUCN (BirdLife International and NatureServe, 2014).

## legislation

legislationUnstructured:

## directThreats

directThreatsUnstructured:

## ancillaryData

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights:  
license:  
modified:  
created:  
mimeType:  
dataType:  
identifier:

## measurementOrFact

relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

abstract: La Garza Patiamarilla mide 61 cm aproximadamente; es totalmente blanca, sus patas y pico son negros y los dedos amarillo brillante. En plumaje nupcial presenta plumas largas, recurvadas, llamadas airones, en la coronilla el pecho y la espalda (Alcaldía de Santiago de Cali Departamento Administrativo de Gestión del Medio Ambiente, 2010). Se alimenta de pequeñas presas que captura removiendo el agua con sus patas a medida que camina (Alcaldía de Santiago de Cali Departamento Administrativo de Gestión del Medio Ambiente, 2010). Generalmente es silenciosa, pero cuando está en temporada reproductiva vocaliza un áspero ahhhh o un uah uah (Alcaldía de Santiago de Cali Departamento Administrativo de Gestión del Medio Ambiente, 2010; Hilty & Brown,



1986). Es una especie bastante común en el Valle del Cauca, sobre todo asociada a humedales y ríos de poca profundidad (Alcaldía de Santiago de Cali Departamento Administrativo de Gestión del Medio Ambiente, 2010), a manglares, estuarios y playones costeros de lodo (Hilty & Brown, 1986). Es una especie gregaria y suele formar bandadas mixtas con otras especies de garzas y aves acuáticas (zancudas) (Alcaldía de Santiago de Cali Departamento Administrativo de Gestión del Medio Ambiente, 2010; McNish, 2007). Es una especie con un amplio rango de distribución y se considera de Preocupación Menor según la IUCN (BirdLife International and NatureServe, 2014).

## associatedParty

role:  
personnelIdentifier:  
personnelDirectory:  
homepage:  
email: enataliagomezm@gmail.com  
phone:  
postalCode:  
country: Colombia  
state: Bogotá Distrito Capital  
city: Bogotá D.C.  
address:  
organisation:  
position:  
lastName: Gómez Melendro  
firstName: Edith Natalia

## ancillaryData

dataObject:  
additionalInformation:

## reference

geoPoint:  
location:  
thumbnailURL:  
mediaURL: <http://avibase.bsc-eoc.org/species.jsp?lang=EN&avibaseid=AA7901D88E7203E5>  
description:

## subject

source: <http://avibase.bsc-eoc.org>

## audience

bibliographicCitation:  
rightsHolder:  
rights:  
license:  
modified:  
created:

## agent

mimeType:  
dataType:  
identifier:

\_id

\_bsontype: ObjectId  
id: VÆ~Eð lg#ç<°