

## Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56d6ebf83c16479905cba913

Fecha de creación: 2016-03-02T13:34:51.006Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

habitatUnstructured: Poco común como transeúnte de otoño y común como residente de invierno y transeúnte de primavera en monte secundario, bordes y montes claros; áreas secas a húmedas (Hilty & Brown 1986). En La Guajira habita en el bosque deciduo tropical, de galería y secundario (Cerrejón 2009-2010).

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

[ancillaryData](#)

type:

language: Francés

name: Cardinal à poitrine rose

creation\_date: 2016-03-02T13:34:51.006Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

fullDescriptionUnstructured:  
briefDescription:

references

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:  
providerLink:

publicationStatus

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

replacementNameFor

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

sanctioned

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

laterHomonymOf

microReference:  
source:  
datatype:  
identifier:  
source:

datatype:  
identifier:  
note:  
ruleConsidered:

[conservedAgainst](#)

basedOn

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

basionym

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

[spellingCorrentionOf](#)

typeNameEntity

source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:

typificacion

typeOfType:

[lectotypeMicroReferenceVoucher](#)

[lectotypePublicationVoucher](#)

[voucherReference](#)

simple:  
microReference:  
year:

publishedIn

simple:  
source:  
datatype:  
identifier:

specialAuthorship

authors

year

simple:  
canonicalAuthorship

authors

year

simple:  
simple: (Linnaeus, 1766)  
canonicalName  
infraspecificEpithet:  
specificEpithet:  
infragenericEpithet:  
linkType:  
ref:  
uninomial:  
simple: Pheucticus ludovicianus  
rank: SPECIES  
simple: Pheucticus ludovicianus (Linnaeus, 1766)

attributes

nomenclaturalCode:  
isAnamorphic:  
id:

synonymsAtomized

ancillaryData

synonymStatus:  
synonymName

anyTwo:

anyOne

providerLink:  
microReference:

publishedIn

source:

datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:

typeNameEntity

source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:

typeVoucherEntity

typeOfType:  
simple:  
microReference:  
year:  
source:  
datatype:  
identifier:

combinationAuthorship

basionymAuthorship

simple:  
authorship  
simple:  
simple: Linnaeus, 1766

epithet

infraspecificEpithet:  
specificEpithet:  
infragenericEpithet:

genus

linkType:

ref:

uninomial:

simple: *Loxia ludovicianus*

rank: SPECIES

simple: *Loxia ludovicianus* Linnaeus, 1766

nomenclaturalCode:

isAnamorphic:

id:

## hierarchy

## ancillaryData

dataObject:

additionalInformation:

## reference

geoPoint:

location:

thumbnailURL:

## mediaURL

description:

## subject

source: The Catalogue of Life, 3rd January 2011

## audience

bibliographicCitation:

rightsHolder:

rights:

license:

title:

modified: 2016-03-02T13:34:31.002Z

created: 2016-03-02T13:34:31.002Z

## agent

homepage:

role:

email:

phone:

address:

position:

organisation:

lastName:

firstName:

mimeType:

dataType:

identifier:

parentTaxon: *Pheucticus*

higherClassification:

infraspecificEpithet:  
specificEpithet:  
taxonRank: SPECIES  
subGenus:  
genus: Pheucticus  
family: Cardinalidae  
order: Passeriformes  
classHierarchy: Aves  
phylum: Chordata  
kingdom: Animalia  
recommended:  
classification:

invasiveness

invasivenessUnstructured:

reproduction

reproductionUnstructured:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Equipo Coordinador del SiB Colombia  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
feedingUnstructured: En La Guajira forrajea en el dosel (Cerrejón 2009-2010) y su dieta se compone de granos



(Cerrejón 2009-2010).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

behaviorUnstructured: Llamado quic agudo y mrtálico (Hilty & Brown 1986).

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

[endemicAtomized](#)

[distribution](#)

[ancillaryData](#)

dataObject:  
additionalInformation:

[reference](#)

geoPoint:  
location:  
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:  
rightsHolder:  
rights: Equipo Coordinador del SiB Colombia  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

[agent](#)

contentType:  
dataType:  
identifier:  
distributionUnstructured: En bases de datos se encontró que: Esta especie se distribuye en: Colombia (Arctos 2012; Collections MNH SI 2012); Canadá, México, El Salvador, Estado Unidos, Guatemala, Honduras (Arctos 2012). En bases de datos, literatura y monitoreos de fauna se encontró que: Esta especie se distribuye en: Antioquia, Bogotá, Bolívar, Boyacá, Chocó, Cundinamarca, La Guajira, Magdalena, Santander, Tolima, Valle del Cauca (ICN 2004); Antioquia, Boyacá, Cesar, La Guajira, Magdalena, Norte de Santander (Collections MNH SI 2012); Boyacá, Huila, Santander, Valle del Cauca (Arctos 2012). Se encuentra en las regiones Andina, Caribe y Pacífica hasta 3800 m (Hilty & Brown 1986).

[distributionAtomized](#)

temporalCoverage

endDate:  
startDate:

distributionScope

type:

threatStatus

ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:

locality:

municipality:

county:

stateProvince:

country: Global

0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementType: IUCN

measurementID:

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:  
rightsHolder:  
rights:  
license:  
modified:  
created:  
mimeType:  
dataType:  
identifier:

## measurementOrFact

relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

abstract: Mide 19 cm aproximadamente. Tiene un pico robusto de color blanco yeso. Los machos cambian de plumaje antes de la migración para la época de reproducción: las partes dorsales, la cabeza y garganta de color negro con dos parches alares extensos blancos, la rabadilla y la cola blancas; las partes ventrales blancas con un collar rosa que se extiende hasta la mitad del pecho. En la época no reproductiva el plumaje es similar pero moteado parduzco dorsalemente; la cabeza con listas ante; las partes inferiores moteadas de marrón y con algunas trazas de rosa. Las hembras son de color marrón en las partes dorsales, estriado de negruzco; con una lista estrecha en la coronilla de color blanco anteado; mejillas marrón; las partes ventrales son blanco anteado con finas estrías marrón (Hilty & Brown 1986).

## associatedParty

role:  
personnelIdentifier:  
personnelDirectory:  
homepage:  
email: enataliagomezm@gmail.com  
phone:  
postalCode:  
country: Colombia  
state: Bogotá Distrito Capital  
city: Bogotá D.C.  
address:  
organisation:  
position:  
lastName: Gómez Melendro  
firstName: Edith Natalia

## ancillaryData

dataObject:  
additionalInformation: Registros biológicos: En el portal de Global Biodiversity Information Facility se reporta que para Colombia hay 453 registros. GBIF.  
Créditos específicos: Ficha elaborada con el apoyo de Carbones del Cerrejón Limited.  
Otros recursos en internet: Arctos, Avibase, AVoCet, BirdLife International, Colecciones en Línea, EoL, iSpecies, IUCN Red List, SI NMNH, xeno-canto

## reference

geoPoint:  
location:  
thumbnailURL:

[mediaURL](#)

description:  
[subject](#)

source:  
[audience](#)

bibliographicCitation:  
rightsHolder:  
rights: Equipo Coordinador del SiB Colombia  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

[agent](#)

mimeType:  
dataType:  
identifier:

\_id  
\_bsontype: ObjectID  
id: VÖëø< G™\°°0