

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56c229d1f0106c67230e7323

Fecha de creación: undefined

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

0

keywords: Actitis macularius,Ampelion rubrocristatus,Anas discors,Anisognathus igniventris,Ardea alba,Arremon assimilis,Atlapetes pallidinucha,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: Actitis macularius,Ampelion rubrocristatus,Anas discors,Anisognathus igniventris,Ardea alba,Arremon assimilis,Atlapetes pallidinucha,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata

abstract:

last_modified:

created:

group_id:

profile_id:

id:

1

keywords: Accipiter striatus,Actitis macularius,Adelomyia melanogenys,Aglaiocercus kingii,Amblycercus holosericeus,Ampelion rubrocristatus,Anas andium,Anas clypeata,Anas discors,Andigena nigrirostris,Anisognathus igniventris,Ardea alba,Ardea herodias,Arremon assimilis,Asio flammeus,Asio stygius,Asthenes flammulata,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bubulcus ibis,Buteo platypterus,Buteo

swainsoni,Butorides striata,Butorides virescens,Cairina moschata,Calidris bairdii,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Cathartes aura,Catharus minimus,Catharus ustulatus,Chaetocercus mulsant,Chaetura pelagica,Chalcostigma heteropogon,Chordeiles minor,Chrysomus icterocephalus,Cinclodes albidiventris,Cinclus leucocephalus,Cinnycerthia unirufa,Cistothorus platensis,Coccyzus pumila,Coccyzus americanus,Coccyzus melacoryphus,Coeligena bonapartei,Colibri coruscans,Colibri thalassinus,Columba livia,Columbina talpacoti,Conirostrum rufum,Conirostrum sitticolor,Contopus cinereus,Contopus fumigatus,Contopus sordidulus,Contopus virens,Coragyps atratus,Crotophaga ani,Crotophaga major,Dendrocygna autumnalis,Dendrocygna viduata,Diglossa albilatera,Diglossa cyanea,Diglossa humeralis,Diglossa lafresnayii,Diglossa sittoides,Dolichonyx oryzivorus,Dubusia taeniata,Egretta thula,Elaenia flavogaster,Elaenia frantzii,Elanus leucurus,Empidonax alnorum,Empidonax traillii,Ensifera ensifera,Eriocnemis cupreoventris,Eriocnemis vestita,Eupsittula pertinax,Falco columbarius,Falco peregrinus,Falco sparverius,Forpus conspicillatus,Fulica americana,Gallinago nobilis,Gallinula galeata,Geothlypis philadelphia,Geranoaetus melanoleucus,Glaucidium jardinii,Grallaria quitesis,Grallaria ruficapilla,Grallaria squamigera,Haplospiza rustica,Heliangelus amethysticollis,Heliangelus exortis,Hellmayrea gularis,Hemispingus supercilialis,Hemispingus verticalis,Henicorhina leucophrys,Hirundo rustica,Icterus chrysater,Icterus icterus,Icterus nigrogularis,Ixobrychus exilis,Jacana jacana,Lafresnaya lafresnayi,Leiostyris alpestris,Leptasthenura andicola,Lesbia nuna,Lesbia victoriae,Machetornis rixosa,Margarornis squamiger,Mecocerculus leucophrys,Megascops choliba,Melanerpes rubricapillus,Merganetta armata,Metallura tyrianthina,Milvago chimachima,Mimus gilvus,Mniotilta varia,Molothrus bonariensis,Muscisaxicola alpinus,Myiarchus crinitus,Myiodynastes luteiventris,Myiotheretes fumigatus,Myiotheretes striaticollis,Myiothlypis nigrocristata,Nomonyx dominicus,Nycticorax nycticorax,Ochthoeca cinnamomeiventris,Ochthoeca diadema,Ochthoeca fumicolor,Ochthoeca rufipectoralis,Orochelidon murina,Oxyechus vociferus,Oxyura jamaicensis,Pachyrhamphus versicolor,Pandion haliaetus,Pardaliparus maculatus,Parkesia noveboracensis,Patagioenas fasciata,Petrochelidon pyrrhonota,Phaetusa simplex,Phalacrocorax brasilianus,Pheucticus aureoventris,Pheucticus ludovicianus,Phimosus infuscatus,Phrygilus unicolor,Phyllomyias uropygialis,Piaya cayana,Picoides fumigatus,Pipraeidea melanonota,Pipreola riefferii,Piranga olivacea,Piranga rubra,Pitangus sulphuratus,Pluvialis dominica,Podilymbus podiceps,Porphyrion martinicus,Porzana carolina,Progne subis,Protonotaria citrea,Pseudocolaptes auratus,Pseudocolaptes auratus,Pseudoscops clamator,Pterophanes cyanopterus,Pygochelidon cyanoleuca,Pyrocephalus rubinus,Pyrrhomyias cinnamomeus,Quiscalus lugubris,Rallus semiplumbeus,Ramphocelus dimidiatus,Ramphomicrodon microhynchus,Riparia riparia,Rupornis magnirostris,Sayornis nigricans,Serpophaga cinerea,Setophaga castanea,Setophaga cerulea,Setophaga petechia,Setophaga ruticilla,Setophaga striata,Sicalis flaveola,Sicalis luteola,Spiza americana,Sporagra spinescens,Sturnella magna,Synallaxis subpudica,Systellura longirostris,Tangara vitriolina,Thraupis episcopus,Thraupis palmarum,Tringa flavipes,Tringa melanoleuca,Tringa solitaria,Troglodytes aedon,Troglodytes solstitialis,Turdus fuscater,Turdus ignobilis,Tyrannus melancholicus,Tyrannus savana,Tyrannus tyrannus,Tyto alba,Vanellus chilensis,Vireo olivaceus,Vultur gryphus,Zenaidura macroura,Zonotrichia capensis

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

pages: 836

issue:

year: 1986

0: Hilty, Steven

1: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

tags: Accipiter striatus,Actitis macularia,Adelomyia melanogenys,Aglaiocercus kingii,Amblycercus holosericeus,Ampelis rubrocrissalis,Anas andium,Anas clypeata,Anas discors,Andigena nigrirostris,Anisognathus igniventris,Ardea alba,Ardea herodias,Arremon assimilis,Asio flammeus,Asio stygius,Asthenes flammulata,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Butorides virescens,Cairina moschata,Calidris bairdii,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Cathartes aura,Catharus minimus,Catharus

ustulatus,Chaetocercus mulsant,Chaetura pelagica,Chalcostigma heteropogon,Chordeiles minor,Chrysomus icterocephalus,Cinclodes albidiventris,Cinclus leucocephalus,Cinnycerthia unirufa,Cistothorus platensis,Coccyzua pumila,Coccyzus americanus,Coccyzus melacoryphus,Coeligena bonapartei,Colibri coruscans,Colibri thalassinus,Columba livia,Columbina talpacoti,Conirostrum rufum,Conirostrum sitticolor,Contopus cinereus,Contopus fumigatus,Contopus sordidulus,Contopus virens,Coragyps atratus,Crotophaga ani,Crotophaga major,Dendrocygna autumnalis,Dendrocygna viduata,Diglossa albilatera,Diglossa cyanea,Diglossa humeralis,Diglossa lafresnayii,Diglossa sittoides,Dolichonyx oryzivorus,Dubusia taeniata,Egretta thula,Elaenia flavogaster,Elaenia frantzii,Elanus leucurus,Empidonax alnorum,Empidonax traillii,Ensifera ensifera,Eriocnemis cupreoventris,Eriocnemis vestita,Eupsittula pertinax,Falco columbarius,Falco peregrinus,Falco sparverius,Forpus conspicillatus,Fulica americana,Gallinago nobilis,Gallinula galeata,Geothlypis philadelphia,Geranoaetus melanoleucus,Glaucidium jardinii,Grallaria quitesis,Grallaria ruficapilla,Grallaria squamigera,Haplospiza rustica,Heliangelus amethysticollis,Heliangelus exortis,Hellmayrea gularis,Hemispingus superciliaris,Hemispingus verticalis,Henicorhina leucophrys,Hirundo rustica,Icterus chrysater,Icterus icterus,Icterus nigrogularis,Ixobrychus exilis,Jacana jacana,Lafresnaya lafresnayi,Leiostyris alpestris,Leptasthenura andicola,Lesbia nuna,Lesbia victoriae,Machetornis rixosa,Margarornis squamiger,Mecocerculus leucophrys,Megascops choliba,Melanerpes rubricapillus,Merganetta armata,Metallura tyrianthina,Milvago chimachima,Mimus gilvus,Mniotilta varia,Molothrus bonariensis,Muscisaxicola alpinus,Myiarchus cinerascens,Myiodynastes luteiventris,Myiotheretes fumigatus,Myiotheretes striatocollis,Myiothlypis nigrocristata,Nomonyx dominicus,Nycticorax nycticorax,Ochthoeca cinnamomeiventris,Ochthoeca diadema,Ochthoeca fumicolor,Ochthoeca rufipectoralis,Orochelidon murina,Oxyechus vociferus,Oxyura jamaicensis,Pachyrhamphus versicolor,Pandion haliaetus,Pardaliparus maculatus,Parkesia noveboracensis,Patagioenas fasciata,Petrochelidon pyrrhonota,Phaetusa simplex,Phalacrocorax brasilianus,Pheucticus aureoventris,Pheucticus ludovicianus,Phimosus infuscatus,Phrygilus unicolor,Phyllomyias uropygialis,Piaya cayana,Picoides fumigatus,Pipraeidea melanonota,Pipreola riefferii,Piranga olivacea,Piranga rubra,Pitangus sulphuratus,Pluvialis dominica,Podilymbus podiceps,Porphyrion martinicus,Porzana carolina,Progne subis,Protonotaria citrea,Pseudocolaptes auratus,Pseudocolaptes cafer,Pseudoscops clamator,Pterophanes cyanopterus,Pygochelidon cyanoleuca,Pyrocephalus rubinus,Pyrrhomyias cinnamomeus,Quiscalus lugubris,Rallus semiplumbeus,Ramphocelus dimidiatus,Ramphomicron microrhynchum,Riparia riparia,Rupornis magnirostris,Sayornis nigricans,Serpophaga cinerea,Setophaga castanea,Setophaga cerulea,Setophaga petechia,Setophaga ruticilla,Setophaga striata,Sicalis flaveola,Sicalis luteola,Spiza americana,Sporagra spinescens,Sturnella magna,Synallaxis subpudica,Systellura longirostris,Tangara vitriolina,Thraupis episcopus,Thraupis palmarum,Tringa flavipes,Tringa melanoleuca,Tringa solitaria,Troglodytes aedon,Troglodytes solstitialis,Turdus fuscater,Turdus ignobilis,Tyrannus melancholicus,Tyrannus savana,Tyrannus tyrannus,Tyto alba,Vanellus chilensis,Vireo olivaceus,Vultur gryphus,Zenaidura macroura,Zonotrichia capensis

abstract:

last_modified:

created:

group_id:

profile_id:

id:

2

link: <http://www.hbw.com/node/61987>

keywords: Atlapetes pallidinucha

0: del Hoyo, J

1: Elliott, A

2: Sargatal, J

3: Christie, D A

4: de Juana, E

institution:

address:

publisher: Lynx Edicions

accessed:

websites:

chapter:

series:

issue:
year: 2011
0: Jaramillo, A
title: Pale-naped Brush-finch (Atlapetes pallidinucha)
source: Handbook of the Birds of the World Alive
type: book_section
tags: Atlapetes pallidinucha
abstract:
last_modified:
created:
group_id:
profile_id:
id:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
habitatUnstructured: La especie habita desde la zona subtropical hasta el páramo (Restall, Rodner, & Lentino, 2006). Es común en bordes de bosque, bosque enano, matorrales hasta el límite de vegetación arbórea y densos chuscales (Chusquea) (Asociación Bogotana de Ornitología, 2000; Hilty & Brown, 1986; Jaramillo, 2011).

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:
usedIn
distributionUnstructured:
endDate:
startDate:

[ancillaryData](#)

type:
language: Español
name: Atlapetes Cabeciblanco

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

0

keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata

abstract:

last_modified:

created:

group_id:

profile_id:

id:

1

keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri

major, *Dendrocygna autumnalis*, *Dendrocygna viduata*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dolichonyx oryzivorus*, *Dubusia taeniata*, *Egretta thula*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreovertris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Geothlypis philadelphia*, *Geranoaetus melanoleucus*, *Glaucidium jardinii*, *Grallaria quitesis*, *Grallaria ruficapilla*, *Grallaria squamigera*, *Haplospiza rustica*, *Heliangelus amethysticollis*, *Heliangelus exortis*, *Hellmayrea gularis*, *Hemispingus superciliaris*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus icterus*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Lafresnaya lafresnayi*, *Leiothlypis peregrina*, *Leptasthenura andicola*, *Lesbia nuna*, *Lesbia victoriae*, *Machetornis rixosa*, *Margarornis squamiger*, *Mecocerculus leucophrys*, *Megascops choliba*, *Melanerpes rubricapillus*, *Merganetta armata*, *Metallura tyrianthina*, *Milvago chimachima*, *Mimus gilvus*, *Mniotilta varia*, *Molothrus bonariensis*, *Muscisaxicola alpinus*, *Myiarchus crinitus*, *Myiodynastes luteiventris*, *Myiotheretes fumigatus*, *Myiotheretes striaticollis*, *Myiothlypis nigrocristata*, *Nomonyx dominicus*, *Nycticorax nycticorax*, *Ochthoeca cinnamomeiventris*, *Ochthoeca diadema*, *Ochthoeca fumicolor*, *Ochthoeca rufipectoralis*, *Orochelidon murina*, *Oxyopogon guerinii*, *Oxyura jamaicensis*, *Pachyrhamphus versicolor*, *Pandion haliaetus*, *Pardirallus maculatus*, *Parkesia noveboracensis*, *Patagioenas fasciata*, *Petrochelidon pyrrhonota*, *Phaetusa simplex*, *Phalacrocorax brasilianus*, *Pheucticus aureoventris*, *Pheucticus ludovicianus*, *Phimosus infuscatus*, *Phrygilus unicolor*, *Phyllomyias uropygialis*, *Piaya cayana*, *Picoides fumigatus*, *Pipraeidea melanonota*, *Pipreola riefferii*, *Piranga olivacea*, *Piranga rubra*, *Pitangus sulphuratus*, *Pluvialis dominica*, *Podilymbus podiceps*, *Porphyrio martinicus*, *Porzana carolina*, *Progne tapera*, *Protonotaria citrea*, *Pseudocolopteryx acutipennis*, *Pseudoscops clamator*, *Pterophanes cyanopterus*, *Pygochelidon cyanoleuca*, *Pyrocephalus rubinus*, *Pyrrhomyias cinnamomeus*, *Quiscalus lugubris*, *Rallus semiplumbeus*, *Ramphocelus dimidiatus*, *Ramphomicron microrhynchum*, *Riparia riparia*, *Rupornis magnirostris*, *Sayornis nigricans*, *Serpophaga cinerea*, *Setophaga castanea*, *Setophaga cerulea*, *Setophaga petechia*, *Setophaga ruticilla*, *Setophaga striata*, *Sicalis flaveola*, *Sicalis luteola*, *Spiza americana*, *Sporagra spinescens*, *Sturnella magna*, *Synallaxis subpudica*, *Systellura longirostris*, *Tangara vitriolina*, *Thraupis episcopus*, *Thraupis palmarum*, *Tringa flavipes*, *Tringa melanoleuca*, *Tringa solitaria*, *Troglodytes aedon*, *Troglodytes solstitialis*, *Turdus fuscater*, *Turdus ignobilis*, *Tyrannus melancholicus*, *Tyrannus savana*, *Tyrannus tyrannus*, *Tyto alba*, *Vanellus chilensis*, *Vireo olivaceus*, *Vultur gryphus*, *Zenaida auriculata*, *Zonotrichia capensis*

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotá de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

fullDescriptionUnstructured: La especie mide aproximadamente 18 cm. Gris teñido de oliva por encima; lados de cabeza y cuello negros (formando una amplia máscara); amplia lista medial en la coronilla naranja canela variando hasta amarillo en la frente desvanecida a blanco en la nuca; garganta amarilla brillante, resto de partes inferiores amarillo oliva opaco; flancos y área perianal oliva (Asociación Bogotá de Ornitología, 2000; Hilty & Brown, 1986).

briefDescription:

[references](#)

keywords: Accipiter striatus, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Anas andium, Andigena nigrirostris, Anisognathus igniventris, Asthenes flammulata, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Calidris bairdii, Chalcostigma heteropogon, Chordeiles minor, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coeligena bonapartei, Colibri thalassinus, Coragyps atratus, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa sittoides, Eriocnemis vestita, Falco peregrinus, Gallinago nobilis, Gallinula galeata, Geranoaetus melanoleucus, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Mecocerculus leucophrys, Mniotilta varia, Molothrus bonariensis, Picoides fumigatus, Piranga olivacea, Pyrocephalus rubinus, Rupornis magnirostris, Sicalis flaveola, Sporagra spinescens, Sturnella magna, Tringa solitaria, Zonotrichia capensis

editors

institution:
address:
publisher: Christopher Helm
accessed:
websites:
chapter:
series:
pages: 880
issue:
year: 2006

authors

authors: Restall, R
authors: Rodner, C
authors: Lentino, M
title: Birds of Northern South America: An Identification Guide, Volume 1: Species Accounts.
source:
type: book
tags: Accipiter striatus, Aglaiocercus kingii, Amblycercus holosericeus, Anas andium, Andigena nigrirostris, Anisognathus igniventris, Asthenes flammulata, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Calidris bairdii, Chalcostigma heteropogon, Chordeiles minor, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coeligena bonapartei, Colibri thalassinus, Coragyps atratus, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa sittoides, Eriocnemis vestita, Falco peregrinus, Gallinago nobilis, Gallinula galeata, Geranoaetus melanoleucus, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Mecocerculus leucophrys, Mniotilta varia, Molothrus bonariensis, Picoides fumigatus, Piranga olivacea, Pyrocephalus rubinus, Rupornis magnirostris, Sicalis flaveola, Sporagra spinescens, Sturnella magna, Tringa solitaria, Zonotrichia capensis
abstract:

identifiers

last_modified:
created:
group_id:
profile_id:
id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:
providerLink:

publicationStatus

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

replacementNameFor

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

sanctioned

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

laterHomonymOf

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

[conservedAgainst](#)

basedOn

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:
canonicalAuthorship

authors

year

simple:
simple: (Boissonneau, 1840)

canonicalName

infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:
ref:
uninomial:
simple: Atlapetes pallidinucha
rank: SPECIES
simple: Atlapetes pallidinucha (Boissonneau, 1840)

attributes

nomenclaturalCode:
isAnamorphic:
id:

synonymsAtomized

hierarchy

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source: The Catalogue of Life, 3rd January 2011

audience

bibliographicCitation:
rightsHolder:
rights:
license:
title:
modified: 2016-02-15T19:36:43.529Z
created: 2016-02-15T19:36:43.529Z

agent

homepage:
role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:
parentTaxon: Atlapetes
higherClassification:
infraspecificEpithet:
specificEpithet:
taxonRank: SPECIES
subGenus:
genus: Atlapetes
family: Emberizidae
order: Passeriformes
classHierarchy: Aves
phylum: Chordata
kingdom: Animalia
recommended:
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:
additionalInformation:

0

link: <http://www.hbw.com/node/61987>

keywords: Atlapetes pallidinucha

0: del Hoyo, J

1: Elliott, A

2: Sargatal, J

3: Christie, D A

4: de Juana, E

institution:

address:

publisher: Lynx Edicions

accessed:

websites:

chapter:

series:

issue:

year: 2011

0: Jaramillo, A

title: Pale-naped Brush-finch (Atlapetes pallidinucha)

source: Handbook of the Birds of the World Alive

type: book_section

tags: *Atlapetes pallidinucha*

abstract:

last_modified:

created:

group_id:

profile_id:

id:

1

keywords: *Accipiter striatus*, *Actitis macularius*, *Adelomyia melanogenys*, *Aglaiocercus kingii*, *Amblycercus holosericeus*, *Ampelion rubrocristatus*, *Anas andium*, *Anas clypeata*, *Anas discors*, *Andigena nigrirostris*, *Anisognathus igniventris*, *Ardea alba*, *Ardea herodias*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Asthenes flammulata*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Butorides virescens*, *Cairina moschata*, *Calidris bairdii*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Cathartes aura*, *Catharus minimus*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chaetura pelagica*, *Chalcostigma heteropogon*, *Chordeiles minor*, *Chrysomus icterocephalus*, *Cinclodes albidiventris*, *Cinclus leucocephalus*, *Cinnycerthia unirufa*, *Cistothorus platensis*, *Coccyzus pumila*, *Coccyzus americanus*, *Coccyzus melacoryphus*, *Coeligena bonapartei*, *Colibri coruscans*, *Colibri thalassinus*, *Columba livia*, *Columbina talpacoti*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cinereus*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Dendrocygna autumnalis*, *Dendrocygna viduata*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dolichonyx oryzivorus*, *Dubusia taeniata*, *Egretta thula*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreovertris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Geothlypis philadelphia*, *Geranoaetus melanoleucus*, *Glaucidium jardinii*, *Grallaria quitesis*, *Grallaria ruficapilla*, *Grallaria squamigera*, *Haplospiza rustica*, *Heliangelus amethysticollis*, *Heliangelus exortis*, *Hellmayrea gularis*, *Hemispingus superciliosus*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus icterus*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Lafresnaya lafresnayi*, *Leiothlypis peregrina*, *Leptasthenura andicola*, *Lesbia nuna*, *Lesbia victoriae*, *Machetornis rixosa*, *Margarornis squamiger*, *Mecocerculus leucophrys*, *Megascops choliba*, *Melanerpes rubricapillus*, *Merganetta armata*, *Metallura tyrianthina*, *Milvago chimachima*, *Mimus gilvus*, *Mniotilta varia*, *Molothrus bonariensis*, *Muscisaxicola alpinus*, *Myiarchus crinitus*, *Myiodynastes luteiventris*, *Myiotheretes fumigatus*, *Myiotheretes striatocollis*, *Myiothlypis nigrocristata*, *Nomonyx dominicus*, *Nycticorax nycticorax*, *Ochthoeca cinnamomeiventris*, *Ochthoeca diadema*, *Ochthoeca fumicolor*, *Ochthoeca rufipectoralis*, *Orochelidon murina*, *Oxyopogon guerini*, *Oxyura jamaicensis*, *Pachyrhamphus versicolor*, *Pandion haliaetus*, *Pardirallus maculatus*, *Parkesia noveboracensis*, *Patagioenas fasciata*, *Petrochelidon pyrrhonota*, *Phaetusa simplex*, *Phalacrocorax brasilianus*, *Pheucticus aureoventris*, *Pheucticus ludovicianus*, *Phimosus infuscatus*, *Phrygilus unicolor*, *Phyllomyias uropygialis*, *Piaya cayana*, *Picoides fumigatus*, *Pipraeidea melanonota*, *Pipreola riefferii*, *Piranga olivacea*, *Piranga rubra*, *Pitangus sulphuratus*, *Pluvialis dominica*, *Podilymbus podiceps*, *Porphyrio martinicus*, *Porzana carolina*, *Progne tapera*, *Protonotaria citrea*, *Pseudocolopteryx acutipennis*, *Pseudoscops clamator*, *Pterophanes cyanopterus*, *Pygochelidon cyanoleuca*, *Pyrocephalus rubinus*, *Pyrrhomyias cinnamomeus*, *Quiscalus lugubris*, *Rallus semiplumbeus*, *Ramphocelus dimidiatus*, *Ramphomicron microrhynchum*, *Riparia riparia*, *Rupornis magnirostris*, *Sayornis nigricans*, *Serpophaga cinerea*, *Setophaga castanea*, *Setophaga cerulea*, *Setophaga petechia*, *Setophaga ruticilla*, *Setophaga striata*, *Sicalis flaveola*, *Sicalis luteola*, *Spiza americana*, *Sporagra spinescens*, *Sturnella magna*, *Synallaxis subpudica*, *Systellura longirostris*, *Tangara vitriolina*, *Thraupis episcopus*, *Thraupis palmarum*, *Tringa flavipes*, *Tringa melanoleuca*, *Tringa solitaria*, *Troglodytes aedon*, *Troglodytes solstitialis*, *Turdus fuscater*, *Turdus ignobilis*, *Tyrannus melancholicus*, *Tyrannus savana*, *Tyrannus tyrannus*, *Tyto alba*, *Vanellus chilensis*, *Vireo olivaceus*, *Vultur gryphus*, *Zenaida auriculata*, *Zonotrichia capensis*

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

pages: 836

issue:

year: 1986

0: Hilty, Steven

1: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigristrostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coccyzus pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerinii, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:

reproductionUnstructured: Ocho individuos en condición reproductiva entre febrero y septiembre en la cordillera Central y Oriental. Se observaron juveniles dependientes en marzo en el PNN Puracé (Hilty & Brown, 1986). También volantones entre marzo y junio, y un nido en el mes de septiembre en Bogotá. Se ha descrito su nido como una copa voluminosa construida con ramitas, pastos secos, recubierto con musgo en su interior y con hojas de bambú en el exterior. El nido contenía un huevo azul verdoso pálido con puntos y manchas cafés y lila concentrados en el extremo más ancho. El nido presentaba las siguientes características: 12 cm diámetro externo, 10 cm altura, 6-5 cm de diámetro interno, 6 cm de profundidad; construido a 40 cm por encima del suelo en un denso matorral de hierbas, con el apoyo de los pastos altos (1-5 m) y un tallo de bambú (Jaramillo, 2011).

0

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

audience

rightsHolder:
rights:
license:
modified:
created:

agent

mimeType:
dataType:
identifier:
relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedBy: TraitBank
measurementDeterminedDate:
measurementUnit:

measurementAccuracy:
measurementValue: Sexual
measurementType: Tipo de Reproducción
measurementID:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:
additionalInformation:

0

link: <http://www.hbw.com/node/61987>

keywords: Atlapetes pallidinucha

0: del Hoyo, J

1: Elliott, A

2: Sargatal, J

3: Christie, D A

4: de Juana, E

institution:

address:

publisher: Lynx Edicions

accessed:

websites:

chapter:

series:

issue:

year: 2011

0: Jaramillo, A

title: Pale-naped Brush-finch (Atlapetes pallidinucha)

source: Handbook of the Birds of the World Alive

type: book_section

tags: Atlapetes pallidinucha

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
feedingUnstructured: Se alimenta de una gran variedad de insectos, semillas y frutos (Jaramillo, 2011).

0

type: Heterótrofos - Omnívoros

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

0

keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata

abstract:

last_modified:

created:

group_id:

profile_id:

id:

1

keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigristrois, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coccyzus pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreovertris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striatocollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerinii, Oxyura jamaicensis, Pachyrhamphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrion martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus

semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

pages: 836

issue:

year: 1986

0: Hilty, Steven

1: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigristrostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coccyzus pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliosus, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striatocollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerini, Oxyura jamaicensis, Pachyrhamphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga

petechia,Setophaga ruticilla,Setophaga striata,Sicalis flaveola,Sicalis luteola,Spiza americana,Sporagra
spinescens,Sturnella magna,Synallaxis subpudica,Systellura longirostris,Tangara vitriolina,Thraupis
episcopus,Thraupis palmarum,Tringa flavipes,Tringa melanoleuca,Tringa solitaria,Troglodytes aedon,Troglodytes
solstitialis,Turdus fuscater,Turdus ignobilis,Tyrannus melancholicus,Tyrannus savana,Tyrannus tyrannus,Tyto
alba,Vanellus chilensis,Vireo olivaceus,Vultur gryphus,Zenaida auriculata,Zonotrichia capensis

abstract:

last_modified:

created:

group_id:

profile_id:

id:

2

keywords: Accipiter striatus,Adelomyia melanogenys,Aglaiocercus kingii,Amblycercus holosericeus,Anas
andium,Andigena nigristrostris,Anisognathus igniventris,Asthenes flammulata,Atlapetes pallidinucha,Atlapetes
schistaceus,Bubulcus ibis,Calidris bairdii,Chalcostigma heteropogon,Chordeiles minor,Cinclus
leucocephalus,Cinnycerthia unirufa,Cistothorus platensis,Coeligena bonapartei,Colibri thalassinus,Coragyps
atratus,Diglossa albilatera,Diglossa cyanea,Diglossa humeralis,Diglossa sittoides,Eriocnemis vestita,Falco
peregrinus,Gallinago nobilis,Gallinula galeata,Geranoaetus melanoleucus,Grallaria quitesis,Grallaria
ruficapilla,Grallaria squamigera,Haplospiza rustica,Mecocerculus leucophrys,Mniotilta varia,Molothrus
bonariensis,Picoides fumigatus,Piranga olivacea,Pyrocephalus rubinus,Rupornis magnirostris,Sicalis
flaveola,Sporagra spinescens,Sturnella magna,Tringa solitaria,Zonotrichia capensis

institution:

address:

publisher: Christopher Helm

accessed:

websites:

chapter:

series:

pages: 880

issue:

year: 2006

0: Restall, R

1: Rodner, C

2: Lentino, M

title: Birds of Northern South America: An Identification Guide, Volume 1: Species Accounts.

source:

type: book

tags: Accipiter striatus,Aglaiocercus kingii,Amblycercus holosericeus,Anas andium,Andigena
nigristrostris,Anisognathus igniventris,Asthenes flammulata,Atlapetes pallidinucha,Atlapetes schistaceus,Bubulcus
ibis,Calidris bairdii,Chalcostigma heteropogon,Chordeiles minor,Cinclus leucocephalus,Cinnycerthia
unirufa,Cistothorus platensis,Coeligena bonapartei,Colibri thalassinus,Coragyps atratus,Diglossa albilatera,Diglossa
cyanea,Diglossa humeralis,Diglossa sittoides,Eriocnemis vestita,Falco peregrinus,Gallinago nobilis,Gallinula
galeata,Geranoaetus melanoleucus,Grallaria quitesis,Grallaria ruficapilla,Grallaria squamigera,Haplospiza
rustica,Mecocerculus leucophrys,Mniotilta varia,Molothrus bonariensis,Picoides fumigatus,Piranga
olivacea,Pyrocephalus rubinus,Rupornis magnirostris,Sicalis flaveola,Sporagra spinescens,Sturnella magna,Tringa
solitaria,Zonotrichia capensis

abstract:

last_modified:

created:

group_id:

profile_id:

id:

3

isbn: 0292719795
keywords: Amblycercus holosericeus,Asthenes flammulata,Atlapetes pallidinucha,Atlapetes schistaceus,Cinclus leucocephalus,Cinnycerthia unirufa,Cistothorus platensis
institution:
address:
publisher: University of Texas Press
accessed:
websites:
chapter:
series:
pages: 750
issue:
year: 2009
0: Ridgely, R
1: Tudor, G
title: Field Guide to the birds of South America: the passerines
source:
type: book
tags: Amblycercus holosericeus,Asthenes flammulata,Atlapetes pallidinucha,Atlapetes schistaceus,Cinclus leucocephalus,Cinnycerthia unirufa,Cistothorus platensis
abstract:
last_modified:
created:
group_id:
profile_id:
id:

4

link: <http://www.hbw.com/node/61987>
keywords: Atlapetes pallidinucha
0: del Hoyo, J
1: Elliott, A
2: Sargatal, J
3: Christie, D A
4: de Juana, E
institution:
address:
publisher: Lynx Edicions
accessed:
websites:
chapter:
series:
issue:
year: 2011
0: Jaramillo, A
title: Pale-naped Brush-finch (Atlapetes pallidinucha)
source: Handbook of the Birds of the World Alive
type: book_section
tags: Atlapetes pallidinucha
abstract:
last_modified:
created:
group_id:
profile_id:
id:

geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:

behaviorUnstructured: Permanece en parejas o familias que saltan en o cerca del suelo, buscan en el follaje y comen frutas; principalmente a menos de 3 m, regularmente siguen bandadas mixtas; menos escondidizas que otros congéneres y usualmente fácil de ver, a veces es curioso (Asociación Bogotana de Ornitología, 2000; Hilty & Brown, 1986). También se observa en grupos de forrajeo en el sotobosque (Restall et al., 2006) y a menudo forrajea en áreas abiertas especialmente en el día (Ridgely & Tudor, 2009). El canto está conformado por una serie muy aguda y débil de trinos y notas silbadas (Hilty & Brown, 1986). Canta al amanecer y las primeras notas son iguales y descendentes. Cantan a menudo en dúo (Restall et al., 2006). En el norte de Perú, el canto territorial es musical y silábico "Wheet-tew-tew-tew"; los llamado son suaves y agudos "tip" y un "pffe" nasal, el cual repite varias veces (Jaramillo, 2011).

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

[endemicAtomized](#)

distribution

ancillaryData

dataObject:

additionalInformation:

reference

link: <http://www.redalyc.org/pdf/491/49131094009.pdf>

keywords: *Anas andium*, *Andigena nigrirostris*, *Atlapetes pallidinucha*, *Chalcostigma heteropogon*, *Cinclodes albidiventris*, *Cinnycerthia unirufa*, *Coeligena bonapartei*, *Conirostrum rufum*, *Eriocnemis cupreiventris*, *Forpus conspicillatus*, *Heliangelus exortis*, *Muscisaxicola alpinus*, *Tangara vitriolina*

editors

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

pages: 235--287

issue:

volume: 14 (2)

year: 2013

authors

authors: Chaparro-Herrera, S

authors: Echeverry-Galvis, M A

authors: Córdoba-Córdoba, Sergio

authors: Sua-Becerra, A

title: Listado actualizado de las aves endémicas y casi-endémicas de Colombia

source: Biota Colombiana

type: article

tags: *Anas andium*, *Andigena nigrirostris*, *Atlapetes pallidinucha*, *Chalcostigma heteropogon*, *Cinclodes albidiventris*, *Cinnycerthia unirufa*, *Coeligena bonapartei*, *Conirostrum rufum*, *Eriocnemis cupreiventris*, *Forpus conspicillatus*, *Heliangelus exortis*, *Muscisaxicola alpinus*, *Tangara vitriolina*

abstract:

identifiers

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

agent

mimeType:

dataType:

identifier:

distributionUnstructured: El Gorrión-montés Cabeciblanco habita entre los 2400 y 3600msnm, pero es más numeroso por encima de los 3000 msnm (Hilty & Brown, 1986). Es una especie casi endémica de Colombia (Chaparro-Herrera et al., 2013) y se distribuye desde el extremo noroccidente de Venezuela a través de la cordillera de los Andes hasta el oriente y occidente de Ecuador y Perú. La especie en Colombia se encuentra en la cordillera Oriental hasta el sur en Cundinamarca. En la cordillera Central se encuentra desde el suroriente de Antioquia por el sur hasta Nariño (BirdLife International and NatureServe, 2014; Hilty & Brown, 1986).

distributionAtomized

stateProvince: Tolima

ancillaryData

locality:

municipality:

country: Colombia

temporalCoverage

endDate:

startDate:

distributionScope

type:

threatStatus

ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:

locality:

municipality:

county:

stateProvince:

country: Global

0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementType: IUCN
measurementID:

territory

territoryUnstructured:
areaOfOccupancy:
extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

abstract: El Gorrión-montés Cabeciblanco tiene plumajes que varían geográficamente. La forma del norte de los Andes venezolanos y de los Andes orientales de Colombia (pallidinucha) es brillante con color leonado en la zona anterior de la banda de la corona; la subespecie papallactae del sur, que se distribuye desde el sur los Andes centrales de Colombia hacia el sur hasta el norte de Perú, es más pálida en la parte de adelante, pero tiene mucho más brillante el amarillo oliva de la garganta y el pecho, además tiene un color amarillento en la franja de la corona

(Schulenberg, 2010). Se alimenta de una gran variedad de insectos, semillas y frutos (Jaramillo, 2011). Permanece en parejas o familias que saltan en o cerca del suelo, buscan en el follaje y comen frutas; principalmente a menos de 3 m, regularmente siguen bandadas mixtas; menos escondidizas que otros congéneres y usualmente fácil de ver, a veces es curioso (Asociación Bogotana de Ornitología, 2000).

associatedParty

role:
personnelIdentifier:
personnelDirectory:
homepage:
email: enataliagomezm@gmail.com
phone:
postalCode:
country: Colombia
state: Bogotá Distrito Capital
city: Bogotá D.C.
address:
organisation:
position:
lastName: Gómez Melendro
firstName: Edith Natalia

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:
mediaURL: <http://www.xeno-canto.org/115468>
description:

subject

source: <http://www.xeno-canto.org>

audience

bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:

agent

contentType:
dataType:
identifier:

_id

_bsontype: ObjectId
id: VÄ)Nð lg#ç20