

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56c4a3fbf0106c67230e7384

Fecha de creación: undefined

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

0

keywords: Accipiter striatus,Adelomyia melanogenys,Aglaiocercus kingii,Amblycercus holosericeus,Anas andium,Andigena nigristrois,Anisognathus igniventris,Asthenes flammulata,Atlapetes pallidinucha,Atlapetes schistaceus,Bubulcus ibis,Calidris bairdii,Catamenia inornata,Chalcostigma heteropogon,Chordeiles minor,Cinclus leucocephalus,Cinnycerthia unirufa,Cistothorus platensis,Coeligena bonapartei,Colibri thalassinus,Coragyps atratus,Diglossa albilatera,Diglossa cyanea,Diglossa humeralis,Diglossa sittoides,Eriocnemis vestita,Falco peregrinus,Gallinago nobilis,Gallinula galeata,Geranoaetus melanoleucus,Grallaria quitesis,Grallaria ruficapilla,Grallaria squamigera,Haplospiza rustica,Mecocerculus leucophrys,Mniotilta varia,Molothrus bonariensis,Picoides fumigatus,Piranga olivacea,Pyrocephalus rubinus,Rupornis magnirostris,Sicalis flaveola,Sporagra spinescens,Sturnella magna,Tringa solitaria,Zonotrichia capensis

institution:

address:

publisher: Christopher Helm

accessed:

websites:

chapter:

series:

pages: 880

issue:

year: 2006

0: Restall, R

1: Rodner, C

2: Lentino, M

title: Birds of Northern South America: An Identification Guide, Volume 1: Species Accounts.

source:

type: book

tags: Accipiter striatus,Aglaiocercus kingii,Amblycercus holosericeus,Anas andium,Andigena nigristrois,Anisognathus igniventris,Asthenes flammulata,Atlapetes pallidinucha,Atlapetes schistaceus,Bubulcus ibis,Calidris bairdii,Catamenia inornata,Chalcostigma heteropogon,Chordeiles minor,Cinclus leucocephalus,Cinnycerthia unirufa,Cistothorus platensis,Coeligena bonapartei,Colibri thalassinus,Coragyps atratus,Diglossa albilatera,Diglossa cyanea,Diglossa humeralis,Diglossa sittoides,Eriocnemis vestita,Falco peregrinus,Gallinago nobilis,Gallinula galeata,Geranoaetus melanoleucus,Grallaria quitesis,Grallaria ruficapilla,Grallaria squamigera,Haplospiza rustica,Mecocerculus leucophrys,Mniotilta varia,Molothrus bonariensis,Picoides fumigatus,Piranga olivacea,Pyrocephalus rubinus,Rupornis magnirostris,Sicalis flaveola,Sporagra spinescens,Sturnella magna,Tringa solitaria,Zonotrichia capensis

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:

habitatUnstructured: Su hábitat está en la selva húmeda característica de tierras altas, allí suele encontrarse en los bordes de bosque o en zonas más abiertas con presencia de vegetación de bajo porte, con arbustos y flores tubulares (Restall et al., 2006).

commonNamesAtomized

ancillaryData

usedBy:
usedIn
distributionUnstructured:
endDate:
startDate:

ancillaryData

type:
language: Alemán
name: Goldbauchmusketier

usesManagementAndConservation

ancillaryData

managementAndConservationUnstructured:

managementAndConservationAtomized

identificationKeys

fullDescription

0

dataObject:
additionalInformation:

0

keywords: Accipiter striatus,Actitis macularius,Adelomyia melanogenys,Agelaiocercus kingii,Amblycercus holosericeus,Ampelion rubrocristatus,Anas andium,Anas clypeata,Anas discors,Andigena nigristrois,Anisognathus igniventris,Ardea alba,Ardea herodias,Arremon assimilis,Asio flammeus,Asio stygius,Asthenes flammulata,Asthenes fuliginosa,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Butorides virescens,Cairina moschata,Calidris bairdii,Calidris melanotos,Cardellina

icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccyzua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreovertris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiostyris alpestris, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus cinerascens, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striatocollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerini, Oxyura jamaicensis, Pachyrhamphus versicolor, Pandion haliaetus, Pardaliparus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne subis, Protonotaria citrea, Pseudocolaptes auratus, Pseudocolaptes auratus, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

fullDescriptionUnstructured: La especie llega a medir hasta 10,9 cm. Posee un pico largo (30 mm) recto y delgado. El macho tiene la frente verde iridiscente, el resto de la coronilla negro; la espalda es verde oscuro brillante y la rabadilla naranja dorado; el pecho y la gorguera son verde iridiscente y las partes bajas son dorado iridiscente. La cola es ligeramente ahorquillada, bronce y las alas son oscuras con una banda rufa. La hembra es verde brillante

por encima, naranja en la rabadilla y por debajo es canela. El pecho es verde y los lados de la garganta manchados de verde mientras que el vientre es manchado de dorado encendido, su cola es bronce con ápice a veces ante (Hilty & Brown, 1986).

briefDescription:

references

link: <http://www.hbw.com/node/55565>

keywords: Coeligena bonapartei

editors

editors: del Hoyo, J

editors: Elliott, A

editors: Sargatal, J

editors: Christie, D A

editors: de Juana, E

institution:

address:

publisher: Lynx Edicions

accessed:

websites:

chapter:

series:

issue:

year: 2013

authors

authors: Z"uchner, T

authors: Motis, A

title: Golden-bellied Starfrontlet (Coeligena bonapartei)

source: Handbook of the Birds of the World Alive

type: book_section

tags: Coeligena bonapartei

abstract:

identifiers

last_modified:

created:

group_id:

profile_id:

id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:

providerLink:

publicationStatus

microReference:

source:

datatype:

identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

replacementNameFor

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

[conservedAgainst](#)

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion
typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:
canonicalAuthorship

authors

year

simple:
simple: (Boissonneau, 1840)

canonicalName

infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:
ref:
uninomial:
simple: Coeligena bonapartei
rank: SPECIES
simple: Coeligena bonapartei (Boissonneau, 1840)

attributes

nomenclaturalCode:
isAnamorphic:
id:

synonymsAtomized

hierarchy

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source: The Catalogue of Life, 3rd January 2011

audience

bibliographicCitation:
rightsHolder:
rights:
license:
title:
modified: 2016-02-17T16:31:10.939Z
created: 2016-02-17T16:31:10.939Z

agent

homepage:
role:

email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:
parentTaxon: Coeligena
higherClassification:
infraspecificEpithet:
specificEpithet:
taxonRank: SPECIES
subGenus:
genus: Coeligena
family: Trochilidae
order: Apodiformes
classHierarchy: Aves
phylum: Chordata
kingdom: Animalia
recommended:
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:
additionalInformation:

0

link: <http://www.hbw.com/node/55565>

keywords: Coeligena bonapartei

0: del Hoyo, J

1: Elliott, A

2: Sargatal, J

3: Christie, D A

4: de Juana, E

institution:

address:

publisher: Lynx Edicions

accessed:

websites:

chapter:

series:

issue:

year: 2013

0: Z"uchner, T

1: Motis, A

title: Golden-bellied Starfrontlet (Coeligena bonapartei)

source: Handbook of the Birds of the World Alive

type: book_section
tags: Coeligena bonapartei
abstract:
last_modified:
created:
group_id:
profile_id:
id:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
reproductionUnstructured: Acerca de su reproducción sólo se conoce que el periodo reproductivo se presenta entre enero y julio (Züchner & Motis, 2013).

0

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

audience

rightsHolder:
rights:
license:
modified:
created:

agent

mimeType:
dataType:
identifier:
relatedTo:
measurementRemarks:
measurementMethod:

measurementDeterminedBy: TraitBank
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue: Sexual
measurementType: Tipo de Reproducción
measurementID:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:

additionalInformation:

0

keywords: Accipiter striatus, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Anas andium, Andigena nigrirostris, Anisognathus igniventris, Asthenes flammulata, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Calidris bairdii, Catamenia inornata, Chalcostigma heteropogon, Chordeiles minor, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coeligena bonapartei, Colibri thalassinus, Coragyps atratus, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa sittoides, Eriocnemis vestita, Falco peregrinus, Gallinago nobilis, Gallinula galeata, Geranoaetus melanoleucus, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Mecocerculus leucophrys, Mniotilta varia, Molothrus bonariensis, Picoides fumigatus, Piranga olivacea, Pyrocephalus rubinus, Rupornis magnirostris, Sicalis flaveola, Sporagra spinescens, Sturnella magna, Tringa solitaria, Zonotrichia capensis

institution:

address:

publisher: Christopher Helm

accessed:

websites:

chapter:

series:

pages: 880

issue:

year: 2006

0: Restall, R

1: Rodner, C

2: Lentino, M

title: Birds of Northern South America: An Identification Guide, Volume 1: Species Accounts.

source:

type: book

tags: Accipiter striatus,Agelaiocercus kingii,Amblycercus holosericeus,Anas andium,Andigena nigrirostris,Anisognathus igniventris,Asthenes flammulata,Atlapetes pallidinucha,Atlapetes schistaceus,Bubulcus ibis,Calidris bairdii,Catamenia inornata,Chalcostigma heteropogon,Chordeiles minor,Cinclus leucocephalus,Cinnycerthia unirufa,Cistothorus platensis,Coeligena bonapartei,Colibri thalassinus,Coragyps atratus,Diglossa albilatera,Diglossa cyanea,Diglossa humeralis,Diglossa sittoides,Eriocnemis vestita,Falco peregrinus,Gallinago nobilis,Gallinula galeata,Geranoaetus melanoleucus,Grallaria quitesis,Grallaria ruficapilla,Grallaria squamigera,Haplospiza rustica,Mecocerculus leucophrys,Mniotilta varia,Molothrus bonariensis,Picoides fumigatus,Piranga olivacea,Pyrocephalus rubinus,Rupornis magnirostris,Sicalis flaveola,Sporagra spinescens,Sturnella magna,Tringa solitaria,Zonotrichia capensis

abstract:

last_modified:

created:

group_id:

profile_id:

id:

1

link: <http://www.hbw.com/node/55565>

keywords: Coeligena bonapartei

0: del Hoyo, J

1: Elliott, A

2: Sargatal, J

3: Christie, D A

4: de Juana, E

institution:

address:

publisher: Lynx Edicions

accessed:

websites:

chapter:

series:

issue:

year: 2013

0: Z"uchner, T

1: Motis, A

title: Golden-bellied Starfrontlet (Coeligena bonapartei)

source: Handbook of the Birds of the World Alive

type: book_section

tags: Coeligena bonapartei

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:
created:
mimeType:
dataType:
identifier:
feedingUnstructured: Visitan flores tubulares, arbustos, epifitas, y árboles de bajo porte en los cuales pueden forrajear de forma solitaria (Restall et al., 2006). También se alimentan cazando insectos y artrópodos en vuelo (Züchner & Motis, 2013).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

0

keywords: Accipiter striatus, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Anas andium, Andigena nigristrois, Anisognathus igniventris, Asthenes flammulata, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Calidris bairdii, Catamenia inornata, Chalcostigma heteropogon, Chordeiles minor, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coeligena bonapartei, Colibri thalassinus, Coragyps atratus, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa sittoides, Eriocnemis vestita, Falco peregrinus, Gallinago nobilis, Gallinula galeata, Geranoaetus melanoleucus, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Mecocerculus leucophrys, Mniotilta varia, Molothrus bonariensis, Picoides fumigatus, Piranga olivacea, Pyrocephalus rubinus, Rupornis magnirostris, Sicalis flaveola, Sporagra spinescens, Sturnella magna, Tringa solitaria, Zonotrichia capensis

institution:

address:

publisher: Christopher Helm

accessed:

websites:

chapter:

series:

pages: 880

issue:

year: 2006

0: Restall, R

1: Rodner, C

2: Lentino, M

title: Birds of Northern South America: An Identification Guide, Volume 1: Species Accounts.

source:

type: book

tags: Accipiter striatus,Agelaiocercus kingii,Amblycercus holosericeus,Anas andium,Andigena nigrirostris,Anisognathus igniventris,Asthenes flammulata,Atlapetes pallidinucha,Atlapetes schistaceus,Bubulcus ibis,Calidris bairdii,Catamenia inornata,Chalcostigma heteropogon,Chordeiles minor,Cinclus leucocephalus,Cinnycerthia unirufa,Cistothorus platensis,Coeligena bonapartei,Colibri thalassinus,Coragyps atratus,Diglossa albilatera,Diglossa cyanea,Diglossa humeralis,Diglossa sittoides,Eriocnemis vestita,Falco peregrinus,Gallinago nobilis,Gallinula galeata,Geranoaetus melanoleucus,Grallaria quitesis,Grallaria ruficapilla,Grallaria squamigera,Haplospiza rustica,Mecocerculus leucophrys,Mniotilta varia,Molothrus bonariensis,Picoides fumigatus,Piranga olivacea,Pyrocephalus rubinus,Rupornis magnirostris,Sicalis flaveola,Sporagra spinescens,Sturnella magna,Tringa solitaria,Zonotrichia capensis

abstract:

last_modified:

created:

group_id:

profile_id:

id:

1

keywords: Actitis macularius,Ampelion rubrocristatus,Anas discors,Anisognathus igniventris,Ardea alba,Arremon assimilis,Asio flammeus,Asio stygius,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Catharus fuscescens,Catharus ustulatus,Chaetocercus mulsant,Chrysomus icterocephalus,Cinnycerthia unirufa,Cistothorus apolinari,Coccyzus americanus,Coeligena bonapartei

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: Actitis macularius,Ampelion rubrocristatus,Anas discors,Anisognathus igniventris,Ardea alba,Arremon assimilis,Asio flammeus,Asio stygius,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Catharus fuscescens,Catharus ustulatus,Chaetocercus mulsant,Chrysomus icterocephalus,Cinnycerthia unirufa,Cistothorus apolinari,Coccyzus americanus,Coeligena bonapartei

abstract:

last_modified:

created:
group_id:
profile_id:
id:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:

behaviorUnstructured: Su comportamiento es similar al de todos los colibríes, son animales solitarios, sólo se reúnen para reproducirse (Restall et al., 2006).
Un suiii suave, especialmente en vuelo. En persecuciones aéreas emite un trino rápido seguido por una serie de silbidos y notas agudas y enérgicas firrrrr fri –fzi –fzi –fzi-fzi –fzi zi zi zi que puede durar hasta medio minuto (Asociación Bogotana de Ornitología, 2000).

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

[endemicAtomized](#)

distribution

ancillaryData

dataObject:

additionalInformation:

reference

keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccyzus pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreovertris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerinii, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

editors

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:
pages: 836
issue:
year: 1986

authors

authors: Hilty, Steven
authors: Brown, William
title: A Guide to the Birds of Colombia
source:
type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccyzua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Mactornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerinii, Oxyura jamaicensis, Pachyrhamphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

abstract:

identifiers

last_modified:
created:

group_id:
profile_id:
id:
geoPoint:
location:
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

[agent](#)

contentType:
dataType:
identifier:
distributionUnstructured: Esta especie es considerada como casi-endémica ya que tiene un rango de vida muy específico, pues solo se distribuye en Venezuela y Colombia (Chaparro-Herrera et al., 2013), allí aunque son aves locales solo presentan dos pequeñas poblaciones alopátricas (Restall et al., 2006). En el país habita las zonas altas de la cordillera Oriental en los departamentos de Santander, Boyacá y Cundinamarca (BirdLife International and NatureServe, 2014), más exactamente desde la Serranía del Perijá en Santander hasta Fusagasugá en Cundinamarca descendiendo por la cordillera Oriental (Hilty & Brown, 1986). El Inca Dorado habita las tierras altas de la cordillera Oriental en Colombia, está reportado entre los 2150 hasta los 3000 msnm. Mientras que en Venezuela la especie se reporta entre los 1400 y los 3200 msnm (Hilty & Brown, 1986).

[distributionAtomized](#)

stateProvince: Santander

[ancillaryData](#)

locality:
municipality:
country: Colombia

temporalCoverage

endDate:
startDate:

distributionScope

type:

[threatStatus](#)

[ancillaryData](#)

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES: Apéndice II

locality:

municipality:

county:

stateProvince:

country: Global

0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:

measurementRemarks:

measurementMethod:

[measurementDeterminedBy](#)

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementType: IUCN

measurementID:

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: La especie es poco común y local en bordes de selva (Hilty & Brown, 1986). Sobre su ecología básica existe poca información, pero se conoce que habita en los bosques enanos y bordes de bosque donde abundan flores tubulares que puedan proveerles néctar. También capturan insectos durante su vuelo y para su identificación se debe tener en cuenta su característico pico recto y largo y la iridiscencia dorada en su vientre y rabadilla (Sedgwick, 2011).

associatedParty

role:

personnelIdentifier:

personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone:

postalCode:

country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address:

organisation:

position:

lastName: Gómez Melendro

firstName: Edith Natalia

ancillaryData

dataObject:

additionalInformation:

reference

geoPoint:

location:

thumbnailURL:

mediaURL: <https://www.flickr.com/photos/ken-ichi/10714446544/>

description:

subject

source: https://farm6.staticflickr.com/5481/10714446544_1da47f0ae6.jpg

audience

bibliographicCitation: The mind of a hummingbird is all murder all the time.

rightsHolder:

rights:

license: <http://creativecommons.org/licenses/by-nc/2.0/>

modified:

created:

agent

contentType:

data Type:

identifier:

_id

_bson type: ObjectId

id: VÄ£ûð lg#ç8@