

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56d59f133c16479905cba8f2

Fecha de creación: 2016-03-01T13:54:29.632Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Lina María Hinestroza B.

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

data Type:

identifier:

habitatUnstructured: Está especie se encuentra ampliamente distribuida en los bordes de bosques caducifolios, de galería, de sabana, secundarios y rastrojos altos de los Llanos de la Orinoquia (McNish 2007).

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

[ancillaryData](#)

type:

language: Alemán

name: Cayenne-Kuckuck

creation_date: 2016-03-01T13:54:29.632Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

fullDescriptionUnstructured:
briefDescription:

references

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:
providerLink:

publicationStatus

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

replacementNameFor

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:

datatype:
identifier:
note:
ruleConsidered:

[conservedAgainst](#)

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

[spellingCorrentionOf](#)

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion

typeOfType:

[lectotypeMicroReferenceVoucher](#)

[lectotypePublicationVoucher](#)

[voucherReference](#)

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:
canonicalAuthorship

authors

year

simple:
simple: (Linnaeus, 1766)
canonicalName
infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:
ref:
uninomial:
simple: Piaya cayana
rank: SPECIES
simple: Piaya cayana (Linnaeus, 1766)

attributes

nomenclaturalCode:
isAnamorphic:
id:

synonymsAtomized

hierarchy

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source: The Catalogue of Life, 3rd January 2011

audience

bibliographicCitation:

rightsHolder:

rights:

license:

title:

modified: 2016-03-01T13:54:09.629Z

created: 2016-03-01T13:54:09.629Z

agent

homepage:

role:

email:

phone:

address:

position:

organisation:

lastName:

firstName:

contentType:

dataType:

identifier:

parentTaxon: Piaya

higherClassification:

infraspecificEpithet:

specificEpithet:

taxonRank: SPECIES

subGenus:

genus: Piaya

family: Cuculidae

order: Cuculiformes

classHierarchy: Aves

phylum: Chordata

kingdom: Animalia

recommended:

classification:

invasiveness

invasivenessUnstructured:

reproduction

reproductionUnstructured:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Lina María Hinestroza B.

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

feedingUnstructured: A esta especie se le puede encontrar en las bandadas mixtas de aves insectívoras (McNish 2007).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:
bibliographicCitation:
rightsHolder:
rights: Lina María Hinestroza B.
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:

created:
mimeType:
dataType:
identifier:

behaviorUnstructured: Presenta movimientos parecidos a los de las ardillas (Cuadros 2004). Se le puede encontrar en solitario, parejas o bandadas (McNish 2007). Piaya cayana tiene varias llamadas distintivas, ninguna frecuente. La más común es Chic, cuah fuerte y seco, un fuerte yiiip-cariir, reminiscente de la llamada del Bichofué (Ridgely 1976), stit fuerte o siit-it en alarma. El canto es juiiip fuerte y resonante, repetido 5-8 veces (Willis & Eisenmann 1979 en Hilty 2001).

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

[endemicAtomized](#)

[distribution](#)

[ancillaryData](#)

dataObject:
additionalInformation:

[reference](#)

geoPoint:
location:
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:

rightsHolder:

rights: Lina María Hinestroza B.

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

[agent](#)

mimeType:

dataType:

identifier:

distributionUnstructured: Se puede visualizar desde México hacia el sur hasta las Guayanas, Venezuela, Brasil, Argentina y Bolivia. Trinidad (Meyer 1964). Reportada en las zonas tropical 0-1000m y montano a más de 2000m (Salaman et al. 2001).

Se encuentra en todo el país en los climas cálidos y medios (Alvarez 1979). En la costa norte de Colombia al este del Golfo de Urabá, hacia el este hasta la parte baja del río Magdalena y al este de los Andes. En el extremo noroeste de Colombia, la costa oeste del Golfo de urabá y el norte de la Costa Pacífica. Al occidente de Nariño, hacia el norte a lo largo de la costa Pacífica hasta la parte alta del río San Juan, el Valle del Cauca y la vertiente oriental de la parte Norte de los Andes Centrales. Al este de los Andes en Arauca y Boyacá; Probablemente hacia el sur hasta el Amazonas (Meyer 1964).

[distributionAtomized](#)

stateProvince: Córdoba

[ancillaryData](#)

locality:

municipality:

country: Colombia

temporalCoverage

endDate:

startDate:

distributionScope

type:

[threatStatus](#)

[ancillaryData](#)

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:

locality:

municipality:

county:

stateProvince:

country: Global

0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementType: IUCN

measurementID:

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

abstract: Está ave mide 43cm, se distingue por el dorso color castaño, más claro en la garganta y la parte superior del pecho, que cambia luego a gris en la parte baja del pecho y el abdomen, llegando a ser negruzco en la región anal; presenta también la cola larga y gradual con manchas blancas escalonadas (Cuadros 2004).

associatedParty

role:
personnelIdentifier:
personnelDirectory:
homepage:
email: enataliagomezm@gmail.com
phone:
postalCode:
country: Colombia
state: Bogotá Distrito Capital
city: Bogotá D.C.
address:
organisation:
position:
lastName: Gómez Melendro
firstName: Edith Natalia

ancillaryData

dataObject:
additionalInformation: Etimología del nombre científico: Probablemente nombre Indígena de Cayena (Cuadros 2004).

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder:
rights: Lina María Hinestroza B.
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

contentType:
dataType:
identifier:

_id

_bsonType: ObjectId

id: $V\check{O}\check{Y} < G^{TM}\backslash^0\bullet$