

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56e0f72a83c45700544e3cfe

Fecha de creación: 2016-03-10T04:25:20.613Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

habitatUnstructured: Esta especie habita en bosques húmedos de Sudamérica.

[commonNamesAtomized](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

[agent](#)

contentType:
dataFormat:
identifier:
usedBy:

usedIn

distributionUnstructured:
endDate:
startDate:

ancillaryData

type:
language: Español
name: Avispa
creation_date: 2016-03-10T04:25:20.613Z

usesManagementAndConservation

ancillaryData

managementAndConservationUnstructured:

managementAndConservationAtomized

identificationKeys

fullDescription

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
contentType:
dataFormat:
identifier:

fullDescriptionUnstructured: Hendidura del músculo propodeal casi circular; metanoto con un tubérculo estrecho a veces débil; tergo abdominal I sin carena transversa o una bastante suave; tergo abdominal II sin tubérculo o con algo bastante débil; especie anaranjada con marcas negras en cabeza y tórax; alas subhialinas con la región costal negruzca; pubescencia anaranjada pegada al cuerpo; nido marrón como una estrella de 6 puntas.

briefDescription:

references

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:
providerLink:

publicationStatus

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

replacementNameFor

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

conservedAgainst

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:
canonicalAuthorship

authors

year

simple:
simple: Fox, 1904
canonicalName
infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:
ref:
uninomial:
simple: Charterginus fulvus
rank: SPECIES
simple: Charterginus fulvus Fox, 1904

attributes

nomenclaturalCode:
isAnamorphic:
id:

synonymsAtomized

hierarchy

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source: The Catalogue of Life, 3rd January 2011

audience

bibliographicCitation:

rightsHolder:
rights:
license:
title:
modified: 2016-03-10T04:25:00.612Z
created: 2016-03-10T04:25:00.612Z

agent

homepage:
role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:
parentTaxon: Charterginus
higherClassification:
infraspecificEpithet:
specificEpithet:
taxonRank: SPECIES
subGenus:
genus: Charterginus
family: Vespidae
order: Hymenoptera
classHierarchy: Insecta
phylum: Arthropoda
kingdom: Animalia
recommended:
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:

dataType:
identifier:
reproductionUnstructured:

0

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

audience

rightsHolder:
rights:
license:
modified:
created:

agent

contentType:
dataType:
identifier:
relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedBy: TraitBank
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue: Sexual
measurementType: Tipo de Reproducción
measurementID:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataFormat:

identifier:

feedingUnstructured:

0

0

strategy: Depredadores

type: Heterótrofos - Omnívoros

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
behaviorUnstructured: Son especies depredadoras con comportamientos eusociales.

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

[endemicAtomized](#)

[distribution](#)

[ancillaryData](#)

dataObject:
additionalInformation:

[reference](#)

geoPoint:
location:
thumbnailURL:

[mediaURL](#)

description:

subject

source:

audience

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

agent

contentType:

dataType:

identifier:

distributionUnstructured: Se ha registrado en Brasil, Perú, Ecuador, Bolivia y Colombia, en áreas que no superan los 700 msnm. En el país se distribuye en las regiones de la Amazonia, Orinoquía y Pacífica.

distributionAtomized

stateProvince: Nariño

ancillaryData

locality:

municipality:

country: Colombia

temporalCoverage

endDate:

startDate:

distributionScope

type:

threatStatus

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

abstract: Presenta el tergo abdominal I con una notoria copa basal delgada, y el área apical más ancha haciendo parte del abdomen restante; gena estrecha, en perfil menos de 1/2 más ancha que el ojo; clípeo más largo que ancho, margen apical casi recto. Hay mucha variación en el color.

associatedParty

role:
personnelIdentifier:
personnelDirectory:
homepage:
email: enataliagomezm@gmail.com
phone:
postalCode:
country: Colombia
state: Bogotá Distrito Capital
city: Bogotá D.C.
address:
organisation:
position:
lastName: Gómez Melendro
firstName: Edith Natalia

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:

thumbnailURL:

mediaURL: <https://www.flickr.com/photos/124722992@N03/14292405190/in/photolist-nLYj89>

description:

subject

source: https://farm4.staticflickr.com/3908/14292405190_e7f8958aa6.jpg

audience

bibliographicCitation: Wasp nest from South America. Ventral view. The nest is attached to the underside of a broad leaf in primary rain forest. The entrance is unusual in being dorsal. The six peripheral projections are sometimes termed 'drip fingers' and possibly help to drain off rain water from the dorsal surface. 'Drip fingers' are also shown to advantage in exhibit XV.

rightsHolder: NHM WASPS

rights:

license: <http://creativecommons.org/licenses/by/2.0/>

modified:

created:

agent

contentType:

dataType:

identifier:

_id

_bsontype: ObjectID

id: Vâ:-*fÄW Dãã