

## Reporte de Fichas para el usuario: Edgar Herrera

Ficha ID: 56e794c783c45700544e41af

Fecha de creación: 2016-03-15T04:52:09.663Z

creation\_date: 2016-03-15T04:52:09.663Z

usesManagementAndConservation

### [ancillaryData](#)

dataObject:

additionalInformation:

### [reference](#)

keywords: Touit stictopterus

### [editors](#)

editors: L. M. Renjifo A. M. Franco, J D Amaya G H Kattan \& B López-Lanús (Eds.).

institution:

address:

publisher: Instituto Alexander von Humboldt, Ministerio del Medio Ambiente

accessed:

websites:

chapter:

series:

pages: 218--220

issue:

year: 2002

### [authors](#)

authors: Rodr\iguez-Mahecha, J. V.

authors: Renjifo, L.M.

title: Touit stictoptera [sic].

source: Libro Rojo de Aves de Colombia

type: book\_section

tags: Touit stictopterus

abstract:

### [identifiers](#)

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

### [mediaURL](#)

description:

### [subject](#)

source:

### [audience](#)

bibliographicCitation:

rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

## agent

mimeType:  
dataType:  
identifier:  
managementAndConservationUnstructured: Aunque actualmente en el país existen áreas protegidas que abarcan la distribución de esta especie, es conveniente estimular mecanismos que garanticen la protección efectiva de estas zonas y el desarrollo de programas y proyectos de conservación; así como adelantar estudios sobre su estado poblacional, su dinámica migratoria, y sus requerimientos mínimos de hábitat (Rodríguez-Mahecha y Renjifo 2002). Igualmente esta especie está incluida en el Apéndice II de la CITES.

## managementAndConservationAtomized

### identificationKeys

### fullDescription

fullDescriptionUnstructured: 18 cm de longitud. Su pico es color gris, cerca del ápice amarillento. El macho presenta coloración principalmente verde; la parte anterior de la cara es color verde amarillenta; las cobertoras alares superiores son color café con los ápices blancos anteados dando una apariencia punteada; presenta un parche color naranja opaco en las cobertoras medias externas; la superficie inferior de las rémiges es color verde azulado; la cola es corta y con forma cuadrada, color verde por encima y amarillo oliva por debajo. La hembra es parecida al macho pero sus cobertoras alares son color verde y están punteadas de color negro, las bridas y la parte anterior de las mejillas son color amarillo más brillante (Hilty y Brown, 2001).  
briefDescription:

## references

link: <http://www.iucnredlist.org/details/full/106001638/0>  
keywords: Touit stictopterus

## editors

institution:  
edition: 2012.1  
address:  
publisher:  
accessed:  
websites:  
chapter:  
series:  
issue:  
year: 2012

## authors

authors: BirdLife International  
title: Touit stictopterus  
source: IUCN Red List of Threatened Species.  
type: miscellany  
tags: Touit stictopterus  
abstract:

identifiers

last\_modified:  
created:  
group\_id:  
profile\_id:  
id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:  
providerLink:

publicationStatus

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

replacementNameFor

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

sanctioned

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

laterHomonymOf

microReference:  
source:  
datatype:  
identifier:

source:  
datatype:  
identifier:  
note:  
ruleConsidered:

conservedAgainst

basedOn

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

basionym

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:  
microReference:

year:  
publishedIn  
simple:  
source:  
datatype:  
identifier:  
specialAuthorship

authors

year  
simple:  
canonicalAuthorship

authors

year  
simple:  
simple: (P. L. Sclater, 1862)  
canonicalName  
infraspecificEpithet:  
specificEpithet:  
infragenericEpithet:  
linkType:  
ref:  
uninomial:  
simple: Touit stictopectus  
rank: SPECIES  
simple: Touit stictopectus (P. L. Sclater, 1862)

attributes  
nomenclaturalCode:  
isAnamorphic:  
id:

synonymsAtomized

ancillaryData

synonymStatus:  
synonymName  
anyTwo:

anyOne

providerLink:  
microReference:  
publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:

identifier:

reletedName

source:

datatype:

identifier:

note:

ruleConsidered:

microReference:

publishedIn

source:

datatype:

identifier:

reletedName

source:

datatype:

identifier:

note:

ruleConsidered:

typeNameEntity

source:

datatype:

identifier:

source:

datatype:

identifier:

source:

datatype:

identifier:

typeVoucherEntity

typeOfTypes:

simple:

microReference:

year:

source:

datatype:

identifier:

[combinationAuthorship](#)

basionymAuthorship

simple:

authorship

simple:

simple: (P. L. Sclater, 1862)

epithet

infraspecificEpithet:

specificEpithet:

infragenericEpithet:

genus

linkType:

ref:

uninomial:

simple: Touit stictoptera

rank: SPECIES

simple: Touit stictoptera (P. L. Sclater, 1862)

nomenclaturalCode:

isAnamorphic:

id:

## [hierarchy](#)

## [ancillaryData](#)

dataObject:

additionalInformation:

## [reference](#)

geoPoint:

location:

thumbnailURL:

## [mediaURL](#)

description:

## [subject](#)

source: The Catalogue of Life, 3rd January 2011

## [audience](#)

bibliographicCitation:

rightsHolder:

rights:

license:

title:

modified: 2016-03-15T04:50:28.956Z

created: 2016-03-15T04:50:28.956Z

## [agent](#)

homepage:

role:

email:

phone:

address:

position:

organisation:

lastName:

firstName:

contentType:

data Type:

identifier:

parentTaxon: Touit



higherClassification:  
infraspecificEpithet:  
specificEpithet:  
taxonRank: SPECIES  
subGenus:  
genus: Touit  
family: Psittacidae  
order: Psittaciformes  
classHierarchy: Aves  
phylum: Chordata  
kingdom: Animalia  
recommended:  
classification:

commonNameAtomized

## ancillaryData

usedBy:  
usedIn  
distributionUnstructured:  
endDate:  
startDate:

## ancillaryData

type:  
language: Inglés  
name: Spot-winged Parrotlet

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:  
additionalInformation:

0

keywords: Touit stictopterus  
0: L. M. Renjifo A. M. Franco, J D Amaya G H Kattan \& B López-Lanús (Eds.).  
institution:  
address:  
publisher: Instituto Alexander von Humboldt, Ministerio del Medio Ambiente  
accessed:  
websites:  
chapter:  
series:  
pages: 218--220  
issue:  
year: 2002  
0: Rodr'iguez-Mahecha, J. V.  
1: Renjifo, L.M.

title: Touit stictoptera [sic].  
source: Libro Rojo de Aves de Colombia  
type: book\_section  
tags: Touit stictopterus  
abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:

1

keywords: Touit stictopterus  
0: del Hoyo, Josep  
1: Elliott, Andrew  
2: Sargatal, Jordi  
institution:  
address:  
publisher: Lynx Edicions  
accessed:  
websites:  
chapter:  
series:  
issue:  
year: 1997  
0: del Hoyo, Josep  
1: Elliott, Andrew  
2: Sargatal, Jordi  
title: Handbook of the Birds of the World. Vol. 4. Sandgrouse to Cuckoos  
source:  
type: book  
tags: Touit stictopterus  
abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
reproductionUnstructured: En Colombia no se conocen datos de su biología reproductiva; pero en Ecuador se ha registrado individuos en condición reproductiva en marzo (Rodríguez-Mahecha y Renjifo 2002; del Hoyo et al. 1997).

molecularData

molecularDataUnstructured:

migratory

0

dataObject:

additionalInformation:

0

keywords: Touit stictopterus

0: L. M. Renjifo A. M. Franco, J D Amaya G H Kattan \& B López-Lanús (Eds.).

institution:

address:

publisher: Instituto Alexander von Humboldt, Ministerio del Medio Ambiente

accessed:

websites:

chapter:

series:

pages: 218--220

issue:

year: 2002

0: Rodr'iguez-Mahecha, J. V.

1: Renjifo, L.M.

title: Touit stictoptera [sic].

source: Libro Rojo de Aves de Colombia

type: book\_section

tags: Touit stictopterus

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataObject:

identifier:

migratoryUnstructured: Al parecer presentan migraciones altitudinales condicionadas por la cosecha de frutos de algunas especies silvestres (Rodríguez-Mahecha y Renjifo 2002).

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:

additionalInformation:

0

keywords: *Actitis macularius*, *Amazilia amabilis*, *Amazilia franciae*, *Amazilia saucerrottei*, *Anas cyanoptera*, *Anthracothorax nigricollis*, *Ara ararauna*, *Ara chloropterus*, *Ara macao*, *Ara militaris*, *Ara severus*, *Aratinga weddellii*, *Bartramia longicauda*, *Brotogeris cyanoptera*, *Brotogeris jugularis*, *Brotogeris sanctithomae*, *Brotogeris versicolurus*, *Bubulcus ibis*, *Buthraupis wetmorei*, *Calidris bairdii*, *Calidris fuscicollis*, *Calidris melanotos*, *Calidris minutilla*, *Catharus fuscescens*, *Catharus minimus*, *Catharus ustulatus*, *Chaetura meridionalis*, *Chalybura buffonii*, *Chlorostilbon mellisugus*, *Chordeiles minor*, *Claravis pretiosa*, *Coccyzus americanus*, *Coccyzus melacoryphus*, *Columba livia*, *Contopus sordidulus*, *Contopus virens*, *Crax globulosa*, *Dacnis hartlaubi*, *Dolichonyx oryzivorus*, *Elaenia gigas*, *Elaenia parvirostris*, *Elaenia spectabilis*, *Elaenia strepera*, *Empidonax traillii*, *Empidonomus aurantioatrocristatus*, *Empidonomus varius*, *Eudocimus ruber*, *Florisuga mellivora*, *Furnarius torridus*, *Galbula pastazae*, *Geotrygon montana*, *Grallaria alleni*, *Grallaria rufocinerea*, *Heliothryx barroti*, *Hirundo rustica*, *Hylomanes mototula*, *Icterus spurius*, *Juliamyia julie*, *Lathrotriccus euleri*, *Lepidothrix coronata*, *Leptosittaca branickii*, *Melanerpes chrysoschelus*, *Myiarchus apicalis*, *Myiodynastes luteiventris*, *Myiozetetes cayanensis*, *Myrmotherula assimilis*, *Numenius phaeopus*, *Nyctidromus albicollis*, *Oporornis agilis*, *Orthopsittaca manilata*, *Pachyramphus rufus* (Boddaert 1783), *Parkesia motacilla*, *Paroaria gularis*, *Passer domesticus*, *Petrochelidon pyrrhonota*, *Phaethornis anthophilus*, *Phaethornis superciliosus*, *Pionus menstruus*, *Pipreola chlorolepidota*, *Pluvialis dominica*, *Porphyrio flavirostris*, *Progne chalybea*, *Progne subis*, *Progne tapera*, *Protonotaria citrea*, *Riparia riparia*, *Sporophila caerulescens*, *Sporophila lineola*, *Tachycineta albiventer* (Boddaert 1783), *Thalurelia colombica*, *Thalurelia furcata*, *Threnetes ruckeri*, *Tinamus osgoodi*, *Touit stictopterus*, *Tringa flavipes*, *Tringa melanoleuca*, *Tringa solitaria*, *Tyrannus savana*, *Tyrannus tyrannus*, *Vireo altiloquus*, *Xiphorhynchus picus*

institution:

edition: 3

address:

publisher:

accessed:

websites:

chapter:

series:

pages: 1034

issue:

year: 2001

0: Hilty, S. L.

1: Brown, W. L.

title: Guía de las aves de Colombia

source:

type: book

tags: *Actitis macularius*, *Amazilia amabilis*, *Amazilia franciae*, *Amazilia saucerrottei*, *Anas cyanoptera*, *Anthracothorax nigricollis*, *Ara ararauna*, *Ara chloropterus*, *Ara macao*, *Ara militaris*, *Ara severus*, *Aratinga weddellii*, *Bartramia longicauda*, *Brotogeris cyanoptera*, *Brotogeris jugularis*, *Brotogeris sanctithomae*, *Brotogeris versicolurus*, *Bubulcus ibis*, *Buthraupis wetmorei*, *Calidris bairdii*, *Calidris fuscicollis*, *Calidris melanotos*, *Calidris minutilla*, *Catharus fuscescens*, *Catharus minimus*, *Catharus ustulatus*, *Chaetura meridionalis*, *Chalybura buffonii*, *Chlorostilbon mellisugus*, *Chordeiles minor*, *Claravis pretiosa*, *Coccyzus americanus*, *Coccyzus melacoryphus*, *Columba livia*, *Contopus sordidulus*, *Contopus virens*, *Crax globulosa*, *Dacnis hartlaubi*, *Dolichonyx oryzivorus*, *Elaenia gigas*, *Elaenia parvirostris*, *Elaenia spectabilis*, *Elaenia strepera*, *Empidonax traillii*, *Empidonomus aurantioatrocristatus*, *Empidonomus varius*, *Eudocimus ruber*, *Florisuga mellivora*, *Furnarius torridus*, *Galbula*

pastazae,Geotrygon montana,Grallaria alleni,Grallaria rufocinerea,Heliothryx barroti,Hirundo rustica,Hylomanes momotula,Icterus spurius,Juliamyia julie,Lathrotriccus euleri,Lepidothrix coronata,Leptosittaca branickii,Melanerpes chrysauchen,Myiarchus apicalis,Myiodynastes luteiventris,Myiozetetes cayanensis,Myrmotherula assimilis,Numenius phaeopus,Nyctidromus albigollis,Oporornis agilis,Orthopsittaca manilata,Pachyramphus rufus (Boddaert 1783),Parkesia motacilla,Paroaria gularis,Passer domesticus,Petrochelidon pyrrhonota,Phaethornis anthophilus,Phaethornis superciliosus,Pionus menstruus,Pipreola chlorolepidota,Pluvialis dominica,Porphyrion flavirostris,Progne chalybea,Progne subis,Progne tapera,Protonotaria citrea,Riparia riparia,Sporophila caerulea,Sporophila lineola,Tachycineta albiventer (Boddaert 1783),Thalurania colombica,Thalurania furcata,Threnetes ruckeri,Tinamus osgoodi,Touit stictopterus,Tringa flavipes,Tringa melanoleuca,Tringa solitaria,Tyrannus savana,Tyrannus tyrannus,Vireo altiloquus,Xiphorhynchus picus

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

1

keywords: Touit stictopterus

institution:

address:

publisher: Pica Press

accessed:

websites:

chapter:

series:

issue:

year: 1998

0: Juniper, T.

1: Parr, M.

title: Parrots: a guide to the parrots of the world.

source:

type: book

tags: Touit stictopterus

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

2

keywords: Touit stictopterus

0: L. M. Renjifo A. M. Franco, J D Amaya G H Kattan \& B López-Lanús (Eds.).

institution:

address:

publisher: Instituto Alexander von Humboldt, Ministerio del Medio Ambiente

accessed:

websites:

chapter:

series:

pages: 218--220

issue:

year: 2002

0: Rodr'iguez-Mahecha, J. V.

1: Renjifo, L.M.  
title: Touit stictoptera [sic].  
source: Libro Rojo de Aves de Colombia  
type: book\_section  
tags: Touit stictopterus  
abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
feedingUnstructured: Se alimenta de frutos de los cauchos (Ficus spp.) y de los capes (Clusia spp.), también de maíz y de “matapalos” (Loranthaceae) (Juniper y Parr 1998; Hilty y Brown, 2001; Rodríguez-Mahecha y Renjifo 2002).

environmentalEnvelope

0

dataObject:  
additionalInformation:

0

keywords: Ara ararauna,Ara chloropterus,Ara macao,Ara severus,Aratinga weddellii,Leptosittaca  
branickii,Orthopsittaca manilata,Touit stictopterus

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

pages: 150

issue:

year: 2005

0: Rodríguez-Mahecha, J. V.

1: Rojas, F.

2: Arzuza, D.

3: González, A.

title: Miniguía Loros, Pericos & Guacamayas Neotropicales.

source:

type: book

tags: Ara ararauna,Ara chloropterus,Ara macao,Ara severus,Aratinga weddellii,Leptosittaca branickii,Orthopsittaca manilata,Touit stictopterus

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

environmentalEnvelopeUnstructured: Bosque semihúmedo nublado (Rodríguez-Mahecha 2005).

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

0

keywords: Actitis macularius,Amazilia amabilis,Amazilia franciae,Amazilia saucerrottei,Anas cyanoptera,Anthracothonax nigricollis,Ara ararauna,Ara chloropterus,Ara macao,Ara militaris,Ara severus,Aratinga weddellii,Bartramia longicauda,Brotogeris cyanoptera,Brotogeris jugularis,Brotogeris sanctithomae,Brotogeris versicolurus,Bubulcus ibis,Buthraupis wetmorei,Calidris bairdii,Calidris fuscicollis,Calidris melanotos,Calidris minutilla,Catharus fuscescens,Catharus minimus,Catharus ustulatus,Chaetura meridionalis,Chalybura buffonii,Chlorostilbon mellisugus,Chordeiles minor,Claravis pretiosa,Coccyzus americanus,Coccyzus melacoryphus,Columba livia,Contopus sordidulus,Contopus virens,Crax globulosa,Dacnis hartlaubi,Dolichonyx oryzivorus,Elaenia gigas,Elaenia parvirostris,Elaenia spectabilis,Elaenia strepera,Empidonax traillii,Empidonax aurantioatrocristatus,Empidonax varius,Eudocimus ruber,Florisuga mellivora,Furnarius torridus,Galbula pastazae,Geotrygon montana,Grallaria alleni,Grallaria rufocinerea,Heliothryx barroti,Hirundo rustica,Hylomanes momotula,Icterus spurius,Juliamyia julie,Lathrotriccus euleri,Lepidothrix coronata,Leptosittaca branickii,Melanerpes chrysauchen,Myiarchus apicalis,Myiodynastes luteiventris,Myiozetetes cayanensis,Myrmotherula assimilis,Numenius phaeopus,Nyctidromus albicollis,Oporornis agilis,Orthopsittaca manilata,Pachyrhamphus rufus (Boddaert 1783),Parkesia motacilla,Paroaria gularis,Passer domesticus,Petrochelidon pyrrhonota,Phaethornis

anthophilus, Phaethornis superciliosus, Pionus menstruus, Pipreola chlorolepidota, Pluvialis dominica, Porphyrio flavirostris, Progne chalybea, Progne subis, Progne tapera, Protonotaria citrea, Riparia riparia, Sporophila caerulescens, Sporophila lineola, Tachycineta albiventer (Boddaert 1783), Thalurania colombica, Thalurania furcata, Threnetes ruckeri, Tinamus osgoodi, Touit stictopterus, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Tyrannus savana, Tyrannus tyrannus, Vireo altiloquus, Xiphorhynchus picus

institution:

edition: 3

address:

publisher:

accessed:

websites:

chapter:

series:

pages: 1034

issue:

year: 2001

0: Hilty, S. L.

1: Brown, W. L.

title: Gu'ia de las aves de Colombia

source:

type: book

tags: Actitis macularius, Amazilia amabilis, Amazilia franciae, Amazilia saucerrottei, Anas cyanoptera, Anthracothorax nigricollis, Ara ararauna, Ara chloropterus, Ara macao, Ara militaris, Ara severus, Aratinga weddellii, Bartramia longicauda, Brotogeris cyanoptera, Brotogeris jugularis, Brotogeris sanctithomae, Brotogeris versicolurus, Bubulcus ibis, Buthraupis wetmorei, Calidris bairdii, Calidris fuscicollis, Calidris melanotos, Calidris minutilla, Catharus fuscescens, Catharus minimus, Catharus ustulatus, Chaetura meridionalis, Chalybura buffonii, Chlorostilbon mellisugus, Chordeiles minor, Claravis pretiosa, Coccyzus americanus, Coccyzus melacoryphus, Columba livia, Contopus sordidulus, Contopus virens, Crax globulosa, Dacnis hartlaubi, Dolichonyx oryzivorus, Elaenia gigas, Elaenia parvirostris, Elaenia spectabilis, Elaenia strepera, Empidonax traillii, Empidonax aurantioatrocristatus, Empidonax varius, Eudocimus ruber, Florisuga mellivora, Furnarius torridus, Galbula pastazae, Geotrygon montana, Grallaria alleni, Grallaria rufocinerea, Heliothryx barroti, Hirundo rustica, Hylomanes momotula, Icterus spurius, Juliomyia julie, Lathrotriccus euleri, Lepidothrix coronata, Leptosittaca branickii, Melanerpes chrysauen, Myiarchus apicalis, Myiodynastes luteiventris, Myiozetetes cayanensis, Myrmotherula assimilis, Numenius phaeopus, Nyctidromus albicollis, Oporornis agilis, Orthopsittaca manilata, Pachyramphus rufus (Boddaert 1783), Parkesia motacilla, Paroaria gularis, Passer domesticus, Petrochelidon pyrrhonota, Phaethornis anthophilus, Phaethornis superciliosus, Pionus menstruus, Pipreola chlorolepidota, Pluvialis dominica, Porphyrio flavirostris, Progne chalybea, Progne subis, Progne tapera, Protonotaria citrea, Riparia riparia, Sporophila caerulescens, Sporophila lineola, Tachycineta albiventer (Boddaert 1783), Thalurania colombica, Thalurania furcata, Threnetes ruckeri, Tinamus osgoodi, Touit stictopterus, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Tyrannus savana, Tyrannus tyrannus, Vireo altiloquus, Xiphorhynchus picus

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:



created:  
mimeType:  
dataType:  
identifier:  
behaviorUnstructured: En el este de Ecuador se ha observado bandadas volando rasantes sobre dosel (Hilty y Brown, 2001). En vuelo ch-ch-ch áspero y repetido (Hilty y Brown, 2001).

#### ancillaryData

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights:  
license:  
modified:  
created:  
mimeType:  
dataType:  
identifier:

#### measurementOrFact

relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

#### annualCycle

annualCycleUnstructured:

#### habitat

0

dataObject:  
additionalInformation:

0

keywords: Actitis macularius,Amazilia amabilis,Amazilia franciae,Amazilia saucerrottei,Anas cyanoptera,Anthracothonax nigricollis,Ara ararauna,Ara chloropterus,Ara macao,Ara militaris,Ara severus,Aratinga weddellii,Bartramia longicauda,Brotogeris cyanoptera,Brotogeris jugularis,Brotogeris sanctithomae,Brotogeris versicolurus,Bubulcus ibis,Buthraupis wetmorei,Calidris bairdii,Calidris fuscicollis,Calidris melanotos,Calidris minutilla,Catharus fuscescens,Catharus minimus,Catharus ustulatus,Chaetura meridionalis,Chalybura buffonii,Chlorostilbon mellisugus,Chordeiles minor,Claravis pretiosa,Coccyzus americanus,Coccyzus melacoryphus,Columba livia,Contopus sordidulus,Contopus virens,Crax globulosa,Dacnis hartlaubi,Dolichonyx oryzivorus,Elaenia gigas,Elaenia parvirostris,Elaenia spectabilis,Elaenia strepera,Empidonax traillii,Empidonax

aurantioatrocristatus, Empidonomus varius, Eudocimus ruber, Florisuga mellivora, Furnarius torridus, Galbula pastazae, Geotrygon montana, Grallaria alleni, Grallaria rufocinerea, Heliothryx barroti, Hirundo rustica, Hylomanes momotula, Icterus spurius, Juliamyia julie, Lathrotriccus euleri, Lepidothrix coronata, Leptosittaca branickii, Melanerpes chrysauchen, Myiarchus apicalis, Myiodynastes luteiventris, Myiozetetes cayanensis, Myrmotherula assimilis, Numenius phaeopus, Nyctidromus albigularis, Oporornis agilis, Orthopsittaca manilata, Pachyrhamphus rufus (Boddaert 1783), Parkesia motacilla, Paroaria gularis, Passer domesticus, Petrochelidon pyrrhonota, Phaethornis anthophilus, Phaethornis superciliosus, Pionus menstruus, Pipreola chlorolepidota, Pluvialis dominica, Porphyrio flavirostris, Progne chalybea, Progne subis, Progne tapera, Protonotaria citrea, Riparia riparia, Sporophila caerulea, Sporophila lineola, Tachycineta albiventer (Boddaert 1783), Thalassidroma colymbica, Thalassidroma furcata, Threnetes ruckeri, Tinamus osgoodi, Touit stictopterus, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Tyrannus savana, Tyrannus tyrannus, Vireo altiloquus, Xiphorhynchus picus

institution:

edition: 3

address:

publisher:

accessed:

websites:

chapter:

series:

pages: 1034

issue:

year: 2001

0: Hilty, S. L.

1: Brown, W. L.

title: Guía de las aves de Colombia

source:

type: book

tags: Actitis macularia, Amazilia amabilis, Amazilia franciae, Amazilia saucerrottei, Anas cyanoptera, Anthracoceros nigricollis, Ara ararauna, Ara chloropterus, Ara macao, Ara militaris, Ara severus, Aratinga weddellii, Bartramia longicauda, Brotogeris cyanoptera, Brotogeris jugularis, Brotogeris sanctithomae, Brotogeris versicolurus, Bubulcus ibis, Butor, Butorides, Calidris bairdii, Calidris fuscicollis, Calidris melanotos, Calidris minutilla, Catharus fuscescens, Catharus minimus, Catharus ustulatus, Chaetura meridionalis, Chalybura buffonii, Chlorostilbon mellisugus, Chordeiles minor, Claravis pretiosa, Coccyzus americanus, Coccyzus melacoryphus, Columba livia, Contopus sordidulus, Contopus virens, Crax globulosa, Dacnis hartlaubi, Dolichonyx oryzivorus, Elaenia gigas, Elaenia parvirostris, Elaenia spectabilis, Elaenia strepera, Empidonax traillii, Empidonomus aurantioatrocristatus, Empidonomus varius, Eudocimus ruber, Florisuga mellivora, Furnarius torridus, Galbula pastazae, Geotrygon montana, Grallaria alleni, Grallaria rufocinerea, Heliothryx barroti, Hirundo rustica, Hylomanes momotula, Icterus spurius, Juliamyia julie, Lathrotriccus euleri, Lepidothrix coronata, Leptosittaca branickii, Melanerpes chrysauchen, Myiarchus apicalis, Myiodynastes luteiventris, Myiozetetes cayanensis, Myrmotherula assimilis, Numenius phaeopus, Nyctidromus albigularis, Oporornis agilis, Orthopsittaca manilata, Pachyrhamphus rufus (Boddaert 1783), Parkesia motacilla, Paroaria gularis, Passer domesticus, Petrochelidon pyrrhonota, Phaethornis anthophilus, Phaethornis superciliosus, Pionus menstruus, Pipreola chlorolepidota, Pluvialis dominica, Porphyrio flavirostris, Progne chalybea, Progne subis, Progne tapera, Protonotaria citrea, Riparia riparia, Sporophila caerulea, Sporophila lineola, Tachycineta albiventer (Boddaert 1783), Thalassidroma colymbica, Thalassidroma furcata, Threnetes ruckeri, Tinamus osgoodi, Touit stictopterus, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Tyrannus savana, Tyrannus tyrannus, Vireo altiloquus, Xiphorhynchus picus

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

1

keywords: Ara ararauna, Ara chloropterus, Ara macao, Ara severus, Aratinga weddellii, Leptosittaca

branickii,Orthopsittaca manilata,Touit stictopterus

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

pages: 150

issue:

year: 2005

0: Rodr\iguez-Mahecha, J. V.

1: Rojas, F.

2: Arzuza, D.

3: González, A.

title: Minigu\ia Loros, Pericos \& Guacamayas Neotropicales.

source:

type: book

tags: Ara ararauna,Ara chloropterus,Ara macao,Ara severus,Aratinga weddellii,Leptosittaca branickii,Orthopsittaca manilata,Touit stictopterus

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

habitatUnstructured: Habita áreas selváticas y bosque húmedo montano y áreas de cultivos estacionales (Hilty y Brown, 2001; Rodríguez-Mahecha 2005).

[endemicAtomized](#)

[distribution](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

keywords: Actitis macularius,Amazilia amabilis,Amazilia franciae,Amazilia saucerrottei,Anas cyanoptera,Anthracothonax nigricollis,Ara ararauna,Ara chloropterus,Ara macao,Ara militaris,Ara severus,Aratinga

weddellii,Bartramia longicauda,Brotogeris cyanoptera,Brotogeris jugularis,Brotogeris sanctithomae,Brotogeris versicolurus,Bubulcus ibis,Buthraupis wetmorei,Calidris bairdii,Calidris fuscicollis,Calidris melanotos,Calidris minutilla,Catharus fuscescens,Catharus minimus,Catharus ustulatus,Chaetura meridionalis,Chalybura buffonii,Chlorostilbon mellisugus,Chordeiles minor,Claravis pretiosa,Coccyzus americanus,Coccyzus melacoryphus,Columba livia,Contopus sordidulus,Contopus virens,Crax globulosa,Dacnis hartlaubi,Dolichonyx oryzivorus,Elaenia gigas,Elaenia parvirostris,Elaenia spectabilis,Elaenia strepera,Empidonax traillii,Empidonomus aurantioatrocristatus,Empidonomus varius,Eudocimus ruber,Florisuga mellivora,Furnarius torridus,Galbula pastazae,Geotrygon montana,Grallaria alleni,Grallaria rufocinerea,Heliodytes barroeti,Hirundo rustica,Hylomanes momotula,Icterus spurius,Juliamyia julie,Lathrotriccus euleri,Lepidothrix coronata,Leptosittaca branickii,Melanerpes chrysarchus,Myiarchus apicalis,Myiodynastes luteiventris,Myiozetetes cayanensis,Myrmotherula assimilis,Numenius phaeopus,Nyctidromus albigularis,Oporornis agilis,Orthopsittacus manilata,Pachyrhamphus rufus (Boddaert 1783),Parkesia motacilla,Paroaria gularis,Passer domesticus,Petrochelidon pyrrhonota,Phaethornis anthophilus,Phaethornis superciliosus,Pionus menstruus,Pipreola chlorolepidota,Pluvialis dominica,Porphyrio flavirostris,Progne subis,Progne subis,Progne tapera,Protonotaria citrea,Riparia riparia,Sporophila caerulea,Sporophila lineola,Tachycineta albiventer (Boddaert 1783),Thalassidroma colymbica,Thalassidroma furcata,Threnetes ruckeri,Tinamus osgoodi,Touit stictopterus,Tringa flavipes,Tringa melanoleuca,Tringa solitaria,Tyrannus savana,Tyrannus tyrannus,Vireo altiloquus,Xiphorhynchus picus

## editors

institution:  
edition: 3  
address:  
publisher:  
accessed:  
websites:  
chapter:  
series:  
pages: 1034  
issue:  
year: 2001

## authors

authors: Hilty, S. L.  
authors: Brown, W. L.  
title: Gu'ia de las aves de Colombia  
source:  
type: book

tags: Actitis macularia,Amazilia amabilis,Amazilia franciae,Amazilia saucerrottei,Anas cyanoptera,Anthracoceros nigricollis,Ara ararauna,Ara chloropterus,Ara macao,Ara militaris,Ara severus,Aratinga weddellii,Bartramia longicauda,Brotogeris cyanoptera,Brotogeris jugularis,Brotogeris sanctithomae,Brotogeris versicolurus,Bubulcus ibis,Buthraupis wetmorei,Calidris bairdii,Calidris fuscicollis,Calidris melanotos,Calidris minutilla,Catharus fuscescens,Catharus minimus,Catharus ustulatus,Chaetura meridionalis,Chalybura buffonii,Chlorostilbon mellisugus,Chordeiles minor,Claravis pretiosa,Coccyzus americanus,Coccyzus melacoryphus,Columba livia,Contopus sordidulus,Contopus virens,Crax globulosa,Dacnis hartlaubi,Dolichonyx oryzivorus,Elaenia gigas,Elaenia parvirostris,Elaenia spectabilis,Elaenia strepera,Empidonax traillii,Empidonomus aurantioatrocristatus,Empidonomus varius,Eudocimus ruber,Florisuga mellivora,Furnarius torridus,Galbula pastazae,Geotrygon montana,Grallaria alleni,Grallaria rufocinerea,Heliodytes barroeti,Hirundo rustica,Hylomanes momotula,Icterus spurius,Juliamyia julie,Lathrotriccus euleri,Lepidothrix coronata,Leptosittaca branickii,Melanerpes chrysarchus,Myiarchus apicalis,Myiodynastes luteiventris,Myiozetetes cayanensis,Myrmotherula assimilis,Numenius phaeopus,Nyctidromus albigularis,Oporornis agilis,Orthopsittacus manilata,Pachyrhamphus rufus (Boddaert 1783),Parkesia motacilla,Paroaria gularis,Passer domesticus,Petrochelidon pyrrhonota,Phaethornis anthophilus,Phaethornis superciliosus,Pionus menstruus,Pipreola chlorolepidota,Pluvialis dominica,Porphyrio flavirostris,Progne subis,Progne subis,Progne tapera,Protonotaria citrea,Riparia riparia,Sporophila caerulea,Sporophila lineola,Tachycineta albiventer (Boddaert 1783),Thalassidroma colymbica,Thalassidroma furcata,Threnetes ruckeri,Tinamus osgoodi,Touit stictopterus,Tringa flavipes,Tringa melanoleuca,Tringa

solitaria,Tyrannus savana,Tyrannus tyrannus,Vireo altiloquus,Xiphorhynchus picus  
abstract:

## identifiers

last\_modified:  
created:  
group\_id:  
profile\_id:  
id:  
geoPoint:  
location:  
thumbnailURL:

## mediaURL

description:

## subject

source:

## audience

bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

## agent

contentType:  
dataType:  
identifier:  
distributionUnstructured: Se distribuye en la vertiente oeste de la Cordillera Central (Cauca), vertiente oeste de la Cordillera Oriental (Cundinamarca), Serranía de la Macarena (Meta); probablemente se encuentra en el este de Nariño (Hilty y Brown, 2001; Rodríguez-Mahecha y Renjifo 2002). Región natural Andina (Salaman et al. 2009). Entre 600 y 2400 msnm (Hilty y Brown, 2001; Rodríguez-Mahecha 2005).

## distributionAtomized

stateProvince: Cauca

## ancillaryData

locality:  
municipality:  
country: Colombia

## temporalCoverage

endDate:  
startDate:

## distributionScope

type:

## threatStatus

## ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES: Apéndice II

locality:

municipality:

county:

stateProvince:

country: Global

0: IUCN

measurementValue: Vulnerable (VU)

relatedTo:

measurementRemarks:

measurementMethod:

## measurementDeterminedBy

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementType: IUCN

measurementID:

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

0

dataObject:

additionalInformation:

0

link: <http://www.iucnredlist.org/details/full/106001638/0>

keywords: Touit stictopterus

institution:

edition: 2012.1

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2012

0: BirdLife International

title: Touit stictopterus

source: IUCN Red List of Threatened Species.

type: miscellany

tags: Touit stictopterus

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

1

keywords: Touit stictopterus

0: L. M. Renjifo A. M. Franco, J D Amaya G H Kattan \& B López-Lanús (Eds.).

institution:

address:

publisher: Instituto Alexander von Humboldt, Ministerio del Medio Ambiente

accessed:

websites:

chapter:

series:

pages: 218--220

issue:

year: 2002

0: Rodr'iguez-Mahecha, J. V.

1: Renjifo, L.M.

title: Touit stictoptera [sic].

source: Libro Rojo de Aves de Colombia

type: book\_section

tags: Touit stictopterus

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

populationBiologyUnstructured: El tamaño de la población de esta especie ha sido estimado entre 1000 y 2500 individuos, su población está disminuyendo rápidamente. En 2002, este periquito había perdido el 70% de su hábitat (BirdLife 2012; Rodríguez-Mahecha y Renjifo 2002)

legislation

legislationUnstructured:

directThreats

0

dataObject:  
additionalInformation:

0

keywords: Ara ararauna,Ara chloropterus,Ara macao,Ara severus,Aratinga weddellii,Leptosittaca  
branickii,Orthopsittaca manilata,Touit stictopterus

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

pages: 150

issue:

year: 2005

0: Rodr'iguez-Mahecha, J. V.

1: Rojas, F.

2: Arzuza, D.

3: González, A.

title: Minigu'ia Loros, Pericos \& Guacamayas Neotropicales.

source:

type: book

tags: Ara ararauna,Ara chloropterus,Ara macao,Ara severus,Aratinga weddellii,Leptosittaca branickii,Orthopsittaca  
manilata,Touit stictopterus

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

1

keywords: Actitis macularius,Amazilia amabilis,Amazilia franciae,Amazilia saucerrottei,Anas  
cyanoptera,Anthracothonax nigricollis,Ara ararauna,Ara chloropterus,Ara macao,Ara militaris,Ara severus,Aratinga  
weddellii,Bartramia longicauda,Brotogeris cyanoptera,Brotogeris jugularis,Brotogeris sanctithomae,Brotogeris  
versicolorus,Bubulcus ibis,Buthraupis wetmorei,Calidris bairdii,Calidris fuscicollis,Calidris melanotos,Calidris  
minutilla,Catharus fuscescens,Catharus minimus,Catharus ustulatus,Chaetura meridionalis,Chalybura  
buffonii,Chlorostilbon mellisugus,Chordeiles minor,Claravis pretiosa,Coccyzus americanus,Coccyzus  
melacoryphus,Columba livia,Contopus sordidulus,Contopus virens,Crax globulosa,Dacnis hartlaubi,Dolichonyx  
oryzivorus,Elaenia gigas,Elaenia parvirostris,Elaenia spectabilis,Elaenia strepera,Empidonax traillii,Empidonax  
aurantioatrocristatus,Empidonax varius,Eudocimus ruber,Florisuga mellivora,Furnarius torridus,Galbula  
pastazae,Geotrygon montana,Grallaria alleni,Grallaria rufocinerea,Heliothryx barroti,Hirundo rustica,Hylomanes  
momotula,Icterus spurius,Juliamyia julie,Lathrotriccus euleri,Lepidothrix coronata,Leptosittaca branickii,Melanerpes  
chrysauchen,Myiarchus apicalis,Myiodynastes luteiventris,Myiozetetes cayanensis,Myrmotherula assimilis,Numenius  
phaeopus,Nyctidromus albigollis,Oporornis agilis,Orthopsittaca manilata,Pachyramphus rufus (Boddaert  
1783),Parkesia motacilla,Paroaria gularis,Passer domesticus,Petrochelidon pyrrhonota,Phaethornis  
anthophilus,Phaethornis superciliosus,Pionus menstruus,Pipreola chlorolepidota,Pluvialis dominica,Porphyrion  
flavivirostris,Progne chalybea,Progne subis,Progne tapera,Protonotaria citrea,Riparia riparia,Sporophila  
caerulescens,Sporophila lineola,Tachycineta albiventer (Boddaert 1783),Thalurania colombica,Thalurania  
furcata,Threnetes ruckeri,Tinamus osgoodi,Touit stictopterus,Tringa flavipes,Tringa melanoleuca,Tringa  
solitaria,Tyrannus savana,Tyrannus tyrannus,Vireo altiloquus,Xiphorhynchus picus

institution:

edition: 3

address:



publisher:

accessed:

websites:

chapter:

series:

pages: 1034

issue:

year: 2001

0: Hilty, S. L.

1: Brown, W. L.

title: Guía de las aves de Colombia

source:

type: book

tags: Actitis macularia, Amazilia amabilis, Amazilia franciae, Amazilia saucerrottei, Anas cyanoptera, Anthracoceros nigricollis, Ara ararauna, Ara chloroptera, Ara macao, Ara militaris, Ara severus, Aratinga weddellii, Bartramia longicauda, Brotogeris cyanoptera, Brotogeris jugularis, Brotogeris sanctithomae, Brotogeris versicolurus, Bubulcus ibis, Butor, Butorides, Calidris bairdii, Calidris fuscicollis, Calidris melanotos, Calidris minutilla, Catharus fuscescens, Catharus minimus, Catharus ustulatus, Chaetura meridionalis, Chalybura buffonii, Chlorostilbon mellisugus, Chordeiles minor, Claravis pretiosa, Coccyzus americanus, Coccyzus melacoryphus, Columba livia, Contopus sordidulus, Contopus virens, Crax globulosa, Dacnis hartlaubi, Dolichonyx oryzivorus, Elaenia gigas, Elaenia parvirostris, Elaenia spectabilis, Elaenia strepera, Empidonax traillii, Empidonax aurantioatrocristatus, Empidonax varius, Eudocimus ruber, Florisuga mellivora, Furnarius torridus, Galbula pastazae, Geotrygon montana, Grallaria alleni, Grallaria rufocinerea, Heliothryx barroti, Hirundo rustica, Hylomanes momotula, Icterus spurius, Juliamyia julie, Lathrotriccus euleri, Lepidothrix coronata, Leptosittaca branickii, Melanerpes chrysaeus, Myiarchus apicalis, Myiodynastes luteiventris, Myiozetetes cayanensis, Myrmotherula assimilis, Numenius phaeopus, Nyctidromus albigularis, Oporornis agilis, Orthopsittacus manilata, Pachyrhamphus rufus (Boddaert 1783), Parkesia motacilla, Paroaria gularis, Passer domesticus, Petrochelidon pyrrhonota, Phaethornis anthophilus, Phaethornis superciliosus, Pionus menstruus, Pipreola chlorolepidota, Pluvialis dominica, Porphyrio flavirostris, Progne subis, Progne tapera, Protonotaria citrea, Riparia riparia, Sporophila caerulea, Sporophila lineola, Tachycineta albiventer (Boddaert 1783), Thalassidroma colymboides, Thalassidroma furcata, Threnetes ruckeri, Tinamus osgoodi, Touit stictoptera, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Tyrannus savana, Tyrannus tyrannus, Vireo altiloquus, Xiphorhynchus picus

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

directThreatsUnstructured: La deforestación en los Andes es la principal amenaza para esta especie (Rodríguez-Mahecha 2005; Hilty y Brown, 2001).

ancillaryData

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights:  
license:  
modified:  
created:  
mimeType:  
dataType:  
identifier:

#### measurementOrFact

relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

abstract: Perico de 17 a 18 cm de longitud. En general, la coloración de su cuerpo es verde, más clara en la zona del abdomen; las alas son color pardo con manchas blancas; la cola es color amarillo-oliva opaco por debajo con el borde azulado. La hembra se diferencia del macho por las plumas de la parte media-baja del ala que son verdes, la frente y el rostro son de color verde amarillento (Rodríguez-Mahecha 2005).

#### [associatedParty](#)

role:  
personnelIdentifier:  
personnelDirectory:  
homepage:  
email: edgarestebanhc@gmail.com  
phone:  
postalCode:  
country:  
state:  
city:  
address:  
organisation:  
position:  
lastName: Herrera Collazos  
firstName: Edgar Esteban

#### [ancillaryData](#)

dataObject:  
additionalInformation: AvianWeb

#### [reference](#)

geoPoint:  
location:  
thumbnailURL:  
mediaURL: <http://www.avianweb.com/spotwingedparrotlets.html>  
description:

## subject

source: <http://www.avianweb.com>

## audience

bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

## agent

contentType:  
dataType:  
identifier:

\_id

\_bsontype: ObjectID  
id: Vç"ÇfÄW Dă ð