

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56d4ac753c16479905cba8a1

Fecha de creación: 2016-02-29T20:39:19.496Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Clara Maria Pérez Gallego

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataObject:

identifier:

habitatUnstructured: Terrestre, en zonas húmedas a muy húmedas. Bordes de selva húmeda y muy húmeda, monte secundario y claros con árboles.

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

[ancillaryData](#)

type:

language: Inglés

name: Andean Emerald

creation_date: 2016-02-29T20:39:19.496Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Clara Maria Pérez Gallego
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:

fullDescriptionUnstructured: El largo total promedio de 9.1 cm, pico recto (23 mm), mandíbula inferior rosa con ápice moreno. Macho por encima verde brillante, coronilla azul violeta iridiscente; lados de cabeza y cuello verde iridiscente; partes inferiores totalmente blanco puro con lados y flancos verdes; supracaudales cobrizas; cola ligeramente ahorquillada verde bronceo con banda subterminal difusa morena y estrecho ápice pálido. Hembra similar pero coronilla verde (no azul).
briefDescription:

references

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:
providerLink:

publicationStatus

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

replacementNameFor

microReference:
source:
datatype:
identifier:
source:

datatype:
identifier:
note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

conservedAgainst

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion
typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:
canonicalAuthorship

authors

year

simple:
simple: (Bourcier & Mulsant, 1846)

canonicalName

infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:

ref:
uninomial:
simple: Amazilia franciae
rank: SPECIES

simple: Amazilia franciae (Bourcier & Mulsant, 1846)

attributes

nomenclaturalCode:
isAnamorphic:
id:

synonymsAtomized

ancillaryData

synonymStatus:
synonymName
anyTwo:

anyOne

providerLink:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:

identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typeVoucherEntity

typeOfType:
simple:
microReference:
year:
source:
datatype:
identifier:

[combinationAuthorship](#)

basionymAuthorship

simple:

authorship

simple:

simple:

epithet

infraspecificEpithet:

specificEpithet:

infragenericEpithet:

genus

linkType:

ref:

uninomial:

simple: Agyrtria franciae

rank: SPECIES

simple: Agyrtria franciae

nomenclaturalCode:

isAnamorphic:

id:

[hierarchy](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:

rightsHolder:
rights:
license:
title:
modified: 2016-02-29T20:38:59.495Z
created: 2016-02-29T20:38:59.495Z

agent

homepage:
role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:
parentTaxon: Amazilia
higherClassification:
infraspecificEpithet:
specificEpithet:
taxonRank: SPECIES
subGenus:
genus: Amazilia
family: Trochilidae
order: Apodiformes
classHierarchy: Aves
phylum: Chordata
kingdom: Animalia
recommended:
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Clara Maria Pérez Gallego
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:

dataType:
identifier:
reproductionUnstructured: No es muy conocida.

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Clara Maria Pérez Gallego
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
feedingUnstructured: Forrajea flores bajas y caza insectos en el follaje.

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Clara Maria Pérez Gallego
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:
created:
mimeType:
dataType:
identifier:
behaviorUnstructured: Diurno y arbóreo hacia el dosel. Usualmente solitario, rutero en flores dispersas a alturas variables; ocasionalmente se congregan en árboles en flor; también caza insectos al aire en cortos vuelos cerca del follaje. Los machos cantan desde perchas expuestas a 7 - 10 m.
Canto es una serie de chillidos fuertes, reminiscente del canto del Mielero común (*Coereba flaveola*), pero menos musical.

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

[endemicAtomized](#)

distribution

ancillaryData

dataObject:

additionalInformation:

reference

geoPoint:

location:

thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:

rightsHolder:

rights: Clara Maria Pérez Gallego

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

agent

contentType:

dataType:

identifier:

distributionUnstructured: Colombia, W Ecuador y N Perú. En el país en la C Occidental, C Central, cabeceras del valle del Magdalena en Huila y vertiente W de C Oriental en Cundinamarca; vertiente del Pacifico en Nariño, desde 1000 m - 2000 m. En la región Andina, Pacífica.

distributionAtomized

temporalCoverage

endDate:

startDate:

distributionScope

type:

threatStatus

ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES: Apéndice II

locality:

municipality:

county:
stateProvince:
country: Global
0: IUCN
measurementValue: Preocupación menor (LC)
relatedTo:
measurementRemarks:
measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementType: IUCN
measurementID:

territory

territoryUnstructured:
areaOfOccupancy:
extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:

measurementAccuracy:
measurementValue:
measurementType:
measurementID:
abstract:

associatedParty

role:
personnelIdentifier:
personnelDirectory:
homepage:
email: enataliagomezm@gmail.com
phone:
postalCode:
country: Colombia
state: Bogotá Distrito Capital
city: Bogotá D.C.
address:
organisation:
position:
lastName: Gómez Melendro
firstName: Edith Natalia

ancillaryData

dataObject:
additionalInformation: Imagen: Amazilia_franciae.jpg

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder: Clara Maria Pérez Gallego
rights:
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

contentType:
dataType:
identifier:

_id

_bsontype: ObjectId

id: VÔ¬u< G™\°Š