

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56ca1d87f0106c67230e7456

Fecha de creación: undefined

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

0

isbn: 13: 978-0-7922-5314-3

keywords: Butorides virescens,Empidonax alnorum,Empidonax traillii,Geothlypis philadelphia,Leiothlypis peregrina,Leucophaeus atricilla,Myiarchus crinitus,Patagioenas fasciata,Petrochelidon pyrrhonota,Setophaga castanea,Setophaga striata,Spiza americana

institution:

address:

publisher: the National Geographic Society

accessed:

websites:

chapter:

series:

pages: 503

issue:

year: 2006

0: Dunn, J

1: Alderfer, J

title: Field guide to the birds of North America. Fifth Edition.

source:

type: book

tags: Butorides virescens,Empidonax alnorum,Empidonax traillii,Geothlypis philadelphia,Leiothlypis peregrina,Leucophaeus atricilla,Myiarchus crinitus,Patagioenas fasciata,Petrochelidon pyrrhonota,Setophaga castanea,Setophaga striata,Spiza americana

abstract:

last_modified:

created:

group_id:

profile_id:

id:

1

link: <http://eol.org/pages/1050919/overview>

keywords: Leiothlypis peregrina

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

0: EOL

title: Leiothlypis peregrina

source: Encyclopedia of life
type: miscellany
tags: *Leiothlypis peregrina*
abstract:
last_modified:
created:
group_id:
profile_id:
id:

2

keywords: *Accipiter striatus*, *Actitis macularius*, *Adelomyia melanogenys*, *Agelaiocercus kingii*, *Amblycercus holosericeus*, *Ampelion rubrocristatus*, *Anas andium*, *Anas clypeata*, *Anas discors*, *Andigena nigrirostris*, *Anisognathus igniventris*, *Ardea alba*, *Ardea herodias*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Asthenes flammulata*, *Asthenes fuliginosa*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Butorides virescens*, *Cairina moschata*, *Calidris bairdii*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Cathartes aura*, *Catharus minimus*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chaetura pelagica*, *Chalcostigma heteropogon*, *Chordeiles minor*, *Chrysomus icterocephalus*, *Cinclodes albidiventris*, *Cinclus leucocephalus*, *Cinnycerthia unirufa*, *Cistothorus apolinari*, *Cistothorus platensis*, *Coccyzua pumila*, *Coccyzus americanus*, *Coccyzus melacoryphus*, *Coeligena bonapartei*, *Colibri coruscans*, *Colibri thalassinus*, *Columba livia*, *Columbina talpacoti*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cinereus*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Dendrocygna autumnalis*, *Dendrocygna viduata*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dolichonyx oryzivorus*, *Dubusia taeniata*, *Egretta caerulea*, *Egretta thula*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Gallinula melanops*, *Geothlypis philadelphia*, *Geranoaetus melanoleucus*, *Glaucidium jardinii*, *Grallaria quitesis*, *Grallaria ruficapilla*, *Grallaria squamigera*, *Haplospiza rustica*, *Heliangelus amethysticollis*, *Heliangelus exortis*, *Hellmayrea gularis*, *Hemispingus superciliaris*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus icterus*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Lafresnaya lafresnayi*, *Leiothlypis peregrina*, *Leptasthenura andicola*, *Lesbia nuna*, *Lesbia victoriae*, *Machetornis rixosa*, *Margarornis squamiger*, *Mecocerculus leucophrys*, *Megascops choliba*, *Melanerpes rubricapillus*, *Merganetta armata*, *Metallura tyrianthina*, *Milvago chimachima*, *Mimus gilvus*, *Mniotilta varia*, *Molothrus bonariensis*, *Muscisaxicola alpinus*, *Myiarchus crinitus*, *Myiodynastes luteiventris*, *Myiotheretes fumigatus*, *Myiotheretes striaticollis*, *Myiothlypis nigrocristata*, *Nomonyx dominicus*, *Nycticorax nycticorax*, *Ochthoeca cinnamomeiventris*, *Ochthoeca diadema*, *Ochthoeca fumicolor*, *Ochthoeca rufipectoralis*, *Orochelidon murina*, *Oxypogon guerinii*, *Oxyura jamaicensis*, *Pachyramphus versicolor*, *Pandion haliaetus*, *Pardirallus maculatus*, *Parkesia noveboracensis*, *Patagioenas fasciata*, *Petrochelidon pyrrhonota*, *Phaetusa simplex*, *Phalacrocorax brasilianus*, *Pheucticus aureoventris*, *Pheucticus ludovicianus*, *Phimosus infuscatus*, *Phrygilus unicolor*, *Phyllomyias uropygialis*, *Piaya cayana*, *Picoides fumigatus*, *Pipraeidea melanonota*, *Pipreola riefferii*, *Piranga olivacea*, *Piranga rubra*, *Pitangus sulphuratus*, *Pluvialis dominica*, *Podilymbus podiceps*, *Porphyrio martinicus*, *Porzana carolina*, *Progne tapera*, *Protonotaria citrea*, *Pseudocolopteryx acutipennis*, *Pseudoscops clamator*, *Pterophanes cyanopterus*, *Pygochelidon cyanoleuca*, *Pyrocephalus rubinus*, *Pyrrhomias cinnamomeus*, *Quiscalus lugubris*, *Rallus semiplumbeus*, *Ramphocelus dimidiatus*, *Ramphomicron microrhynchum*, *Riparia riparia*, *Rupornis magnirostris*, *Sayornis nigricans*, *Serpophaga cinerea*, *Setophaga castanea*, *Setophaga cerulea*, *Setophaga petechia*, *Setophaga ruticilla*, *Setophaga striata*, *Sicalis flaveola*, *Sicalis luteola*, *Spiza americana*, *Sporagra spinescens*, *Sturnella magna*, *Synallaxis subpudica*, *Systemellura longirostris*, *Tangara vitriolina*, *Thraupis episcopus*, *Thraupis palmarum*, *Tringa flavipes*, *Tringa melanoleuca*, *Tringa solitaria*, *Troglodytes aedon*, *Troglodytes solstitialis*, *Turdus fuscater*, *Turdus ignobilis*, *Tyrannus melancholicus*, *Tyrannus savana*, *Tyrannus tyrannus*, *Tyto alba*, *Vanellus chilensis*, *Vireo olivaceus*, *Vultur gryphus*, *Zenaida auriculata*, *Zonotrichia capensis*
institution:
address:
publisher: Princeton University Press
accessed:

websites:

chapter:

series:

pages: 836

issue:

year: 1986

0: Hilty, Steven

1: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

[illegible]

abstract:

last_modified:

created:

group_id:

profile_id:

id:

3

link: http://awsassets.panda.org/downloads/migratorias{_}aves{_}42{_}final.pdf

keywords: *Leiothlypis peregrina*

0: Naranjo, L G

1: Amaya, J D

2: Eusse-González, D

3: Cifuentes-Sarmiento, Y

institution:

address:

publisher: Ministerio de Ambiente y Desarrollo Sostenible / WWF Colombia

accessed:

websites:

chapter:

series:

pages: 708

issue:

volume: 1

year: 2012

0: Peraza, Camilo

title: *Leiothlypis peregrina*

source: Guía de las Especies Migratorias de la Biodiversidad en Colombia Aves

type: book_section

tags: *Leiothlypis peregrina*

abstract:

last_modified:

created:

group_id:

profile_id:

id:

4

keywords: *Actitis macularius*, *Ampelion rubrocristatus*, *Anas discors*, *Anisognathus igniventris*, *Ardea alba*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bartramia longicauda*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Catharus fuscescens*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chrysomus icterocephalus*, *Cinnycerthia unirufa*, *Cistothorus apolinari*, *Coccyzus americanus*, *Coeligena bonapartei*, *Colibri coruscans*, *Columba livia*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cooperi*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dubusia taeniata*, *Egretta caerulea*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Gallinula melanops*, *Geothlypis philadelphia*, *Glaucidium jardinii*, *Grallaria ruficapilla*, *Hemispingus supercilialis*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Leiothlypis peregrina*

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D
institution:
edition: Primera Ed
address:
publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca
accessed:
websites:
chapter:
series:
pages: 276
issue:
year: 2000

0: Asociación Bogotana de Ornitología
title: Aves de la Sabana de Bogotá: guía de campo
source:

type: book
tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccyzus americanus, Coeligena bonapartei, Colibri coruscans, Columba livia, Conirostrum rufum, Conirostrum sitticolor, Contopus cooperi, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dubusia taeniata, Egretta caerulea, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Glaucidium jardinii, Grallaria ruficapilla, Hemispingus supercilialis, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Leiothlypis peregrina

abstract:
last_modified:

created:
group_id:
profile_id:
id:

geoPoint:
location:
thumbnailURL:

description:
source:
bibliographicCitation:
rightsHolder:

rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:

created:
mimeType:
dataType:
identifier:

habitatUnstructured: En sus zonas de reproducción habita en bosques caducifolios, coníferas, alisos y sauce (Dunn & Aldefer, 2006). Cuando reside en las épocas de invierno es posible registrarla en áreas selváticas, montes claros, cultivos de café (EOL, n.d.; Hilty & Brown, 1986), principalmente piedemonte y montañas (Peraza, 2012). En Colombia esta especie es común en la Sierra Nevada de Santa Marta, muy dispersa por los Andes en áreas boscosas (Hilty & Brown, 1986). En la Sabana de Bogotá frecuenta árboles nativos e introducidos en los parques, bordes de ríos, cercas vivas, bordes de cultivos, potreros en fincas y áreas residenciales (Asociación Bogotana de

Ornitología, 2000; Peraza, 2012).

[commonNamesAtomized](#)

[ancillaryData](#)

dataObject:
additionalInformation:

[reference](#)

geoPoint:
location:
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

[agent](#)

contentType:
dataType:
identifier:
usedBy:

usedIn

distributionUnstructured: Colombia
endDate:
startDate:

[ancillaryData](#)

type:
language: Español
name: Reinita Verdilla

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

0

link: http://awsassets.panda.org/downloads/migratorias{_}aves{_}42{_}final.pdf

keywords: *Leiothlypis peregrina*

0: Naranjo, L G

1: Amaya, J D

2: Eusse-González, D

3: Cifuentes-Sarmiento, Y

institution:

address:

publisher: Ministerio de Ambiente y Desarrollo Sostenible / WWF Colombia

accessed:

websites:

chapter:

series:

pages: 708

issue:

volume: 1

year: 2012

0: Peraza, Camilo

title: *Leiothlypis peregrina*

source: Guía de las Especies Migratorias de la Biodiversidad en Colombia Aves

type: book_section

tags: *Leiothlypis peregrina*

abstract:

last_modified:

created:

group_id:

profile_id:

id:

1

link: <http://www.museum.lsu.edu/{~}Remsen/SACCprop571.htm>

keywords: *Cardellina canadensis*, *Geothlypis philadelphia*, *Leiothlypis peregrina*, *Myiothlypis nigrocristata*, *Setophaga castanea*, *Setophaga cerulea*, *Setophaga fusca*, *Setophaga petechia*, *Setophaga striata*

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2012

0: Banks, R

title: Proposal (571) to South American Classification Committee

source: South American Classification Committee

type: miscellany

tags: *Cardellina canadensis*, *Geothlypis philadelphia*, *Leiothlypis peregrina*, *Myiothlypis nigrocristata*, *Setophaga*

castanea,Setophaga cerulea,Setophaga fusca,Setophaga petechia,Setophaga striata

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

fullDescriptionUnstructured: La Reinita Verdilla mide aproximadamente 12 cm; el plumaje de invierno es muy parecido entre los sexos, verde oliva amarillento por encima, con línea superciliar amarillo opaco con blanco y línea ocular negruzca. Márgenes contrastantes verde oliva sobre fondo oscuro en las primarias. Partes inferiores amarillo opaco con blanco en parte baja del abdomen y la cola. Hacia febrero comienzan a mudar el plumaje (Peraza, 2012). El plumaje reproductivo del macho es gris en la parte superior de la cabeza y resto de las partes verde oliva brillante. Márgenes contrastantes verde oliva brillante sobre fondo oscuro en las primarias. Tenue barra blanca en el ala en muchos individuos, blanco por debajo. Las hembras son similares, pero sin gris en la cabeza y con ceja amarillenta. Tonos verde amarillentos en pecho y parte del abdomen (Peraza, 2012). Anteriormente Vermivora peregrina (Wilson, A, 1811). Cambió de acuerdo con la propuesta (571) del South American Classification Committee (Banks, 2012).

briefDescription:

references

isbn: 978-0-679-45122-8

keywords: Anas clypeata,Ardea herodias,Buteo swainsoni,Cairina moschata,Crotophaga ani,Dendrocygna autumnalis,Dendrocygna viduata,Egretta caerulea,Egretta thula,Empidonax alnorum,Geothlypis philadelphia,Ixobrychus exilis,Leiothlypis peregrina,Myiodynastes luteiventris,Nomonyx dominicus,Parkesia noveboracensis,Petrochelidon pyrrhonota,Pheucticus ludovicianus,Protonotaria citrea,Sayornis nigricans

editors

institution:

edition: 1 st ed.

address:

publisher: National Audubon Society

accessed:

websites:

chapter:

series:

pages: 545

issue:

year: 2000

authors

authors: Sibley, D A
title: The SIBLEY Guide to Birds
source:
type: book
tags: Anas clypeata,Ardea herodias,Buteo swainsoni,Cairina moschata,Crotophaga ani,Dendrocygna autumnalis,Dendrocygna viduata,Egretta caerulea,Egretta thula,Empidonax alnorum,Geothlypis philadelphia,Ixobrychus exilis,Leiothlypis peregrina,Myiodynastes luteiventris,Nomonyx dominicus,Parkesia noveboracensis,Petrochelidon pyrrhonota,Pheucticus ludovicianus,Protonotaria citrea,Sayornis nigricans
abstract:

identifiers

last_modified:
created:
group_id:
profile_id:
id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:
providerLink:

publicationStatus

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

replacementNameFor

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:

identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

conservedAgainst

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:

canonicalAuthorship

authors

year

simple:
simple: (A. Wilson, 1811)

canonicalName

infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:
ref:
uninomial:
simple: Leiothlypis peregrina
rank: SPECIES
simple: Leiothlypis peregrina (A. Wilson, 1811)

attributes

nomenclaturalCode:
isAnamorphic:
id:

synonymsAtomized

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder:
rights:
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

mimeType:
dataType:
identifier:
synonymStatus:

synonymName

anyTwo:

anyOne

providerLink:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:

datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:

note:
ruleConsidered:

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typeVoucherEntity

typeOfType:
simple:
microReference:
year:
source:
datatype:
identifier: (Wilson, A, 1811)

[combinationAuthorship](#)

basionymAuthorship

simple:

authorship

simple:
simple:

epithet

infraspecificEpithet:
specificEpithet:
infragenericEpithet:

genus

linkType:
ref:
uninomial:
simple: Vermivora peregrina
rank:
simple:
nomenclaturalCode:
isAnamorphic:
id:

[hierarchy](#)

[ancillaryData](#)

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder:
rights:
license:
title:
modified: 2016-02-21T20:20:55.548Z
created: 2016-02-21T20:20:55.548Z

agent

homepage:
role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:
parentTaxon: Leiothlypis
higherClassification:
infraspecificEpithet:
specificEpithet:
taxonRank: SPECIES
subGenus:
genus: Leiothlypis
family: Parulidae
order: Passeriformes
classHierarchy: Aves
phylum: Chordata
kingdom: Animalia
recommended:
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:
additionalInformation:

0

link: <http://eol.org/pages/1050919/overview>

keywords: *Leiothlypis peregrina*

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

0: EOL

title: *Leiothlypis peregrina*

source: Encyclopedia of life

type: miscellany

tags: *Leiothlypis peregrina*

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

reproductionUnstructured: Al igual que otras especies de reinitas, la Reinita Verdilla es monógama. El nido es una copa pequeña, construido de musgo, que por lo general se ubica en la parte inferior de arbustos o árboles pequeños. La incubación dura entre once a doce días; depositan entre cuatro a seis huevos blancos, algunas veces manchados con castaño o marcas púrpura pálido (EOL, n.d.).

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:

additionalInformation:

0

link: <http://eol.org/pages/1050919/overview>

keywords: *Leiothlypis peregrina*

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

0: EOL

title: *Leiothlypis peregrina*

source: Encyclopedia of life

type: miscellany

tags: *Leiothlypis peregrina*

abstract:

last_modified:

created:

group_id:

profile_id:

id:

1

keywords: *Actitis macularius*, *Ampelion rubrocristatus*, *Anas discors*, *Anisognathus igniventris*, *Ardea alba*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bartramia longicauda*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Catharus fuscescens*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chrysomus icterocephalus*, *Cinnycerthia unirufa*, *Cistothorus apolinari*, *Coccyzus americanus*, *Coeligena bonapartei*, *Colibri coruscans*, *Columba livia*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cooperi*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dubusia taeniata*, *Egretta caerulea*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Gallinula melanops*, *Geothlypis philadelphia*, *Glaucidium jardinii*, *Grallaria ruficapilla*, *Hemispingus supercilialis*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Leiothlypis peregrina*

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccyzus americanus, Coeligena bonapartei, Colibri coruscans, Columba livia, Conirostrum rufum, Conirostrum sitticolor, Contopus cooperi, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dubusia taeniata, Egretta caerulea, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Glaucidium jardinii, Grallaria ruficapilla, Hemispingus supercilialis, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Leiothlypis peregrina

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

feedingUnstructured: Es una especie omnívora; se alimenta de pequeños insectos; también consume frutos pequeños y néctar, especialmente de las flores de los eucaliptos (Asociación Bogotana de Ornitología, 2000; EOL,

n.d.). También se ha visto en comederos alimentándose de los plátanos (EOL, n.d.).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

0

keywords: *Actitis macularius*, *Ampelion rubrocristatus*, *Anas discors*, *Anisognathus igniventris*, *Ardea alba*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bartramia longicauda*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Catharus fuscescens*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chrysomus icterocephalus*, *Cinnycerthia unirufa*, *Cistothorus apolinari*, *Coccyzus americanus*, *Coeligena bonapartei*, *Colibri coruscans*, *Columba livia*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cooperi*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dubusia taeniata*, *Egretta caerulea*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Gallinula melanops*, *Geothlypis philadelphia*, *Glaucidium jardinii*, *Grallaria ruficapilla*, *Hemispingus supercilialis*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Leiothlypis peregrina*

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:
year: 2000
0: Asociación Bogotana de Ornitología
title: Aves de la Sabana de Bogotá: guía de campo
source:
type: book
tags: *Actitis macularius*, *Ampelion rubrocristatus*, *Anas discors*, *Anisognathus igniventris*, *Ardea alba*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bartramia longicauda*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Catharus fuscescens*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chrysomus icterocephalus*, *Cinnycerthia unirufa*, *Cistothorus apolinari*, *Coccyzus americanus*, *Coeligena bonapartei*, *Colibri coruscans*, *Columba livia*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cooperi*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dubusia taeniata*, *Egretta caerulea*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Gallinula melanops*, *Geothlypis philadelphia*, *Glaucidium jardinii*, *Grallaria ruficapilla*, *Hemispingus superciliaris*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Leiothlypis peregrina*
abstract:
last_modified:
created:
group_id:
profile_id:
id:

1

link: http://awsassets.panda.org/downloads/migratorias{_}aves{_}42{_}final.pdf
keywords: *Leiothlypis peregrina*
0: Naranjo, L G
1: Amaya, J D
2: Eusse-González, D
3: Cifuentes-Sarmiento, Y
institution:
address:
publisher: Ministerio de Ambiente y Desarrollo Sostenible / WWF Colombia
accessed:
websites:
chapter:
series:
pages: 708
issue:
volume: 1
year: 2012
0: Peraza, Camilo
title: *Leiothlypis peregrina*
source: Guía de las Especies Migratorias de la Biodiversidad en Colombia Aves
type: book_section
tags: *Leiothlypis peregrina*
abstract:
last_modified:
created:
group_id:
profile_id:

id:

2

isbn: 0-618-57424-7

keywords: *Anas clypeata*, *Anas platyrhynchos*, *Ardea herodias*, *Bartramia longicauda*, *Buteo swainsoni*, *Cairina moschata*, *Calidris melanotos*, *Columbina talpacoti*, *Contopus cooperi*, *Contopus virens*, *Crotophaga ani*, *Dendrocygna autumnalis*, *Dolichonyx oryzivorus*, *Egretta caerulea*, *Egretta thula*, *Empidonax alnorum*, *Empidonax traillii*, *Falco columbarius*, *Falco sparverius*, *Geothlypis philadelphia*, *Hirundo rustica*, *Leiothlypis peregrina*, *Myiarchus crinitus*, *Myiodynastes luteiventris*, *Nomonyx dominicus*, *Nycticorax nycticorax*, *Pandion haliaetus*, *Parkesia noveboracensis*, *Petrochelidon pyrrhonota*, *Podilymbus podiceps*, *Protonotaria citrea*, *Sayornis nigricans*

institution:

edition: Hillstar E

address:

publisher:

accessed:

websites:

chapter:

series:

pages: 391

issue:

year: 2005

O: Kaufman, K

title: Guía de campo a las aves de Norteamérica

source:

type: book

tags: *Anas clypeata*, *Anas platyrhynchos*, *Ardea herodias*, *Bartramia longicauda*, *Buteo swainsoni*, *Cairina moschata*, *Calidris melanotos*, *Columbina talpacoti*, *Contopus cooperi*, *Contopus virens*, *Crotophaga ani*, *Dendrocygna autumnalis*, *Dolichonyx oryzivorus*, *Egretta caerulea*, *Egretta thula*, *Empidonax alnorum*, *Empidonax traillii*, *Falco columbarius*, *Falco sparverius*, *Geothlypis philadelphia*, *Hirundo rustica*, *Leiothlypis peregrina*, *Myiarchus crinitus*, *Myiodynastes luteiventris*, *Nomonyx dominicus*, *Nycticorax nycticorax*, *Pandion haliaetus*, *Parkesia noveboracensis*, *Petrochelidon pyrrhonota*, *Podilymbus podiceps*, *Protonotaria citrea*, *Sayornis nigricans*

abstract:

last_modified:

created:

group_id:

profile_id:

id:

3

isbn: 978-0-679-45122-8

keywords: *Anas clypeata*, *Ardea herodias*, *Buteo swainsoni*, *Cairina moschata*, *Crotophaga ani*, *Dendrocygna autumnalis*, *Dendrocygna viduata*, *Egretta caerulea*, *Egretta thula*, *Empidonax alnorum*, *Geothlypis philadelphia*, *Ixobrychus exilis*, *Leiothlypis peregrina*, *Myiodynastes luteiventris*, *Nomonyx dominicus*, *Parkesia noveboracensis*, *Petrochelidon pyrrhonota*, *Pheucticus ludovicianus*, *Protonotaria citrea*, *Sayornis nigricans*

institution:

edition: 1 st ed.

address:

publisher: National Audubon Society

accessed:

websites:

chapter:

series:

pages: 545

issue:

year: 2000

contentType:

dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

[endemicAtomized](#)

[distribution](#)

[ancillaryData](#)

dataObject:
additionalInformation:

[reference](#)

link: http://awsassets.panda.org/downloads/migratorias{_}aves{_}42{_}final.pdf
keywords: Leiothlypis peregrina

[editors](#)

editors: Naranjo, L G
editors: Amaya, J D
editors: Eusse-González, D
editors: Cifuentes-Sarmiento, Y
institution:
address:
publisher: Ministerio de Ambiente y Desarrollo Sostenible / WWF Colombia
accessed:
websites:
chapter:
series:
pages: 708
issue:
volume: 1
year: 2012

[authors](#)

authors: Peraza, Camilo
title: Leiothlypis peregrina
source: Guía de las Especies Migratorias de la Biodiversidad en Colombia Aves
type: book_section
tags: Leiothlypis peregrina
abstract:

[identifiers](#)

last_modified:

created:
group_id:
profile_id:
id:
geoPoint:
location:
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

[agent](#)

contentType:
dataType:
identifier:
distributionUnstructured: Esta especie se distribuye desde el sur de Canadá (Alberta, Manitoba, sur de Ontario, Quebec y New Brunswick) y noroeste de Estados Unidos (Maine); inverte en Florida (Estados Unidos) y las Antillas Mayores hasta Jamaica. Presente en todos los países de Centroamérica desde el sur de México, Colombia y región noroccidental de Venezuela, Aruba, Curazao y Bonaire. Algunos registros en el norte de Ecuador y Guyana Francesa (Peraza, 2012). La distribución en Colombia viene desde límites con Panamá, por las tierras bajas del Caribe, regiones Pacífica (con excepción de suroccidente de Nariño) y Andina. En la cordillera Oriental hasta el Piedemonte llanero (Peraza, 2012).
Principalmente hasta los 2600 msnm (Sabana de Bogotá), rara vez a mayores elevaciones (Asociación Bogotana de Ornitología, 2000; Peraza, 2012).

[distributionAtomized](#)

stateProvince: Córdoba

[ancillaryData](#)

locality:
municipality:
country: Colombia

temporalCoverage

endDate:
startDate:

distributionScope

type:

[threatStatus](#)

ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:

locality:

municipality:

county:

stateProvince:

country: Global

0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementType: IUCN

measurementID:

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

0

dataObject:

additionalInformation:

0

link: <http://maps.iucnredlist.org/map.html?id=22721621>

keywords: Leiothlypis peregrina

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2014

0: BirdLife International

1: NatureServe

title: Bird Species Distribution Maps of the World. 2012. Vermivora peregrina. The IUCN Red List of Threatened Species. Version 2014.3

source:

type: miscellany

tags: Leiothlypis peregrina

abstract:
last_modified:
created:
group_id:
profile_id:
id:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
populationBiologyUnstructured: La Reinita Verdilla tiene un amplio rango de distribución, la tendencia de la población es estable por lo cual se encuentra en categoría de Preocupación Menor (BirdLife International and NatureServe, 2014).

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: La Reinita Verdilla mide aproximadamente 12 cm; es una especie pequeña, activa, de pico fino y colores opacos. Su plumaje verde oliva por encima, blanco teñido de amarillo opaco o verdoso por debajo; una ceja fina amarillenta o blanquecina y una línea ocular oscura son las únicas marcas distintivas; a partir de febrero adquiere su plumaje de cría con la coronilla más gris y la ceja más blanca, especialmente en los machos, los cuales son más blancos por debajo. Pico color cuerno oscuro, patas grises (Asociación Bogotana de Ornitología, 2000). Es una especie simple, con un ruidoso y titubeante canto; pasa el verano en pantanos y claros en los bosques del norte; el invierno principalmente en Centro y Sudamérica (Kaufman, 2005). Es posible verla en grupos mixtos, forrajeando activamente entre el follaje (Hilty & Brown, 1986). Es una especie omnívora; se alimenta de pequeños insectos; también consume frutos pequeños y néctar, especialmente de las flores de los eucaliptos (Asociación Bogotana de Ornitología, 2000; EOL, n.d.). También se ha visto en comederos alimentándose de los plátanos (EOL, n.d.). La Reinita Verdilla tiene un amplio rango de distribución, la tendencia de la población es estable por lo cual se encuentra en categoría de Preocupación Menor (BirdLife International and NatureServe, 2014).

associatedParty

role:

personnelIdentifier:

personnelDirectory:

homepage:

email: enataliagomez@gmail.com

phone:

postalCode:

country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address:

organisation:

position:

lastName: Gómez Melendro

firstName: Edith Natalia

ancillaryData

dataObject:

additionalInformation:

reference

geoPoint:

location:

thumbnailURL:

mediaURL: <http://avibase.bsc-eoc.org/species.jsp?lang=ES&avibaseid=D204A9309E8CCC3A>

description:

subject

source: <http://avibase.bsc-eoc.org>

audience

bibliographicCitation:

rightsHolder:

rights:

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

agent

contentType:
dataType:
identifier:

_id

_bsonType: ObjectId

id: VÊ ¢ð lg#çE`