Ficha ID: 56c26f58f0106c67230e732c Fecha de creación: undefined annualCycles annualCycleUnstructured: habitats 0 dataObject: additionalInformation: keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos 0: Stiles. F G 1: Bohórquez, C I 2: Cadena, C D 3: De la Zerda, S 4: Hernández, M 5: Roselli. L 6: Kelsey, M 7: Valencia, I D 8: Knapp, D institution: edition: Primera Ed address: publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca accessed: websites: chapter: series: pages: 276 issue: year: 2000 0: Asociación Bogotana de Ornitología title: Aves de la Sabana de Bogotá: guía de campo source: type: book tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos abstract: last_modified: created: group_id: profile_id: id: keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus

Reporte de Fichas para el usuario: edith gómez

keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo

swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis.Catamenia analis.Catamenia inornata.Cathartes aura.Catharus minimus.Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus.Contopus fumigatus.Contopus sordidulus.Contopus virens.Coragyps atratus.Crotophaga ani.Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis. Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes Iuteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax, nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina,Oxypogon guerinii,Oxyura jamaicensis,Pachyramphus versicolor,Pandion haliaetus,Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis institution:

address:

publisher: Princeton University Press

accessed: websites: chapter: series: pages: 836 issue: year: 1986 0: Hilty, Steven 1: Brown, William

title: A Guide to the Birds of Colombia

source: type: book

tags: Accipiter striatus,Actitis macularius,Adelomyia melanogenys,Aglaiocercus kingii,Amblycercus holosericeus,Ampelion rubrocristatus,Anas andium,Anas clypeata,Anas discors,Andigena nigrirostris,Anisognathus igniventris,Ardea alba,Ardea herodias,Arremon assimilis,Asio flammeus,Asio stygius,Asthenes flammulata,Asthenes fuliginosa,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Butorides virescens,Cairina moschata,Calidris bairdii,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Cathartes aura,Catharus minimus,Catharus

ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis.Diglossa lafresnavii.Diglossa sittoides.Dolichonyx oryzivorus.Dubusia taeniata.Egretta thula.Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes Iuteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina,Oxypogon guerinii,Oxyura jamaicensis,Pachyramphus versicolor,Pandion haliaetus,Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris. Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis abstract:

last modified:

created:

group_id:

profile id:

id:

isbn: 0963699636

keywords: Anas clypeata, Anas platyrynchos, Ardea herodias, Astragalinus psaltria, Bartramia longicauda, Buteo swainsoni, Cairina moschata, Calidris melanotos, Cardellina canadensis, Cathartes aura, Catharus minimus, Coccyzus americanus, Columbina talpacoti, Contopus cooperi, Contopus virens, Coragyps atratus, Crotophaga ani, Dendrocygna autumnalis, Egretta caerulea, Egretta thula, Empidonax alnorum, Empidonax traillii, Falco columbarius, Fulica americana, Geothlypis philadelphia, Hirundo rustica, Ixobrychus exilis, Jacana jacana, Mniotilta varia, Nycticorax nycticorax, Oxyura jamaicensis, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Phalacrocorax brasilianus, Piranga olivacea, Pitangus sulphuratus, Podilymbus podiceps, Porzana carolina, Pyrocephalus rubinus, Rupornis magnirostris, Setophaga fusca, Sturnella magna, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon

0: Alderfer, Jonathan

institution: address:

publisher: Washington D.C. National Geographic Society

accessed: websites: chapter: series: pages: 664 issue: year: 2006 0: Alderfer, J 1: Zickefoose, J 2: Littlehales, B

title: National Geographic Complete Birds of North America

source: type: book

tags: Anas clypeata,Anas platyrynchos,Ardea herodias,Astragalinus psaltria,Bartramia longicauda,Buteo swainsoni,Cairina moschata,Calidris melanotos,Cardellina canadensis,Cathartes aura,Catharus minimus,Coccyzus americanus,Columbina talpacoti,Contopus cooperi,Contopus virens,Coragyps atratus,Crotophaga ani,Dendrocygna autumnalis,Egretta caerulea,Egretta thula,Empidonax alnorum,Empidonax traillii,Falco columbarius,Fulica americana,Geothlypis philadelphia,Hirundo rustica,Ixobrychus exilis,Jacana jacana,Mniotilta varia,Nycticorax nycticorax,Oxyura jamaicensis,Pandion haliaetus,Pardirallus maculatus,Parkesia noveboracensis,Phalacrocorax brasilianus,Piranga olivacea,Pitangus sulphuratus,Podilymbus podiceps,Porzana carolina,Pyrocephalus rubinus,Rupornis magnirostris,Setophaga fusca,Sturnella magna,Tringa flavipes,Tringa melanoleuca,Tringa solitaria,Troglodytes aedon

abstract: last_modified: created: group_id: profile_id:

id:

isbn: 958 33-2118-4

keywords: Bartramia longicauda, Calidris bairdii, Calidris melanotos

institution:

edition: Primera ed

address:

accessed:

publisher: American Bird Conservancy, WWF-US, Humedales para las Américas y Manomet Conservation Science,

Asociación Calidris

websites: chapter: series: pages: 141 issue: year: 2001 0: Canevari, P 1: Castro, G 2: Sallaberry, M 3: Naranjo, L

title: Guía de los Chorlos y Playeros de la Región Neotropical

source: type: book

tags: Bartramia longicauda, Calidris bairdii, Calidris melanotos

abstract: last_modified: created:

group_id: profile_id: id: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Asociación Bogotana de Ornitología license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: habitatUnstructured: Esta especie forrajea en pequeños grupos en bancos de lodo con pasto esparcido, charcos en potreros anegados, praderas, marismas, campos inundados, áreas abiertas húmedas y bordes de humedales, ocasionalmente se observa en playas y planos lodosos de la costa, ríos y lagunas (Asociación Bogotana de Ornitología, 2000; Hilty & Brown, 1986). Las zonas de anidación son la tundra y por lo general en las franjas de vegetación alrededor de los humedales (Alderfer et al., 2006; Canevari et al., 2001). commonNamesAtomized
ancillaryData
usedBy:
usedIn
distributionUnstructured: endDate: startDate:
ancillaryData
type: language: Polaco name: Biegus arktyczny
usesManagementAndConservation
ancillaryData
managementAndConservationUnstructured:
managementAndConservationAtomized
identificationKeys
fullDescription
0
dataObject: additionalInformation:

isbn: 958 33-2118-4

keywords: Bartramia longicauda, Calidris bairdii, Calidris melanotos

institution:

edition: Primera ed

address:

publisher: American Bird Conservancy, WWF-US, Humedales para las Américas y Manomet Conservation Science,

Asociación Calidris

accessed:
websites:
chapter:
series:
pages: 141
issue:
year: 2001
0: Canevari, P
1: Castro, G
2: Sallaberry, M
3: Naranjo, L

title: Guía de los Chorlos y Playeros de la Región Neotropical

source: type: book

tags: Bartramia longicauda, Calidris bairdii, Calidris melanotos

abstract: last_modified: created: group_id: profile_id: id:

-

isbn: 0963699636

keywords: Anas clypeata, Anas platyrynchos, Ardea herodias, Astragalinus psaltria, Bartramia longicauda, Buteo swainsoni, Cairina moschata, Calidris melanotos, Cardellina canadensis, Cathartes aura, Catharus minimus, Coccyzus americanus, Columbina talpacoti, Contopus cooperi, Contopus virens, Coragyps atratus, Crotophaga ani, Dendrocygna autumnalis, Egretta caerulea, Egretta thula, Empidonax alnorum, Empidonax traillii, Falco columbarius, Fulica americana, Geothlypis philadelphia, Hirundo rustica, Ixobrychus exilis, Jacana jacana, Mniotilta varia, Nycticorax nycticorax, Oxyura jamaicensis, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Phalacrocorax brasilianus, Piranga olivacea, Pitangus sulphuratus, Podilymbus podiceps, Porzana carolina, Pyrocephalus rubinus, Rupornis magnirostris, Setophaga fusca, Sturnella magna, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon

0: Alderfer, Jonathan

institution: address:

publisher: Washington D.C. National Geographic Society

accessed: websites: chapter: series: pages: 664 issue: year: 2006 0: Alderfer, J 1: Zickefoose, J

2: Littlehales, B

title: National Geographic Complete Birds of North America

source: type: book

tags: Anas clypeata, Anas platyrynchos, Ardea herodias, Astragalinus psaltria, Bartramia longicauda, Buteo swainsoni, Cairina moschata, Calidris melanotos, Cardellina canadensis, Cathartes aura, Catharus minimus, Coccyzus americanus, Columbina talpacoti, Contopus cooperi, Contopus virens, Coragyps atratus, Crotophaga ani, Dendrocygna autumnalis.Egretta caerulea.Egretta thula.Empidonax alnorum.Empidonax traillii.Falco columbarius.Fulica americana, Geothlypis philadelphia, Hirundo rustica, Ixobrychus exilis, Jacana jacana, Mniotilta varia, Nycticorax nycticorax,Oxyura jamaicensis,Pandion haliaetus,Pardirallus maculatus,Parkesia noveboracensis,Phalacrocorax brasilianus, Piranga olivacea, Pitangus sulphuratus, Podilymbus podiceps, Porzana carolina, Pyrocephalus rubinus, Rupornis magnirostris, Setophaga fusca, Sturnella magna, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon

abstract: last modified: created: group id: profile id:

isbn: 0-618-57424-7

keywords: Anas clypeata, Anas platyrynchos, Ardea herodias, Bartramia longicauda, Buteo swainsoni, Cairina moschata, Calidris melanotos, Columbina talpacoti, Contopus cooperi, Contopus virens, Crotophaga ani, Dendrocygna autumnalis, Dolichonyx oryzivorus, Egretta caerulea, Egretta thula, Empidonax alnorum, Empidonax traillii, Falco columbarius, Falco sparverius, Geothlypis philadelphia, Hirundo rustica, Leiothlypis peregrina, Myiarchus crinitus.Mviodynastes luteiventris.Nomonyx dominicus.Nycticorax nycticorax.Pandion haliaetus.Parkesia noveboracensis, Petrochelidon pyrrhonota, Podilymbus podiceps, Protonotaria citrea, Sayornis nigricans institution:

edition: Hillstar E

address: publisher: accessed: websites: chapter: series: pages: 391 issue: year: 2005 0: Kaufman, K

title: Guía de campo a las aves de Norteamérica

source: type: book

tags: Anas clypeata, Anas platyrynchos, Ardea herodias, Bartramia longicauda, Buteo swainsoni, Cairina moschata, Calidris melanotos, Columbina talpacoti, Contopus cooperi, Contopus virens, Crotophaga ani, Dendrocygna autumnalis, Dolichonyx oryzivorus, Egretta caerulea, Egretta thula, Empidonax alnorum, Empidonax traillii, Falco columbarius, Falco sparverius, Geothlypis philadelphia, Hirundo rustica, Leiothlypis peregrina, Myiarchus crinitus. Myiodynastes luteiventris, Nomonyx dominicus, Nycticorax, nycticorax, Pandion haliaetus, Parkesia noveboracensis, Petrochelidon pyrrhonota, Podilymbus podiceps, Protonotaria citrea, Sayornis nigricans abstract:

last modified:

created: group id: profile id: id:

geoPoint:

location:

thumbnail URL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created: mimeType: dataType: identifier:

fullDescriptionUnstructured: El Correlimos Pechirrayado es el playero más grande que la mayoría de las especies del género. Tiene un tamaño entre 20 y 23 cm, cuello proporcionalmente largo, pico corto y ligeramente decurvado, patas amarillo verdoso y una marcada separación del color del pecho con respecto al blanco del abdomen (Canevari et al., 2001). Por encima fuertemente estriado de negro; cabeza, cuello y pecho ante estriado café en fuerte contraste con partes inferiores blancas. En vuelo muestra una barra alar blanquecina y centro negro ancho en rabadilla y cola (Canevari et al., 2001). En época reproductiva muestra algunas plumas de la cabeza y el dorso con estriado negruzco y canela (Alderfer et al., 2006; Kaufman, 2005).

briefDescription:

references

isbn: 958 33-2118-4

keywords: Bartramia longicauda, Calidris bairdii, Calidris melanotos

editors

institution:

edition: Primera ed

address:

publisher: American Bird Conservancy, WWF-US, Humedales para las Américas y Manomet Conservation Science,

Asociación Calidris

accessed: websites: chapter: series: pages: 141 issue: year: 2001

authors

authors: Canevari, P authors: Castro, G authors: Sallaberry, M authors: Naranjo, L

title: Guía de los Chorlos y Playeros de la Región Neotropical

source: type: book

tags: Bartramia longicauda, Calidris bairdii, Calidris melanotos

abstract:

identifiers

last_modified:

created:

group_i profile_i id:	
taxonR	ecordName
ancilla	ryData
provide anyTwo provide	
publica	tionStatus
source: datatype identifie source: datatype identifie note:	r: э: r:
	sidered:
microRe source: datatype identifie source: datatype identifie note:	r: e:
sanctio	ned
microRe source: datatype identifie source: datatype identifie	r: e:

note:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered: laterHomonymOf microReference:

ruleConsidered:

conservedAgainst

basedOn

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

basionym

microReference:

source:
datatype:
identifier:
source:
datatype:
identifier:
note:

ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:

microReference:

year:

publishedIn

simple:

source: datatype: identifier:	
specialAuthorship	
authors	
year	
simple:	
canonicalAuthorship	
authors	
year	
simple: simple: (Vieillot, 1819)	
canonicalName	
infraspecificEpithet: specificEpithet: infragenericEpithet: linkType: ref:	
uninomial: simple: Calidris melanotos rank: SPECIES simple: Calidris melanotos (Vieillot, 1819))
attributes	
nomenclaturalCode: isAnamorphic: id:	
synonymsAtomized	
ancillaryData	
synonymStatus:	
synonymName	
anyTwo:	
anyOne	
providerLink: microReference:	
publishedIn	

source: datatype: identifier:

reletedName

source: datatype: identifier:

note:

ruleConsidered: microReference:

publishedIn

source: datatype: identifier:

reletedName

source: datatype: identifier: note:

ruleConsidered: microReference:

publishedIn

source: datatype: identifier:

reletedName

source: datatype: identifier: note:

ruleConsidered: microReference:

publishedIn

source: datatype: identifier:

reletedName

source: datatype: identifier: note:

ruleConsidered: microReference:

publishedIn

source: datatype: identifier:

reletedName

source: datatype: identifier: note: ruleConsidered: microReference: publishedIn source: datatype: identifier: reletedName source: datatype: identifier: note: ruleConsidered: typeNameEntity source: datatype: identifier: source: datatype: identifier: source: datatype: identifier: typeVoucherEntity typeOfType: simple: microReference: year: source: datatype: identifier: combinationAuthorship basionymAuthorship simple: authorship simple: simple: Vieillot, 1819 epithet infraspecificEpithet: specificEpithet: infragenericEpithet:

genus

linkType: ref: uninomial:

simple: Tringa melanotos

rank: SPECIES

simple: Tringa melanotos Vieillot, 1819

nomenclaturalCode: isAnamorphic:

id:

hierarchy

ancillaryData

dataObject:

additionalInformation:

reference

geoPoint: location: thumbnailURL:

mediaURL

description:

subject

source: The Catalogue of Life, 3rd January 2011

audience

bibliographicCitation:

rigthsHolder:

rights: license: title:

modified: 2016-02-16T00:26:28.867Z created: 2016-02-16T00:26:28.867Z

agent

homepage:

role: email: phone: address: position: organisation: lastName: firstName: mimeType: dataType:

identifier:

parentTaxon: Calidris higherClassification: infraspecificEpithet:

specificEpithet:

taxonRank: SPECIES

subGenus: genus: Calidris family: Scolopacidae order: Charadriiformes classHierarchy: Aves phylum: Chordata kingdom: Animalia recommended: classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:

additionalInformation:

0

isbn: 0-618-57424-7

keywords: Anas clypeata, Anas platyrynchos, Ardea herodias, Bartramia longicauda, Buteo swainsoni, Cairina moschata, Calidris melanotos, Columbina talpacoti, Contopus cooperi, Contopus virens, Crotophaga ani, Dendrocygna autumnalis, Dolichonyx oryzivorus, Egretta caerulea, Egretta thula, Empidonax alnorum, Empidonax traillii, Falco columbarius, Falco sparverius, Geothlypis philadelphia, Hirundo rustica, Leiothlypis peregrina, Myiarchus crinitus, Myiodynastes luteiventris, Nomonyx dominicus, Nycticorax nycticorax, Pandion haliaetus, Parkesia noveboracensis, Petrochelidon pyrrhonota, Podilymbus podiceps, Protonotaria citrea, Sayornis nigricans institution:

edition: Hillstar E

address: publisher: accessed: websites: chapter: series: pages: 391 issue: year: 2005

0: Kaufman, K

title: Guía de campo a las aves de Norteamérica

source: type: book

tags: Anas clypeata,Anas platyrynchos,Ardea herodias,Bartramia longicauda,Buteo swainsoni,Cairina moschata,Calidris melanotos,Columbina talpacoti,Contopus cooperi,Contopus virens,Crotophaga ani,Dendrocygna autumnalis,Dolichonyx oryzivorus,Egretta caerulea,Egretta thula,Empidonax alnorum,Empidonax traillii,Falco columbarius,Falco sparverius,Geothlypis philadelphia,Hirundo rustica,Leiothlypis peregrina,Myiarchus crinitus,Myiodynastes luteiventris,Nomonyx dominicus,Nycticorax nycticorax,Pandion haliaetus,Parkesia noveboracensis,Petrochelidon pyrrhonota,Podilymbus podiceps,Protonotaria citrea,Sayornis nigricans

abstract:

last modified:

created:

group_id:

profile_id:

id:

source:

link: http://eol.org/pages/1049474/overview keywords: Calidris melanotos 0: Frankis, Michael institution: address: publisher: accessed: websites: chapter: series: issue: 0: EOL title: Calidris melanotos source: Encyclopedia of life type: miscellany tags: Calidris melanotos abstract: last_modified: created: group_id: profile_id: id: link: http://bna.birds.cornell.edu/bna/species/348/articles/introduction keywords: Calidris melanotos 0: Poole, A institution: address: publisher: accessed: websites: chapter: series: issue: year: 2013 0: Farmer, A 1: Holmes, R 2: Pitelka, F title: Pectoral Sandpiper (Calidris melanotos) source: The Birds of North America Online type: miscellany tags: Calidris melanotos abstract: last_modified: created: group_id: profile_id: id: geoPoint: location: thumbnailURL: description:

bibliographicCitation: rightsHolder: rights: Asociación Bogotana de Ornitología license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: reproductionUnstructured: Su temporada reproductiva inicia a finales de mayo y principios de junio. Tiene un sistema de apareamiento promiscuo, los machos se aparean con varias hembras sucesivamente y estas visitan los territorios de varios machos para aparearse (EOL, n.d.; Kaufman, 2005). La hembra incuba los huevos durante cuatro días, los pichones dejan el nido y la mayoría de veces el macho abandona el área de reproducción, la hembra se mantiene con el juvenil hasta que empieza a volar, alrededor de los 21 días (EOL, n.d.; Farmer et al., 2013).
0
dataObject: additionalInformation:
reference
geoPoint: location: thumbnailURL:
mediaURL
description:
subject
audience
rightsHolder: rights: license: modified: created:
agent
mimeType: dataType: identifier: relatedTo: measurementRemarks: measurementMethod: measurementDeterminedBy: TraitBank measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: Sexual measurementType: Tipo de Reproducción measurementID:

molecularData

molecularDataUnstructured:
migratory
migratoryUnstructured:
lifeForm
lifeFormUnstructured:
lifeCycle
lifeCycleUnstructured:
interactions
interactionsUnstructured:
feeding
0
dataObject: additionalInformation:
0
isbn: 0-618-57424-7 keywords: Anas clypeata,Anas platyrynchos,Ardea herodias,Bartramia longicauda,Buteo swainsoni,Cairina moschata,Calidris melanotos,Columbina talpacoti,Contopus cooperi,Contopus virens,Crotophaga ani,Dendrocygna autumnalis,Dolichonyx oryzivorus,Egretta caerulea,Egretta thula,Empidonax alnorum,Empidonax traillii,Falco columbarius,Falco sparverius,Geothlypis philadelphia,Hirundo rustica,Leiothlypis peregrina,Myiarchus crinitus,Myiodynastes luteiventris,Nomonyx dominicus,Nycticorax nycticorax,Pandion haliaetus,Parkesia noveboracensis,Petrochelidon pyrrhonota,Podilymbus podiceps,Protonotaria citrea,Sayornis nigricans institution: edition: Hillstar E address: publisher: accessed: websites: chapter: series: pages: 391 issue: year: 2005 0: Kaufman, K title: Guía de campo a las aves de Norteamérica source: type: book tags: Anas clypeata,Anas platyrynchos,Ardea herodias,Bartramia longicauda,Buteo swainsoni,Cairina moschata,Calidris melanotos,Columbina talpacoti,Contopus cooperi,Contopus virens,Crotophaga ani,Dendrocygna
autumnalis, Dolichonyx oryzivorus, Egretta caerulea, Egretta thula, Empidonax alnorum, Empidonax traillii, Falco columbarius, Falco sparverius, Geothlypis philadelphia, Hirundo rustica, Leiothlypis peregrina, Myiarchus crinitus, Myiodynastes luteiventris, Nomonyx dominicus, Nycticorax nycticorax, Pandion haliaetus, Parkesia noveboracensis, Petrochelidon pyrrhonota, Podilymbus podiceps, Protonotaria citrea, Sayornis nigricans abstract:
last_modified:
created: group_id:
profile_id:

id: isbn: 958 33-2118-4 keywords: Bartramia longicauda, Calidris bairdii, Calidris melanotos institution: edition: Primera ed address: publisher: American Bird Conservancy, WWF-US, Humedales para las Américas y Manomet Conservation Science, Asociación Calidris accessed: websites: chapter: series: pages: 141 issue: year: 2001 0: Canevari, P 1: Castro, G 2: Sallaberry, M 3: Naranjo, L title: Guía de los Chorlos y Playeros de la Región Neotropical source: type: book tags: Bartramia longicauda, Calidris bairdii, Calidris melanotos abstract: last_modified: created: group_id: profile_id: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Asociación Bogotana de Ornitología license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: feedingUnstructured: El Correlimos Pechirrayado se alimenta principalmente de invertebrados, incluyendo en su dieta insectos, arácnidos, crustáceos y moluscos, pero también come algas y semillas que obtiene del sondeo con su pico en el lodo (Canevari et al., 2001; Kaufman, 2005). 0

type: Heterótrofos - Omnívoros

environmentalEnvelopeUnstructured:

environmentalEnvelope

ecologicalSignificance
ecologicalSignificanceUnstructured:
dispersal
dispersalUnstructured: distance: structureDispersed: type: purpose:
behavior
0
dataObject: additionalInformation:
0
keywords: Actitis macularius,Ampelion rubrocristatus,Anas discors,Anisognathus igniventris,Ardea alba,Arremon assimilis,Asio flammeus,Asio stygius,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Calidris melanotos 0: Stiles, F G 1: Bohórquez, C I 2: Cadena, C D 3: De la Zerda, S 4: Hernández, M 5: Roselli, L 6: Kelsey, M 7: Valencia, I D 8: Knapp, D institution: edition: Primera Ed address: publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca
accessed: websites:
chapter:
series: pages: 276
issue: year: 2000
0: Asociación Bogotana de Ornitología
title: Aves de la Sabana de Bogotá: guía de campo source:
type: book
tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos abstract:
last_modified:
created:
group_id: profile_id:
id:

1

isbn: 0-618-57424-7

keywords: Anas clypeata,Anas platyrynchos,Ardea herodias,Bartramia longicauda,Buteo swainsoni,Cairina moschata,Calidris melanotos,Columbina talpacoti,Contopus cooperi,Contopus virens,Crotophaga ani,Dendrocygna autumnalis,Dolichonyx oryzivorus,Egretta caerulea,Egretta thula,Empidonax alnorum,Empidonax traillii,Falco columbarius,Falco sparverius,Geothlypis philadelphia,Hirundo rustica,Leiothlypis peregrina,Myiarchus crinitus,Myiodynastes luteiventris,Nomonyx dominicus,Nycticorax nycticorax,Pandion haliaetus,Parkesia noveboracensis,Petrochelidon pyrrhonota,Podilymbus podiceps,Protonotaria citrea,Sayornis nigricans institution:

edition: Hillstar E

address: publisher: accessed: websites: chapter: series: pages: 391 issue: year: 2005 0: Kaufman, K

title: Guía de campo a las aves de Norteamérica

source: type: book

tags: Anas clypeata,Anas platyrynchos,Ardea herodias,Bartramia longicauda,Buteo swainsoni,Cairina moschata,Calidris melanotos,Columbina talpacoti,Contopus cooperi,Contopus virens,Crotophaga ani,Dendrocygna autumnalis,Dolichonyx oryzivorus,Egretta caerulea,Egretta thula,Empidonax alnorum,Empidonax traillii,Falco columbarius,Falco sparverius,Geothlypis philadelphia,Hirundo rustica,Leiothlypis peregrina,Myiarchus crinitus,Myiodynastes luteiventris,Nomonyx dominicus,Nycticorax nycticorax,Pandion haliaetus,Parkesia noveboracensis,Petrochelidon pyrrhonota,Podilymbus podiceps,Protonotaria citrea,Sayornis nigricans abstract:

last_modified: created:

group_id: profile_id:

id:

2

keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris. Ardea alba. Ardea herodias. Arremon assimilis. Asio flammeus. Asio stygius. Asthenes flammulata. Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis.Diglossa lafresnavii.Diglossa sittoides.Dolichonyx oryzivorus.Dubusia taeniata.Egretta thula.Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus

verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes Iuteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris. Ochthoeca diadema. Ochthoeca fumicolor. Ochthoeca rufipectoralis. Orochelidon murina,Oxypogon guerinii,Oxyura jamaicensis,Pachyramphus versicolor,Pandion haliaetus,Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis institution:

address:

accessed:

publisher: Princeton University Press

websites: chapter: series: pages: 836 issue: year: 1986 0: Hilty, Steven 1: Brown, William

title: A Guide to the Birds of Colombia

source: type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis.Henicorhina leucophrys.Hirundo rustica.Icterus chrysater.Icterus icterus.licterus nigrogularis.Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia

victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina,Oxypogon guerinii,Oxyura jamaicensis,Pachyramphus versicolor,Pandion haliaetus,Pardirallus maculatus.Parkesia noveboracensis.Patagioenas fasciata.Petrochelidon pyrrhonota.Phaetusa simplex.Phalacrocorax brasilianus.Pheucticus aureoventris.Pheucticus Iudovicianus.Phimosus infuscatus.Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis abstract:

last modified:

created:

group_id:

profile_id:

id:

3

link: http://bna.birds.cornell.edu/bna/species/348/articles/introduction

keywords: Calidris melanotos

0: Poole, A

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2013

0: Farmer, A

1: Holmes, R

2: Pitelka. F

title: Pectoral Sandpiper (Calidris melanotos) source: The Birds of North America Online

type: miscellany

tags: Calidris melanotos

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:
source:
bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:

behaviorUnstructured: Es una especie pasiva, por lo general se puede observar solitario o en pequeños grupos forrajeando sobre bancos de lodo con pasto y bordes de humedales; cuando se levanta vuela en zigzag como la Caica (Gallinago nobilis) (Asociación Bogotana de Ornitología, 2000).

Su reclamo un yrrk! o dsrrt! vibrante y ronco al que le agrega notas dsrrt-d´dyrrt en staccato durante el vuelo, también un meloso chriit (Kaufman, 2005). En las zonas de parada y de migración es común escuchar una serie de vocalizaciones como prrip prrip que realiza cuando aterriza o sale en vuelo, igualmente un tr´r´r´r´k o ch´r´r´t´ áspero y raspante cuando se encuentra con otros playeros (Asociación Bogotana de Ornitología, 2000; Hilty & Brown, 1986). En cortejo da unas vocalizaciones graves ju´s (Farmer et al., 2013).

ancillaryData

dataObject:

identifier:

additionalInformation:

geoPoint: location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

mimeType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

endemicAtomized

distribution

ancillaryData

dataObject: additionalInformation:

reference

isbn: 0963699636

keywords: Anas clypeata, Anas platyrynchos, Ardea herodias, Astragalinus psaltria, Bartramia longicauda, Buteo swainsoni, Cairina moschata, Calidris melanotos, Cardellina canadensis, Cathartes aura, Catharus minimus, Coccyzus americanus, Columbina talpacoti, Contopus cooperi, Contopus virens, Coragyps atratus, Crotophaga ani, Dendrocygna autumnalis, Egretta caerulea, Egretta thula, Empidonax alnorum, Empidonax traillii, Falco columbarius, Fulica americana, Geothlypis philadelphia, Hirundo rustica, Ixobrychus exilis, Jacana jacana, Mniotilta varia, Nycticorax nycticorax, Oxyura jamaicensis, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Phalacrocorax brasilianus, Piranga olivacea, Pitangus sulphuratus, Podilymbus podiceps, Porzana carolina, Pyrocephalus rubinus, Rupornis magnirostris, Setophaga fusca, Sturnella magna, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon

editors

editors: Alderfer, Jonathan

institution: address:

publisher: Washington D.C. National Geographic Society

accessed: websites: chapter: series: pages: 664 issue: year: 2006

authors

authors: Alderfer, J authors: Zickefoose, J authors: Littlehales, B

title: National Geographic Complete Birds of North America

source: type: book

tags: Anas clypeata,Anas platyrynchos,Ardea herodias,Astragalinus psaltria,Bartramia longicauda,Buteo swainsoni,Cairina moschata,Calidris melanotos,Cardellina canadensis,Cathartes aura,Catharus minimus,Coccyzus americanus,Columbina talpacoti,Contopus cooperi,Contopus virens,Coragyps atratus,Crotophaga ani,Dendrocygna autumnalis,Egretta caerulea,Egretta thula,Empidonax alnorum,Empidonax traillii,Falco columbarius,Fulica americana,Geothlypis philadelphia,Hirundo rustica,Ixobrychus exilis,Jacana jacana,Mniotilta varia,Nycticorax nycticorax,Oxyura jamaicensis,Pandion haliaetus,Pardirallus maculatus,Parkesia noveboracensis,Phalacrocorax brasilianus,Piranga olivacea,Pitangus sulphuratus,Podilymbus podiceps,Porzana carolina,Pyrocephalus rubinus,Rupornis magnirostris,Setophaga fusca,Sturnella magna,Tringa flavipes,Tringa melanoleuca,Tringa solitaria,Troglodytes aedon

abstract:

identifiers

last_modified: created:

group_id: profile id:

id:

geoPoint: location:

thumbnailURL:
mediaURL
description:
subject
source:
audience
bibliographicCitation: rightsHolder: rights: Asociación Bogotana de Ornitología license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created:
agent
mimeType: dataType: identifier: distributionUnstructured: Es una especie migratoria de larga distancia; se reproduce en el círculo polar ártico en Siberia y Norteamérica (Alaska y Canadá), desde la península Yamal, en el oeste de Rusia hasta la bahía de Hudson. Transeúnte en Centroamérica, Antillas Mayores y Menores, Trinidad y Tobago, Colombia, Venezuela y Ecuador (Alderfer et al., 2006). Frecuenta pantanos, planicies con pasto, campos inundados, charcos desde el nivel del mar hasta los 3100 msnm (Asociación Bogotana de Ornitología, 2000; Ruíz-Guerra, 2012). Residente de invierno en Perú, sur de Bolivia, sur de Brasil y norte de Argentina y Uruguay hasta el centro de Chile (Narosky & Yzurieta, 2010).
distributionAtomized
stateProvince: Córdoba
ancillaryData
locality: municipality: country: Colombia
temporalCoverage
endDate: startDate:
distributionScope
type:
threatStatus
ancillaryData

apendiceCITES: locality: municipality:

threatStatusUnstructured: threatStatusAtomized

county: stateProvince: country: Global 0: IUCN measurementValue: Preocupación menor (LC) relatedTo: measurementRemarks: measurementMethod: measurementDeterminedBy measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementType: IUCN measurementID: territory territoryUnstructured: areaOfOccupancy: extentOfOccurrence: populationBiology 0 dataObject: additionalInformation: link: http://maps.iucnredlist.org/map.html?id=22693408 keywords: Calidris melanotos institution: address: publisher: accessed: websites: chapter: series: issue: year: 2014 0: BirdLife International 1: NatureServe title: Bird Species Distribution Maps of the World. 2012. Calidris melanotos. The IUCN Red List of Threatened Species. Version 2014.3 source: type: miscellany tags: Calidris melanotos abstract: last_modified: created: group_id: profile_id: geoPoint: location:

thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Asociación Bogotana de Ornitología license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: populationBiologyUnstructured: Es una especie común en las zonas de anidación, tiene un amplio rango de distribución y la tendencia de la población parece estar estable (BirdLife International and NatureServe, 2014).
legislation
legislationUnstructured:
directThreats
directThreatsUnstructured:
ancillaryData
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified: created: mimeType: dataType: identifier:
measurementOrFact
relatedTo: measurementRemarks: measurementMethod: measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: measurementType: measurementID: abstract: El Correlimos Pechirrayado es una especie de playero migratorio boreal. El nombre griego Kalidris fue aplicado por Aristóteles a un ave de orilla, el nombre de la especie proviene del griego melas = negro y notos = espaldar, para describir detalles del plumaje. Se reproduce en el círculo polar ártico en Siberia y Norteamérica (Alaska y Canadá), desde la península Yamal, en el oeste de Rusia hasta la bahía de Hudson. Transeúnte en Centroamérica, Antillas Mayores y Menores, Trinidad y Tobago, Colombia, Venezuela y Ecuador. Residente de invierno en Perú, sur de Bolivia, sur de Brasil y norte de Argentina y Uruguay hasta el centro de Chile (Canevari et

al., 2001). Más grande que la mayoría de las especies del género, mide entre 20 y 24 cm, el plumaje por encima café oscuro listado con café claro, con dos listas anchas anteadas a lo largo de cada lado del dorso (Canevari et al., 2001). Generalmente habita en bordes de cuerpos de agua, potreros y charcos donde busca su alimento (Asociación Bogotana de Ornitología, 2000; Hilty & Brown, 1986).

associatedParty

role:

personnelldentifier: personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone: postalCode: country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address: organisation: position:

lastName: Gómez Melendro firstName: Edith Natalia

ancillaryData

dataObject:

additionalInformation:

reference

geoPoint: location: thumbnailURL:

mediaURL: http://neotropical.birds.cornell.edu/portal/species/overview?p%0dp_spp=153141

description:

subject

source: http://neotropical.birds.cornell.edu

audience

bibliographicCitation:

rightsHolder: rights:

license: modified: created:

agent

mimeType: dataType: identifier:

_id

_bsontype: ObjectID id: VÂoXð Ig#ç2À