Reporte de Fichas para el usuario: edith gómez
Ficha ID: 56e2cefb83c45700544e3d79 Fecha de creación: 2016-03-11T13:58:26.367Z
annualCycles
annualCycleUnstructured:
habitats
habitatUnstructured:
commonNamesAtomized
ancillaryData
dataObject: additionalInformation:
reference
geoPoint: location: thumbnailURL:
mediaURL
description:
subject
source:
audience
bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created:
agent
mimeType: dataType: identifier: usedBy:
usedIn
distributionUnstructured: endDate: startDate:
ancillaryData
type: language: Español name: Avispa creation_date: 2016-03-11T13:58:26.367Z

uses Management And Conservation

## ancillaryData

managementAndConservationUnstructured:

## managementAndConservationAtomized

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

geoPoint: location:

thumbnailURL: description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created: mimeType: dataType: identifier:

fullDescriptionUnstructured: Hembra oscura con dientes mandibulares y margen ventral del clípeo rojos. Parte inferior del flagelo antenal y parte superior del segmento 3 rojo oscuro. Parte superior del clípeo rojiza con difusas manchas oscuras. Tégula café, alas hialinas con región costal amarillo oscuro y venación café, longitud 10 mm. Clípeo moderadamente angular, ápice débilmente truncado, superficie toscamente granulada. Frente con punteaduras, ocelos posteriores separados por un poco menos que dos diámetros, gena más ancha que la mitad del ancho del seno ocular, espacio malar ligermante más largo que en M. wagneri. Quilla pronotal recta, un poco curva en la mitad, aguda a los lados. Propodeo con un surco largo pero sin quilla. Tercer segmento del taro medio moderadamente asimétrico. Macho muy parecido a la hembra. Clípeo débilmente angular, superficie con pelos plateados. Espacio malar ligeramente corto. Lado interno de la tibia posterior sin mancha amarilla. briefDescription:

### references

keywords: Agelaia angulata angulata,Agelaia angulicollis,Agelaia brevistigma,Agelaia cajennensis,Agelaia fulvofasciata,Agelaia hamiltoni,Agelaia testacea,Angiopolybia obidensis,Angiopolybia pallens,Angiopolybia paraensis,Angiopolybia zischkai,Apoica thoracica,Asteloeca traili,Brachygastra albula,Brachygastra augusti,Brachygastra lecheguana,Brachygastra propodealis,Brachygastra scutellaris,Chartergellus amazonicus,Chartergellus punctatior,Chartergus chartarius,Chartergus globiventris,Chartergus metanotalis,Clypearia sulcata,Clypearia weyrauchi,Epipona tatua,Leipomeles dorsata,Metapolybia acincta,Metapolybia docilis,Metapolybia nigra,Metapolybia suffusa,Mischocyttarus adolphi,Mischocyttarus alfkenii excrucians,Mischocyttarus alfkenii trinitatis,Mischocyttarus carbonarius carbonarius,Mischocyttarus flavicans,Mischocyttarus foveatus,Mischocyttarus injucundus tingomariae,Mischocyttarus insolitus,Mischocyttarus labiatus,Mischocyttarus latissimus,Mischocyttarus metoecus,Mischocyttarus petiolatus,Mischocyttarus pseudomimeticus,Mischocyttarus punctatus,Mischocyttarus tomentosus,Mischocyttarus vaqueroi,Parachartergus amazonensis,Parachartergus colobopterus,Parachartergus fasciipennis,Parachartergus fraternus,Parachartergus fulgidipennis,Parachartergus griseus,Parachartergus richardsi,Parachartergus weyrauchi,Polybia bistriata,Polybia catillifex,Polybia eberhardae,Polybia emaciata,Polybia fastidiosuscula,Polybia jurinei,Polybia liliacea,Polybia micans,Polybia occidentalis,Polybia platycephala

sylvestris, Polybia quadricincta, Polybia raui raui, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis, Polybia sericea, Polybia striata, Polybia velutina, Protopolybia acutiscutis, Protopolybia amarella, Protopolybia chartergoides chartergoides, Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea

### editors

institution: address: publisher: accessed: websites: chapter: series: pages: 585 issue: year: 1978

#### authors

authors: Richards, O. W.

title: The social wasps of the Americas excluding the Vespinae

source: type: book

tags: Agelaia angulata angulata, Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis, Angiopolybia zischkai, Apoica thoracica, Asteloeca traili, Brachygastra albula, Brachygastra augusti, Brachygastra lecheguana, Brachygastra propodealis, Brachygastra scutellaris, Chartergellus amazonicus, Chartergellus punctatior, Chartergus chartarius, Chartergus globiventris, Chartergus metanotalis, Clypearia sulcata, Clypearia weyrauchi, Epipona tatua, Leipomeles dorsata, Metapolybia acincta. Metapolybia docilis. Metapolybia nigra. Metapolybia suffusa. Mischocyttarus adolphi. Mischocyttarus alfkeni, Mischocyttarus alfkenii excrucians, Mischocyttarus alfkenii trinitatis, Mischocyttarus carbonarius carbonarius, Mischocyttarus flavicans, Mischocyttarus foveatus, Mischocyttarus injucundus tingomariae, Mischocyttarus insolitus, Mischocyttarus labiatus, Mischocyttarus latissimus, Mischocyttarus metoecus, Mischocyttarus petiolatus, Mischocyttarus pseudomimeticus, Mischocyttarus punctatus, Mischocyttarus tomentosus, Mischocyttarus vaqueroi, Parachartergus amazonensis, Parachartergus colobopterus, Parachartergus fasciipennis, Parachartergus fraternus, Parachartergus fulgidipennis, Parachartergus griseus, Parachartergus richardsi, Parachartergus weyrauchi, Polybia bistriata, Polybia catillifex, Polybia eberhardae, Polybia emaciata, Polybia fastidiosuscula, Polybia jurinei, Polybia liliacea, Polybia micans, Polybia occidentalis, Polybia platycephala sylvestris, Polybia quadricincta, Polybia raui raui, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis.Polybia sericea.Polybia striata.Polybia velutina.Protopolybia acutiscutis.Protopolybia amarella.Protopolybia chartergoides chartergoides, Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea abstract:

#### identifiers

last\_modified: created: group\_id: profile\_id:

id:

taxonRecordName

# ancillaryData

# providerSpecificData anyTwo: providerLink: publicationStatus microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: replacementNameFor microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: sanctioned microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: laterHomonymOf microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: conservedAgainst

basedOn

## authors

## year

simple:

canonicalAuthorship

### authors

## year

simple:

simple: Richards, 1978

canonicalName

infraspecificEpithet: specificEpithet: infragenericEpithet:

linkType: ref:

uninomial:

simple: Mischocyttarus petiolatus

rank: SPECIES

simple: Mischocyttarus petiolatus Richards, 1978

### attributes

nomenclaturalCode: isAnamorphic:

id.

# synonymsAtomized

# hierarchy

# ancillaryData

dataObject:

additionalInformation:

### reference

geoPoint: location: thumbnailURL:

## mediaURL

description:

## subject

source: The Catalogue of Life, 3rd January 2011

## audience

bibliographicCitation:

rigthsHolder: rights: license: title: modified: 2016-03-11T13:58:06.365Z created: 2016-03-11T13:58:06.365Z agent homepage: role: email: phone: address: position: organisation: lastName: firstName: mimeType: dataType: identifier: parentTaxon: Mischocyttarus higherClassification: infraspecificEpithet: specificEpithet: taxonRank: SPECIES subGenus: genus: Mischocyttarus family: Vespidae order: Hymenoptera classHierarchy: Insecta phylum: Arthropoda kingdom: Animalia recommended: classification: invasiveness invasivenessUnstructured: reproduction dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created: mimeType:

dataType: identifier: reproductionUnstructured:
0
dataObject: additionalInformation:
reference
geoPoint: location: thumbnailURL:
mediaURL
description:
subject
audience
rightsHolder: rights: license: modified: created:
agent
mimeType: dataType: identifier: relatedTo: measurementRemarks: measurementMethod: measurementDeterminedBy: TraitBank measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: Sexual measurementType: Tipo de Reproducción measurementID:
molecularData
molecularDataUnstructured:
migratory
migratoryUnstructured:
lifeForm
lifeFormUnstructured:
lifeCycle
lifeCycleUnstructured:
interactions

interactionsUnstructured:
feeding
0
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: feedingUnstructured: Las avispas de este género caminan a lo largo de los hilos de las telarañas, presumiblemente gracias a una especial adaptación en los lóbulos tarsales de las patas medias y posteriores. Después de esto las avispas forrajeras retornan al nido y allí mastican la comida compartiéndola con otras avispas de la colonia, las larvas son alimentadas por regurgitación. El néctar recolectado de una amplia variedad de flores también es llevado al nido y guardado en sus celdas para luego ser consumido por los adultos.  0
0
strategy: Depredadores type: Heterótrofos - Omnívoros
environmentalEnvelope
environmentalEnvelopeUnstructured:
ecologicalSignificance
ecologicalSignificanceUnstructured:
dispersal
dispersalUnstructured: distance: structureDispersed: type: purpose:
behavior
0
dataObject: additionalInformation: geoPoint: location: thumbnailURL:

description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: behaviorUnstructured: Es una especie de avispa depredadora con hábitos sociales.
ancillaryData
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified: created: mimeType: dataType: identifier:
measurementOrFact
relatedTo: measurementRemarks: measurementMethod: measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: measurementType: measurementID:
endemicAtomized
distribution
ancillaryData
dataObject: additionalInformation:
reference
geoPoint: location: thumbnailURL:

mediaURL
description:
subject
source:
audience
bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created:
agent
mimeType: dataType: identifier: distributionUnstructured: Se distribuye en áreas de hasta 1100 msnm. En el país se ha registrado en la región de la Amazonia.
distributionAtomized
ancillaryData
locality: municipality: country: Perú
temporalCoverage
endDate: startDate:
distributionScope
type:
threatStatus
territory
territoryUnstructured: areaOfOccupancy: extentOfOccurrence:
populationBiology
populationBiologyUnstructured:
legislation
legislationUnstructured:
directThreats
directThreatsUnstructured:
ancillaryData

dataObject	Ċ.
additionalli	Π

nformation:

geoPoint: location: thumbnailURL: description: source:

bibliographicCitation:

rightsHolder:

rights: license: modified: created: mimeType: dataType: identifier:

### measurementOrFact

relatedTo:

measurementRemarks: measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: Es una especie de avispa depredadora que se distribuye en áreas de hasta 1100 msnm de altura en Colombia y Perú. La hembra presenta una coloración oscura y alas hialinas con la región costal amarilla oscura y la venación café. Es una especie con hábitos sociales que vive y se desarrolla en colonias.

### associatedParty

role:

personnelldentifier: personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone: postalCode: country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address: organisation: position:

lastName: Gómez Melendro firstName: Edith Natalia

## ancillaryData

### dataObject:

additionalInformation: Información de tipos: El holotipo de esta especie está depositado en el British Museum of Natural History (BMNH), Londres, Inglaterra.

Créditos específicos: Al Instituto Amazónico de Investigaciones Científicas Sinchi. Al Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIAT-AC. La elaboración de esta ficha fue posible gracias al apoyo del gobierno de Finlandia a través del convenio entre la Secretaria General de la CAN y el Instituto Sinchi para el

proyecto "Sistema Regional Amazónico de Información Ambiental con énfasis en Biodiversidad de los países miembros de la CAN, (programa BIOCAN), componente SIAT-AC".

reference

geoPoint:
location:
thumbnailURL:

## mediaURL

description:

# subject

source:

## audience

bibliographicCitation: rightsHolder:

rights:

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

## agent

mimeType: dataType: identifier:

\_id

\_bsontype: ObjectID id: VâÎûfÄW Dãx•