

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56d4ca783c16479905cba8c6

Fecha de creación: 2016-02-29T22:47:22.933Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Clara Maria Pérez Gallego

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataObject:

identifier:

habitatUnstructured: Esta especie se encuentra desde selva hasta áreas intervenidas. Esta especie habita claro con matorrales, áreas cultivadas y bordes de selva.

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

[ancillaryData](#)

type:

language: Español

name: Asoma Terciopelo

creation_date: 2016-02-29T22:47:22.933Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Clara Maria Pérez Gallego
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
fullDescriptionUnstructured: Esta especie tienen un largo total promedio 18 cm. Macho con mandíbula inferior blanco plateado reluciente, cabeza, manto, garganta y pecho rojo marrón intenso, gradado a carmesí brillante en baja espalda, rabadilla y bajas partes inferiores; alas y cola negros;, tibias y centro de abdomen negros. La hembra como el macho pero más opaca, casi negruzca en garganta y pecho, pero retiene rabadilla y partes inferiores rojas; pico negruzco.
briefDescription:

references

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:
providerLink:

publicationStatus

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

replacementNameFor

microReference:
source:
datatype:
identifier:
source:

datatype:
identifier:
note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

conservedAgainst

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion
typeOfType:

[lectotypeMicroReferenceVoucher](#)

[lectotypePublicationVoucher](#)

[voucherReference](#)

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

[authors](#)

[year](#)

simple:
canonicalAuthorship

[authors](#)

[year](#)

simple:
simple: Lafresnaye, 1837

canonicalName

infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:

ref:
uninomial:
simple: Ramphocelus dimidiatus
rank: SPECIES

simple: Ramphocelus dimidiatus Lafresnaye, 1837

attributes

nomenclaturalCode:

isAnamorphic:

id:

synonymsAtomized

hierarchy

ancillaryData

dataObject:

additionalInformation:

reference

geoPoint:

location:

thumbnailURL:

mediaURL

description:

subject

source: The Catalogue of Life, 3rd January 2011

audience

bibliographicCitation:

rightsHolder:

rights:

license:

title:

modified: 2016-02-29T22:47:02.932Z

created: 2016-02-29T22:47:02.932Z

agent

homepage:

role:

email:

phone:

address:

position:

organisation:

lastName:

firstName:

mimeType:

dataType:

identifier:

parentTaxon: Ramphocelus

higherClassification:

infraspecificEpithet:

specificEpithet:

taxonRank: SPECIES
subGenus:
genus: Ramphocelus
family: Thraupidae
order: Passeriformes
classHierarchy: Aves
phylum: Chordata
kingdom: Animalia
recommended:
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Clara Maria Pérez Gallego
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:

reproductionUnstructured: Hay registros de nidos de esta especie entre enero y marzo con dos huevos azules con puntos oscuros.

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Clara Maria Pérez Gallego
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
feedingUnstructured: Esta especie se alimenta de frutas e insectos.

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Clara Maria Pérez Gallego
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
behaviorUnstructured: Esta especie es diurna y en grupos en el sotobosque. Esta especie se encuentra usualmente en grupos en sotobosque, a lo largo de jardines y claros. Esta especie canta fuerte chink y tzuiip.

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

[endemicAtomized](#)

[distribution](#)

[ancillaryData](#)

dataObject:
additionalInformation:

[reference](#)

geoPoint:
location:
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:
rightsHolder:
rights: Clara Maria Pérez Gallego
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:

created:

[agent](#)

mimeType:

dataType:

identifier:

distributionUnstructured: Esta especie se encuentra desde W Panamá hasta W Venezuela. En el país se distribuye en todo el valle del Magdalena, Cordillera Occidental y región Caribe hasta Sierra Nevada Santa Marta hasta los 1500 msnm.

[distributionAtomized](#)

temporalCoverage

endDate:

startDate:

distributionScope

type:

[threatStatus](#)

[ancillaryData](#)

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:

locality:

municipality:

county:

stateProvince:

country: Global

0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:

measurementRemarks:

measurementMethod:

[measurementDeterminedBy](#)

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementType: IUCN

measurementID:

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

mimeType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract:

[associatedParty](#)

role:

personnelIdentifier:

personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone:

postalCode:

country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address:

organisation:

position:

lastName: Gómez Melendro

firstName: Edith Natalia

[ancillaryData](#)

dataObject:

additionalInformation: Registros biológicos: En la Central Hidroeléctrica Miel I se registró esta especie en los bosques de El Tigre, La Clara, Campiña y Cariaño, en el rastrojo del Encanto y en las áreas intervenidas de Cantera y Terraplen.
Imagen: Ramphocelus_dimidiatus.jpg

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder: Clara Maria Pérez Gallego
rights:
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

contentType:
dataType:
identifier:

_id
_bsontype: ObjectId
id: VÔËx< G™\ºœ`