

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56d5f4433c16479905cba905  
Fecha de creación: 2016-03-01T19:57:57.679Z

annualCycles  
annualCycleUnstructured:  
habitats  
habitatUnstructured:  
[commonNamesAtomized](#)

[ancillaryData](#)  
usedBy:  
usedIn  
distributionUnstructured:  
endDate:  
startDate:

[ancillaryData](#)  
type:  
language: Alemán  
name: Seidenkuhstärling  
creation\_date: 2016-03-01T19:57:57.679Z  
usesManagementAndConservation

[ancillaryData](#)  
managementAndConservationUnstructured:  
[managementAndConservationAtomized](#)

identificationKeys  
fullDescription

0  
dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Equipo Coordinador del SiB Colombia  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:

fullDescriptionUnstructured: Se reconocen siete subespecies:  
Molothrus bonariensis aequatorialis Chapman, 1915  
Molothrus bonariensis bonariensis (Gmelin, 1789)  
Molothrus bonariensis cabanisii Cassin, 1866  
Molothrus bonariensis minimus Dalmas, 1900  
Molothrus bonariensis occidentalis Berlepsch & Stolzmann, 1892  
Molothrus bonariensis riparius Griscom & Greenway, 1937  
Molothrus bonariensis venezuelensis Stone, 1891  
briefDescription:

references

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:  
providerLink:

publicationStatus

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

replacementNameFor

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

sanctioned

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

laterHomonymOf

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

[conservedAgainst](#)

basedOn

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

basionym

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

[spellingCorrentionOf](#)

typeNameEntity

source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:

typificacion

typeOfType:

[lectotypeMicroReferenceVoucher](#)

[lectotypePublicationVoucher](#)

[voucherReference](#)

simple:  
microReference:  
year:  
  
publishedIn  
  
simple:  
source:  
datatype:  
identifier:  
  
specialAuthorship

[authors](#)

[year](#)

simple:  
  
canonicalAuthorship

[authors](#)

[year](#)

simple:  
simple: (Gmelin, 1789)  
  
canonicalName  
  
infraspecificEpithet:  
specificEpithet:  
infragenericEpithet:  
linkType:  
ref:  
uninomial:  
simple: *Molothrus bonariensis*  
rank: SPECIES  
simple: *Molothrus bonariensis* (Gmelin, 1789)  
  
attributes  
  
nomenclaturalCode:  
isAnamorphic:  
id:

[synonymsAtomized](#)

[hierarchy](#)

[ancillaryData](#)

dataObject:  
additionalInformation:

[reference](#)

geoPoint:  
location:  
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:  
rightsHolder:  
rights:  
license:  
title:  
modified: 2016-03-01T19:57:37.678Z  
created: 2016-03-01T19:57:37.678Z

[agent](#)

homepage:  
role:  
email:  
phone:  
address:  
position:  
organisation:  
lastName:  
firstName:  
mimeType:  
dataType:  
identifier:  
parentTaxon: Molothrus  
higherClassification:  
infraspecificEpithet:  
specificEpithet:  
taxonRank: SPECIES  
subGenus:  
genus: Molothrus  
family: Icteridae  
order: Passeriformes  
classHierarchy: Aves  
phylum: Chordata  
kingdom: Animalia  
recommended:  
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Equipo Coordinador del SiB Colombia  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
reproductionUnstructured: Parásito de cría y promiscuo. Pone en nidos de especies más pequeñas, posiblemente hasta 5 huevos por año, estos de color blanquecino con marcas variables (Hilty & Brown 1986).

#### molecularData

molecularDataUnstructured:

#### migratory

migratoryUnstructured:

#### lifeForm

lifeFormUnstructured:

#### lifeCycle

lifeCycleUnstructured:

#### interactions

interactionsUnstructured:

#### feeding

0

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Equipo Coordinador del SiB Colombia  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
feedingUnstructured: En La Guajira habita en los bosques deciduos tropical, de galería y secundario, en matorrales

o rastros y en pastizales (Cerrejón 2009-2010).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataFormat:

identifier:

behaviorUnstructured: Camina con la cola ligeramente erecta (Hilty & Brown 1986). El canto es un gorjeo musical líquido, a veces interrumpido por notas más ásperas y usualmente acompañado por despliegue de las plumas del cuello. Permanece solitario o más frecuentemente en pequeños grupos laxos; los grandes grupos son raros con excepción de los dormideros. (Hilty & Brown 1986).

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:  
identifier:

measurementOrFact

relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

[endemicAtomized](#)

[distribution](#)

[ancillaryData](#)

dataObject:  
additionalInformation:

[reference](#)

geoPoint:  
location:  
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:  
rightsHolder:  
rights: Equipo Coordinador del SiB Colombia  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

[agent](#)

contentType:  
dataType:  
identifier:  
distributionUnstructured: En bases de datos se encontró que:  
Esta especie se distribuye en: Argentina, Ecuador (Arctos 2012)  
Las siguientes subespecies se distribuyen en:  
Molothrus bonariensis aequatorialis  
Molothrus bonariensis bonariensis Argentina, Brasil, Chile (Arctos 2012)  
Molothrus bonariensis cabanisii Colombia (Arctos 2012; Collections MNH SI 2012)  
Molothrus bonariensis minimus Trinidad y Tobago (Collections MNH SI 2012)



Molothrus bonariensis occidentalis Perú (Collections MNH SI 2012).  
En bases de datos y monitoreos de fauna se encontró que:  
Esta especie se distribuye en: La Guajira (Cerrejón 2009-2010)  
Las siguientes subespecies se distribuyen en Colombia:  
Molothrus bonariensis aequatorialis Cauca (ICN 2004); Caquetá (Collections MNH SI 2012)  
Molothrus bonariensis cabanisii Antioquia, Atlántico, Bogotá, Bolívar, Boyacá, Caldas, Cauca, Córdoba, Cundinamarca, Quindío, Santander, Tolima, Valle del Cauca (ICN 2004); Antioquia, Atlántico, Bolívar, Boyacá, Caldas, Cauca, Cesar, Chocó, Córdoba, Huila, Magdalena, Norte de Santander, Valle, Valle del Cauca (Collections MNH SI 2012); Huila, Santander, Valle del Cauca (Arctos 2012)  
Molothrus bonariensis minimus Amazonas (ICN 2004)  
Molothrus bonariensis venezuelensis Amazonas, Boyacá, Caquetá, Cundinamarca, Meta, Norte de Santander, Santander (ICN 2004); Norte de Santander (Collections MNH SI 2012).

distributionAtomized

stateProvince: Córdoba

ancillaryData

locality:  
municipality:  
country: Colombia

temporalCoverage

endDate:  
startDate:

distributionScope

type:

threatStatus

ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:  
locality:  
municipality:  
county:  
stateProvince:  
country: Global  
0: IUCN  
measurementValue: Preocupación menor (LC)  
relatedTo:  
measurementRemarks:  
measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementType: IUCN  
measurementID:

## territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

## populationBiology

populationBiologyUnstructured:

## legislation

legislationUnstructured:

## directThreats

directThreatsUnstructured:

## ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

## measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: Mide unos 22 cm. Tiene el pico corto y cónico y ojos oscuros. Los machos tienen todo el plumaje de color negro púrpura lustroso. Las hembras son de color marrón grisáceo opaco en las partes dorsales, mucho más pálido en las partes ventrales; tienen una banda superciliar corta y tenue blanco grisáceo (Hilty & Brown 1986).

## associatedParty

role:

personnelIdentifier:

personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone:

postalCode:

country: Colombia  
state: Bogotá Distrito Capital  
city: Bogotá D.C.  
address:  
organisation:  
position:  
lastName: Gómez Melendro  
firstName: Edith Natalia

## ancillaryData

dataObject:  
additionalInformation: Registros biológicos: En el portal de Global Biodiversity Information Facility se reporta que para Colombia hay 753 registros. GBIF  
Créditos específicos: Ficha elaborada con el apoyo de Carbones del Cerrejón Limited.  
Metadatos: GBIF  
Otros recursos en Internet: Arctos, Avibase, AVoCet, BirdLife International, Colecciones en Línea, EoL, iSpecies, IUCN Red List, SI NMNH, xeno-canto

## reference

geoPoint:  
location:  
thumbnailURL:

## mediaURL

description:

## subject

source:

## audience

bibliographicCitation:  
rightsHolder:  
rights: Equipo Coordinador del SiB Colombia  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

## agent

contentType:  
dataType:  
identifier:

\_id

\_bsontype: ObjectId  
id: VÖöC< G™\º•