

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56d6f4dd3c16479905cba916

Fecha de creación: 2016-03-02T14:12:47.548Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataObject:

identifier:

habitatUnstructured: Residente de invierno localmente común pero errática en muchas regiones; terreno abierto y áreas agrícolas, especialmente en rosales (Hilty & Brown 1986). En La Guajira habita en matorral o rastrojo y pastizales (Cerrejón 2009-2010).

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

[ancillaryData](#)

type:

language: Español

name: Arrocero americano

creation_date: 2016-03-02T14:12:47.548Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

fullDescriptionUnstructured:
briefDescription:

references

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:
providerLink:

publicationStatus

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

replacementNameFor

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:

datatype:
identifier:
note:
ruleConsidered:

conservedAgainst

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:
canonicalAuthorship

authors

year

simple:
simple: (Gmelin, 1789)
canonicalName
infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:
ref:
uninomial:
simple: Spiza americana
rank: SPECIES
simple: Spiza americana (Gmelin, 1789)

attributes

nomenclaturalCode:
isAnamorphic:
id:

synonymsAtomized

hierarchy

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:

rightsHolder:

rights:

license:

title:

modified: 2016-03-02T14:12:27.548Z

created: 2016-03-02T14:12:27.548Z

[agent](#)

homepage:

role:

email:

phone:

address:

position:

organisation:

lastName:

firstName:

contentType:

dataType:

identifier:

parentTaxon: Spiza

higherClassification:

infraspecificEpithet:

specificEpithet:

taxonRank: SPECIES

subGenus:

genus: Spiza

family: Cardinalidae

order: Passeriformes

classHierarchy: Aves

phylum: Chordata

kingdom: Animalia

recommended:

classification:

invasiveness

invasivenessUnstructured:

reproduction

reproductionUnstructured:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

feedingUnstructured: En La Guajira se observó que los individuos forrajean entre el suelo y el dosel y su dieta es granívora e insectívora (Cerrejón 2009-2010).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:
bibliographicCitation:
rightsHolder:
rights: Equipo Coordinador del SiB Colombia
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
behaviorUnstructured: En pequeños grupos o en densas bandadas de varios centenares o miles en áreas de alimentación favorable (Hilty & Brown 1986).

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

endemicAtomized

distribution

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder:
rights: Equipo Coordinador del SiB Colombia
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

mimeType:
dataType:
identifier:
distributionUnstructured: En bases de datos se encontró que: Esta especie se distribuye en: Estados Unidos, Honduras, México (Arctos 2012).
En bases de datos, literatura y monitoreos de fauna se encontró que: Esta especie se distribuye en: La Guajira (Cerrejón 2009-2010); Antioquia, Boyacá, Casanare, Meta, Tolima (ICN 2004); Antioquia, Bolívar, Cesar, Chocó, Córdoba, La Guajira (Collections MNH SI 2012). En las regiones Andina, Caribe, Orinoquía y Pacífica hasta los 1600 m (Hilty & Brown 1986).

distributionAtomized

stateProvince: La Guajira

ancillaryData

locality:
municipality:
country: Colombia

temporalCoverage

endDate:
startDate:

distributionScope

type:

threatStatus

ancillaryData

threatStatusUnstructured:
threatStatusAtomized

apendiceCITES:
locality:
municipality:
county:
stateProvince:

country: Global
0: IUCN
measurementValue: Preocupación menor (LC)
relatedTo:
measurementRemarks:
measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementType: IUCN
measurementID:

territory

territoryUnstructured:
areaOfOccupancy:
extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:

measurementType:

measurementID:

abstract: Mide 15 cm. Los machos tienen un plumaje distintivo durante la época de reproducción: cabeza grisáceo con lista ocular amarilla; el resto de las partes dorsales pardo claro estriado de parduzco; hombros rufos; garganta blanca con un estrecha línea malar negra conectada a una V pectoral; pecho amarillo desvanecido a blanquecino en abdomen. El plumaje no reproductivo es más opaco, con menos amarillo, V pectoral reducida o ausente. Las hembras como los machos en temporada no reproductiva tienen lista ocular más débil; blanco sucio debajo tinte amarillo en el pecho; malar parduzco que se conecta con estriado difuso en los lados (Hilty & Brown 1986).

associatedParty

role:

personnelIdentifier:

personnelDirectory:

homepage:

email: enataliagomez@gmail.com

phone:

postalCode:

country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address:

organisation:

position:

lastName: Gómez Melendro

firstName: Edith Natalia

ancillaryData

dataObject:

additionalInformation: Registros biológicos: En el portal de Global Biodiversity Information Facility se reporta que para Colombia hay 93 registros. GBIF

Créditos específicos: Ficha elaborada con el apoyo de Carbones del Cerrejón Limited.

Otros recursos en internet: Arctos, Avibase, AVoCet, BirdLife International, Colecciones en Línea, EoL, iSpecies, IUCN Red List, SI NMNH, xeno-canto

reference

geoPoint:

location:

thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

agent

contentType:
dataType:
identifier:

_id

_bsonType: ObjectId

id: VÖôÝ< GTM\0⁶