Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56ca344ff0106c67230e7458

Fecha de creación: undefined

annualCycles

annualCycleUnstructured:

habitats

0

dataObject: additionalInformation:

0

keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis.Coccycua pumila.Coccyzus americanus.Coccyzus melacoryphus.Coeligena bonapartei.Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta caerulea, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria guitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon querinii, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota.Phaetusa simplex.Phalacrocorax brasilianus.Pheucticus aureoventris.Pheucticus ludovicianus.Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus,Ramphomicron microrhynchum,Riparia riparia,Rupornis magnirostris,Sayornis nigricans,Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis institution: address:

publisher: Princeton University Press

accessed: websites: chapter: series: pages: 836 issue: year: 1986

year: 1986 0: Hilty, Steven 1: Brown, William

title: A Guide to the Birds of Colombia

source: type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis.Diglossa lafresnavii.Diglossa sittoides.Dolichonyx oryzivorus.Dubusia taeniata.Egretta caerulea.Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima.Mimus gilvus.Mniotilta varia.Molothrus bonariensis.Muscisaxicola alpinus.Mviarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon querinii, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus,Ramphomicron microrhynchum,Riparia riparia,Rupornis magnirostris,Sayornis nigricans,Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

abstract: last_modified: created:

group_id: profile_id: id: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Asociación Bogotana de Ornitología license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: habitatUnstructured: La especie es común en áreas húmedas cubiertas de hierba, especialmente con arbustos dispersos y Espeletia o en los bordes de bosque de Polylepis; también en bosques achaparrados en el ecotono con el páramo (Hilty & Brown, 1986).
commonNamesAtomized
ancillaryData
usedBy:
usedIn
distributionUnstructured: endDate: startDate:
ancillaryData
type: language: Inglés name: Andean Tit-Spinetail
usesManagementAndConservation
ancillaryData
managementAndConservationUnstructured:
managementAndConservationAtomized
identificationKeys
fullDescription
0
dataObject:
additionalInformation:
0

keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis.Catamenia analis.Catamenia inornata,Cathartes aura,Catharus minimus,Catharus ustulatus.Chaetocercus mulsant.Chaetura pelagica.Chalcostigma heteropogon.Chordeiles minor.Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta caerulea, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor.Ochthoeca rufipectoralis.Orochelidon murina.Oxypogon querinii.Oxyura jamaicensis.Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus.Ramphomicron microrhynchum,Riparia riparia,Rupornis magnirostris,Sayornis nigricans,Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola. Sicalis luteola. Spiza americana. Sporagra spinescens. Sturnella magna. Synallaxis subpudica. Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis institution:

address:

accessed:

publisher: Princeton University Press

websites: chapter: series: pages: 836 issue: year: 1986 0: Hilty, Steven 1: Brown, William

title: A Guide to the Birds of Colombia

source: type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus

holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus. Cinclodes albidiventris. Cinclus leucocephalus. Cinnycerthia unirufa. Cistothorus apolinari. Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta caerulea, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon querinii, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus,Ramphomicron microrhynchum,Riparia riparia,Rupornis magnirostris,Sayornis nigricans,Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris. Tangara vitriolina. Thraupis episcopus. Thraupis palmarum. Tringa flavipes. Tringa melanoleuca. Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis abstract:

last modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnail URL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

mimeType: dataType: identifier:

fullDescriptionUnstructured: Mide 16.5 cm. Cola larga (8.9 cm). Plumas centrales de la cola puntiagudas. Corona castaño rojizo listado con negro; ceja y barbilla blancas. Fuertemente rayado con marrón oscuro y blanco por encima y por debajo; alas y cola café oscuro; plumas externas de la cola con bordes pálidos (Hilty & Brown, 1986). briefDescription:

references

link: http://www.birdlife.org/datazone/species/factsheet/22702192

keywords: Leptasthenura andicola

editors

institution: address: publisher: accessed: websites: chapter: series: issue: year: 2014

authors

authors: BirdLife International

title: Species factsheet: Leptasthenura andicola

source:

type: miscellany

tags: Leptasthenura andicola

abstract:

identifiers

last_modified: created: group_id: profile_id: id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo: providerLink:

publicationStatus

microReference:

source: datatype: identifier: source: datatype:

note: ruleConsidered:
replacementNameFor
microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered:
sanctioned
microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered:
laterHomonymOf
microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered:
conservedAgainst
basedOn microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: basionym microReference:
source: datatype:

identifier:

identifier: source: datatype: identifier:
note: ruleConsidered:
spellingCorrentionOf
typeNameEntity
source: datatype: identifier: source: datatype: identifier: source: datatype: identifier:
typificacion
typeOfType:
lectotypeMicroReferenceVoucher
lectotypePublicationVoucher
voucherReference
simple: microReference: year:
publishedIn
simple: source: datatype: identifier:
specialAuthorship
authors
year
simple:
canonicalAuthorship
authors
year
simple:

simple: P. L. Sclater, 1870

canonicalName

infraspecificEpithet: specificEpithet: infragenericEpithet:

linkType: ref: uninomial:

simple: Leptasthenura andicola

rank: SPECIES

simple: Leptasthenura andicola P. L. Sclater, 1870

attributes

nomenclaturalCode: isAnamorphic:

id:

synonymsAtomized

hierarchy

ancillaryData

dataObject:

additionalInformation:

reference

geoPoint: location: thumbnailURL:

mediaURL

description:

subject

source: The Catalogue of Life, 3rd January 2011

audience

bibliographicCitation:

rigthsHolder: rights:

license: title:

modified: 2016-02-21T20:44:54.168Z created: 2016-02-21T20:44:54.168Z

agent

homepage:

role: email: phone: address: position: organisation: lastName: firstName: mimeType: dataType: identifier: parentTaxon: Leptasthenura higherClassification: infraspecificEpithet: specificEpithet: taxonRank: SPECIES subGenus: genus: Leptasthenura family: Furnariidae order: Passeriformes classHierarchy: Aves phylum: Chordata kingdom: Animalia recommended: classification: invasiveness invasivenessUnstructured: reproduction 0 dataObject: additionalInformation: link: http://www.hbw.com/node/56411 keywords: Leptasthenura andicola 0: del Hoyo, J 1: Elliott, A 2: Sargatal, J 3: Christie, D A 4: de Juana, E institution: address: publisher: Lynx Edicions accessed: websites: chapter: series: issue: year: 2003 0: Remsen, J.V., Jr title: Andean Tit-spinetail (Leptasthenura andicola) source: Handbook of the Birds of the World Alive type: book_section tags: Leptasthenura andicola

abstract: last_modified:

created: group_id:
profile_id: id:
geoPoint:
location: thumbnailURL:
description:
source: bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created: mimeType:
dataType:
identifier: reproductionUnstructured: Es una especie presumiblemente monógama. El nido es una bola elaborada con
fragmentos de plantas de 16 cm de diámetro, entrada lateral y cámara profusamente llena de plumas y pelo. La hembra pone dos huevos (Remsen, 2003).
molecularData
molecularDataUnstructured:
migratory
migratoryUnstructured:
lifeForm
lifeFormUnstructured:
lifeCycle
lifeCycleUnstructured:
interactions
interactionsUnstructured:
feeding
0
dataObject: additionalInformation:
0
link: http://www.hbw.com/node/56411 keywords: Leptasthenura andicola
0: del Hoyo, J 1: Elliott, A
2: Sargatal, J
3: Christie, D A 4: de Juana, E
institution:
address: publisher: Lynx Edicions

accessed: websites: chapter: series: issue: year: 2003 0: Remsen, J.V., Jr title: Andean Tit-spinetail (Leptasthenura andicola) source: Handbook of the Birds of the World Alive type: book_section tags: Leptasthenura andicola abstract: last_modified: created: group_id: profile_id: id: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rights: Asociación Bogotana de Ornitología license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: feedingUnstructured: Consume una variedad de artrópodos. Forrajea en todos los niveles y por lo general
permanece en parejas o en pequeños grupos, de vez en cuando asociado a bandadas mixtas. Rebusca en el follaje de los arbustos, las ramas pequeñas y sus epífitas, en el follaje e inflorescencias de Espeletia, y en flores (especialmente de Senecio); de vez en cuando se cuelga boca abajo acrobáticamente (Remsen, 2003).
environmentalEnvelope
environmentalEnvelopeUnstructured:
ecologicalSignificance
ecologicalSignificanceUnstructured:
dispersal
dispersalUnstructured: distance: structureDispersed: type: purpose:
behavior
0
dataObject: additionalInformation:

keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis.Coccycua pumila.Coccyzus americanus,Coccyzus melacoryphus,Coeligena bonapartei,Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta caerulea, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria guitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor,Ochthoeca rufipectoralis,Orochelidon murina,Oxypogon guerinii,Oxyura jamaicensis,Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus,Ramphomicron microrhynchum,Riparia riparia,Rupornis magnirostris,Sayornis nigricans,Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis institution:

address:

accessed:

publisher: Princeton University Press

websites: chapter: series: pages: 836 issue: year: 1986 0: Hilty, Steven 1: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis.Catamenia analis.Catamenia inornata.Cathartes aura.Catharus minimus.Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta caerulea, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria guitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus. Nycticorax nycticorax. Ochthoeca cinnamomeiventris. Ochthoeca diadema. Ochthoeca fumicolor,Ochthoeca rufipectoralis,Orochelidon murina,Oxypogon guerinii,Oxyura jamaicensis,Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus,Ramphomicron microrhynchum,Riparia riparia,Rupornis magnirostris,Sayornis nigricans,Serpophaga cinerea. Setophaga castanea. Setophaga cerulea. Setophaga petechia. Setophaga ruticilla. Setophaga striata. Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis abstract:

abstract: last_modified: created: group_id: profile id:

id:

1

link: http://www.hbw.com/node/56411 keywords: Leptasthenura andicola

0: del Hoyo, J 1: Elliott, A 2: Sargatal, J 3: Christie, D A

4: de Juana, E institution: address: publisher: Lynx Edicions accessed: websites: chapter: series: issue: year: 2003 0: Remsen, J.V., Jr tttle: Andean Tit-spinetail (Leptasthenura andicola) source: Handbook of the Birds of the World Alive type: book_section tags: Leptasthenura andicola abstract: last_modified: created: group_id: profile_id: id: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rights-Holder: rights-Asociación Bogotana de Ornitología license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: detaTiype: identifier: behaviorUnstructured: La especie prefiere permanecer furtivo y algunas veces nervioso desciende cortas distancias y desaparece rápidamente. Algunas veces con bandadas mixtas o en pequeños grupos con Mecocerculus
ancillaryData
ancillaryData dataObject:
additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified:

created: mimeType: dataType: identifier:

measurementOrFact

relatedTo:

measurementRemarks: measurementMethod:

measurementDeterminedDate:

measurementUnit: measurementAccuracy: measurementValue: measurementType: measurementID:

endemicAtomized

distribution

ancillaryData

dataObject:

additionalInformation:

reference

link: http://maps.iucnredlist.org/map.html?id=22702192

keywords: Leptasthenura andicola

editors

institution: address: publisher: accessed: websites: chapter: series: issue: year: 2014

authors

authors: BirdLife International

authors: NatureServe

title: Bird Species Distribution Maps of the World. 2012. Leptasthenura andicola. The IUCN Red List of Threatened

Species. Version 2014.3

source:

type: miscellany

tags: Leptasthenura andicola

abstract:

identifiers

last_modified: created: group_id:

profile_id: id: geoPoint: location: thumbnailURL:
mediaURL
description:
subject
source:
audience
bibliographicCitation: rightsHolder: rights: Asociación Bogotana de Ornitología license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created:
agent
mimeType: dataType: identifier: distributionUnstructured: La especie se distribuye principalmente en Sudamérica desde el norte de Venezuela hasta el norte de Bolivia. En Colombia se registra en Santa Marta, Cundinamarca y la cordillera Central desde Tolima (Nevado del Ruiz) hasta Nariño (BirdLife International and NatureServe, 2014; Hilty & Brown, 1986). La especie se encuentra principalmente entre los 3000 y los 4500 msnm (Hilty & Brown, 1986).
distributionAtomized
stateProvince: Tolima
ancillaryData
locality: municipality: country: Colombia
temporalCoverage
endDate: startDate:
distributionScope
type:

ancillaryData

threatStatus

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES: locality:

municipality: county: stateProvince: country: Global 0: IUCN measurementValue: Preocupación menor (LC) relatedTo: measurementRemarks: measurementMethod: measurementDeterminedBy measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementType: IUCN measurementID: territory territoryUnstructured: areaOfOccupancy: extentOfOccurrence: populationBiology dataObject: additionalInformation: 0 link: http://www.birdlife.org/datazone/species/factsheet/22702192 keywords: Leptasthenura andicola institution: address: publisher: accessed: websites: chapter: series: issue: year: 2014 0: BirdLife International title: Species factsheet: Leptasthenura andicola source: type: miscellany tags: Leptasthenura andicola abstract: last_modified: created: group_id: profile_id: id: geoPoint: location: thumbnailURL:

description: source: bibliographicCitation: rightsHolder: rights: Asociación Bogotana de Ornitología license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: populationBiologyUnstructured: El tamaño de la población mundial no se ha cuantificado, pero esta especie se ha descrito como "bastante común" (BirdLife International, 2014).
legislation
legislationUnstructured:
directThreats
directThreatsUnstructured:
ancillaryData
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified: created: mimeType: dataType: identifier:
measurementOrFact
relatedTo: measurementRemarks: measurementMethod: measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: measurementType: measurementID: abstract: El Coludito Paramuno es una especie de furnárido originario de Sudamérica y que se distribuye desde Venezuela hasta Bolivia (BirdLife International and NatureServe, 2014; Hilty & Brown, 1986). Prefiere áreas

Venezuela hasta Bolivia (BirdLife International and NatureServe, 2014; Hilty & Brown, 1986). Prefiere áreas húmedas especialmente con arbustos dispersos y Espeletia o en los bordes de bosque de Polylepis; también en bosques achaparrados en el ecotono con el páramo, principalmente entre los 3000 y 4500 msnm (Hilty & Brown, 1986). Consume una variedad de artrópodos. Forrajea en todos los niveles y por lo general permanece en parejas o en pequeños grupos, de vez en cuando asociado a bandadas mixtas (Remsen, 2003).

associatedParty

role:

personnelldentifier: personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone: postalCode: country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address: organisation: position:

lastName: Gómez Melendro firstName: Edith Natalia

ancillaryData

dataObject:

additionalInformation:

reference

geoPoint: location: thumbnailURL:

mediaURL: http://www.xeno-canto.org/37083

description:

subject

source: http://www.xeno-canto.org

audience

bibliographicCitation:

rightsHolder:

rights:

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

agent

mimeType: dataType: identifier:

_id

_bsontype: ObjectID id: VÊ4Oð Ig#çE€