Reporte de Fichas para el usuario: edith gómez Ficha ID: 56d5c5653c16479905cba8f8 Fecha de creación: 2016-03-01T16:37:59.069Z annualCycles annualCycleUnstructured: habitats 0 dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Equipo Coordinador del SiB Colombia license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: habitatUnstructured: Residente de invierno, se encuentran en números muy pequeños en estuarios, manglares, lagos y pantanos de agua dulce (Hilty & Brown 1986). commonNamesAtomized ancillaryData usedBy: usedIn distributionUnstructured: endDate: startDate: ancillaryData type: language: Español name: Garza morena creation date: 2016-03-01T16:37:59.069Z usesManagementAndConservation ancillaryData

managementAndConservationUnstructured:

managementAndConservationAtomized

identificationKeys

#### fullDescription

0

dataObject:

additionalInformation:

geoPoint: location:

thumbnailURL: description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created: mimeType: dataType: identifier:

fullDescriptionUnstructured: Mide 132 cm. y pesa 2.5 kg. Es la garza más grande de Costa Rica, de aleteo más lento. Los adultos son de cabeza blanca. Las anchas rayas negras a los lados de la coronilla se extienden hasta la parte posterior de la nuca y las largas plumas occipitales. El cuello es gris parduzco con una raya blanca y negra por el centro de la parte delantera. El cuerpo y las alas son principalmente gris azulado, con un parche negro al lado del pecho. Presentan rayas blancas por debajo y los muslos rojizos. El iris es amarillo, el pico es amarillento y las patas son negruzcas (Elizondo, 2000).

briefDescription:

#### references

#### taxonRecordName

## ancillaryData

providerSpecificData

anyTwo: providerLink:

publicationStatus

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

replacementNameFor

microReference:

source: datatype: identifier:

datatype: identifier: note: ruleConsidered:
sanctioned
microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered:
laterHomonymOf
microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered:
conservedAgainst
basedOn
microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered:
basionym
microReference: source: datatype: identifier: source: datatype: identifier: note:
ruleConsidered:

source:

spellingCorrentionOf

typeNameEntity
source: datatype: identifier: source: datatype: identifier: source: datatype: identifier: source: datatype: identifier:
typificacion
typeOfType:
lectotypeMicroReferenceVoucher
lectotypePublicationVoucher
voucherReference
simple: microReference: year:
publishedIn
simple: Linnaeus, C. (1758). Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Editio decima, reformata. Laurentius Salvius: Holmiae. ii, 824 pp. source: datatype: identifier:
specialAuthorship
authors
year
simple:
canonicalAuthorship
authors
year
simple: simple: Linnaeus, 1758
canonicalName
infraspecificEpithet: specificEpithet: infragenericEpithet: linkType: ref:

uninomial:

simple: Ardea herodias

rank: SPECIES

simple: Ardea herodias Linnaeus, 1758

attributes

nomenclaturalCode: isAnamorphic:

id:

# synonymsAtomized

# hierarchy

# ancillaryData

dataObject:

additionalInformation:

### reference

geoPoint: location: thumbnailURL:

#### mediaURL

description:

# subject

source: The Catalogue of Life, 3rd January 2011

### audience

bibliographicCitation:

rigthsHolder:

rights: license: title:

modified: 2016-03-01T16:37:39.069Z created: 2016-03-01T16:37:39.069Z

### agent

homepage:

role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:

parentTaxon: Ardea

higherClassification: infraspecificEpithet: specificEpithet:

taxonRank: SPECIES

subGenus: genus: Ardea family: Ardeidae order: Pelecanifo

order: Pelecaniformes classHierarchy: Aves phylum: Chordata kingdom: Animalia recommended: classification:

invasiveness

invasivenessUnstructured:

reproduction

reproductionUnstructured:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

feedingUnstructured:

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type: purpose:

behavior

dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Equipo Coordinador del SiB Colombia license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: behaviorUnstructured: Generalmente solitario. Estático o vadeando en aguas someras (Hilty & Brown 1986).
ancillaryData
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified: created: mimeType: dataType: identifier:
measurementOrFact
relatedTo: measurementRemarks: measurementMethod: measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: measurementType: measurementID:
endemicAtomized
distribution
ancillaryData

dataObject: additionalInformation:

#### reference

geoPoint: location:

thumbnailURL:

#### mediaURL

description:

## subject

source:

#### audience

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

#### agent

mimeType: dataType: identifier:

distributionUnstructured: En bases de datos y monitoreos de fauna se encontró que:

Esta especie se distribuye en: Estados Unidos, Korea (Arctos 2012).

Para las siguientes subespecies la distribución es:

Ardea herodias cognata Ecuador (Arctos 2012)

Ardea herodias fannini Canadá, Estados Unidos (Arctos 2012)

Ardea herodias herodias Canadá, Colombia, Estados Unidos, México (Arctos 2012; Collections MNH SI 2012)

Ardea herodias hyperonca Estados Unidos (Arctos 2012) Ardea herodias lessonii Colombia (Collections MNH SI 2012)

Ardea herodias occidentalis Estados Unidos (Arctos 2012)

Ardea herodias sanctilucae México (Arctos 2012)

Ardea herodias treganzai Estados Unidos, México (Arctos 2012)

Ardea herodias wardi Estados Unidos (Arctos 2012)

En bases de datos y monitoreos de fauna se encontró que:

Esta especie se distribuye en: La Guajira (Cerrejón 2009-2010).

Para las siguientes subespecies la distribución en Colombia es:

Ardea herodias herodias Cundinamarca (ICN 2004)

Ardea herodias lessonii Magdalena (Collections MNH SI 2012), en las regiones Andina y Caribe de Colombia hasta 2600 m (Hilty & Brown 1986).

#### distributionAtomized

stateProvince: La Guajira

# ancillaryData

locality: municipality: country: Colombia

temporalCoverage

endDate:

startDate:
distributionScope
type:
threatStatus
ancillaryData
threatStatusUnstructured:
threatStatusAtomized
apendiceCITES: locality: municipality: county: stateProvince: country: Global 0: IUCN measurementValue: Preocupación menor (LC) relatedTo: measurementRemarks: measurementMethod:
measurementDeterminedBy
measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementType: IUCN measurementID:
territory
territoryUnstructured: areaOfOccupancy: extentOfOccurrence:
populationBiology
populationBiologyUnstructured:
legislation
legislationUnstructured:
directThreats
directThreatsUnstructured:
ancillaryData
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation:

rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

سماملما المقطية

#### measurementOrFact

relatedTo:

measurementRemarks: measurementMethod:

measurementDeterminedDate:

measurementUnit: measurementAccuracy: measurementValue: measurementType: measurementID:

abstract: Mide de 102 a 130 cm. Esbelta y de cuello largo. Pico amarillo opaco; patas amarillo verdoso oscuro. Cabeza y garganta blancas, área occipital y lados de la coronilla negros (también la cresta occipital elongada en plumaje nupcial); cuello grisáceo, estriado al frente de negro y blanco, resto de las partes dorsales y alas gris azuloso; las partes ventrales más o menos estriadas de negro y blanco; tibias castaño (Hilty & Brown 1986).

# associatedParty

role:

personnelldentifier: personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone: postalCode: country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address: organisation: position:

lastName: Gómez Melendro firstName: Edith Natalia

# ancillaryData

dataObject:

additionalInformation: Registros biológicos: Portal de datos GBIF

Otros recursos en Internet: Arctos, Avibase, AVoCet, Colecciones en Línea, EoL, iSpecies, IUCN Red List, SI NMNH

xeno-canto

Créditos específicos: Ficha elaborada con el apoyo de Carbones del Cerrejón Limited.

#### reference

geoPoint: location: thumbnailURL:

# mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder:
rights: Equipo Coordinador del SiB Colombia
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
agent
mimeType:
dataType:
identifier:
\_id

\_bsontype: ObjectID id: VÕÅe< G™\0•€