

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56c398e7f0106c67230e7351

Fecha de creación: undefined

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

0

keywords: Actitis macularius,Ampelion rubrocristatus,Anas discors,Anisognathus igniventris,Ardea alba,Arremon assimilis,Asio flammeus,Asio stygius,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Catharus fuscescens,Catharus ustulatus,Chaetocercus mulsant,Chrysomus icterocephalus

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: Actitis macularius,Ampelion rubrocristatus,Anas discors,Anisognathus igniventris,Ardea alba,Arremon assimilis,Asio flammeus,Asio stygius,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Catharus fuscescens,Catharus ustulatus,Chaetocercus mulsant,Chrysomus icterocephalus

abstract:

last_modified:

created:

group_id:

profile_id:

id:

1

maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

pages: 836

issue:

year: 1986

0: Hilty, Steven

1: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Agelaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigristrostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coccyzus pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiostyris alpestris, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Mactornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striatocollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerini, Oxyura jamaicensis, Pachyrhamphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus

unicolor,Phyllomyias uropygialis,Piaya cayana,Picoides fumigatus,Pipraeidea melanonota,Pipreola riefferii,Piranga olivacea,Piranga rubra,Pitangus sulphuratus,Pluvialis dominica,Podilymbus podiceps,Porphyrion martinicus,Porzana carolina,Progne tapera,Protonotaria citrea,Pseudocolopteryx acutipennis,Pseudoscops clamator,Pterophanes cyanopterus,Pygochelidon cyanoleuca,Pyrocephalus rubinus,Pyrrhomyias cinnamomeus,Quiscalus lugubris,Rallus semiplumbeus,Ramphocelus dimidiatus,Ramphomicron microrhynchum,Riparia riparia,Rupornis magnirostris,Sayornis nigricans,Serpophaga cinerea,Setophaga castanea,Setophaga cerulea,Setophaga petechia,Setophaga ruticilla,Setophaga striata,Sicalis flaveola,Sicalis luteola,Spiza americana,Sporagra spinescens,Sturnella magna,Synallaxis subpudica,Systellura longirostris,Tangara vitriolina,Thraupis episcopus,Thraupis palmarum,Tringa flavipes,Tringa melanoleuca,Tringa solitaria,Troglodytes aedon,Troglodytes solstitialis,Turdus fuscater,Turdus ignobilis,Tyrannus melancholicus,Tyrannus savana,Tyrannus tyrannus,Tyto alba,Vanellus chilensis,Vireo olivaceus,Vultur gryphus,Zenaida auriculata,Zonotrichia capensis

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

habitatUnstructured: La Monjita es común en cuerpos de agua dulce, tierras inundadas y orillas de los ríos, pastizales de várzea, tierras agrícolas como arrozales y especialmente en regiones abiertas (Fraga, 2011; Hilty & Brown, 1986). En la Sabana de Bogotá prefiere los humedales con abundantes juncos (Scirpus), con frecuencia se le ve parado en ellos o recorriéndolos (Asociación Bogotana de Ornitología, 2000).

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

[ancillaryData](#)

type:

language: Inglés

name: Yellow-hooded Blackbird

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

managementAndConservationAtomized

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

0

keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus

abstract:

last_modified:

created:

group_id:

profile_id:

id:

1

link: <http://www.hbw.com/species/yellow-hooded-blackbird-chrysomus-icterocephalus>

keywords: Chrysomus icterocephalus

0: del Hoyo, J

1: Elliott, A

2: Sargatal, J

3: Christie, D A

4: de Juana, E

institution:

address:

publisher: Lynx Edicions

accessed:

websites:

chapter:

series:

issue:

year: 2011

0: Fraga, R

title: Yellow-hooded Blackbird (Chrysomus icterocephalus)

source: Handbook of the Birds of the World Alive

type: book_section

tags: Chrysomus icterocephalus

abstract:

last_modified:

created:

group_id:

profile_id:

id:

2

keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigristrostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidnucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coccyzus pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliosus, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Mactornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striatocollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerini, Oxyura jamaicensis, Pachyrhamphus versicolor, Pandion haliaetus, Pardirallus

maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

pages: 836

issue:

year: 1986

0: Hilty, Steven

1: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Agelaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigristrostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coccyzus pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiostyris peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Mactornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striatocollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerinii, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus

unicolor,Phyllomyias uropygialis,Piaya cayana,Picoides fumigatus,Pipraeidea melanonota,Pipreola riefferii,Piranga olivacea,Piranga rubra,Pitangus sulphuratus,Pluvialis dominica,Podilymbus podiceps,Porphyrion martinicus,Porzana carolina,Progne tapera,Protonotaria citrea,Pseudocolopteryx acutipennis,Pseudoscops clamator,Pterophanes cyanopterus,Pygochelidon cyanoleuca,Pyrocephalus rubinus,Pyrrhomyias cinnamomeus,Quiscalus lugubris,Rallus semiplumbeus,Ramphocelus dimidiatus,Ramphomicron microrhynchum,Riparia riparia,Rupornis magnirostris,Sayornis nigricans,Serpophaga cinerea,Setophaga castanea,Setophaga cerulea,Setophaga petechia,Setophaga ruticilla,Setophaga striata,Sicalis flaveola,Sicalis luteola,Spiza americana,Sporagra spinescens,Sturnella magna,Synallaxis subpudica,Systellura longirostris,Tangara vitriolina,Thraupis episcopus,Thraupis palmarum,Tringa flavipes,Tringa melanoleuca,Tringa solitaria,Troglodytes aedon,Troglodytes solstitialis,Turdus fuscater,Turdus ignobilis,Tyrannus melancholicus,Tyrannus savana,Tyrannus tyrannus,Tyto alba,Vanellus chilensis,Vireo olivaceus,Vultur gryphus,Zenaida auriculata,Zonotrichia capensis

abstract:

last_modified:

created:

group_id:

profile_id:

id:

3

link: <http://www.museum.lsu.edu/{~}Remsen/SACCprop135.htm>

keywords: Chrysomus icterocephalus

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2004

0: Schulenberg, Tom

title: Proposal (135) to South American Classification Committee

source: South American Classification Committee

type: miscellany

tags: Chrysomus icterocephalus

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

fullDescriptionUnstructured: La Monjita mide entre 17 y 19 cm, la hembra es un poco más pequeña, 16 cm en promedio (Asociación Bogotana de Ornitología, 2000; Hilty & Brown, 1986); el macho tiene un peso promedio de 35.9 g y la hembra de 26.8 g. En la subespecie *C. i. icterocephalus* el macho tiene la cabeza, garganta y la parte superior del pecho amarillo brillante, el área loreal y el resto de plumaje es negro brillante (Asociación Bogotana de Ornitología, 2000; Fraga, 2011); el iris es marrón oscuro; pico cónico, agudo y negro (Hilty & Brown, 1986); patas negras. La hembra tiene área superciliar, garganta y la parte alta del pecho amarillo, el resto de la cara y la corona es amarillo verdoso. Las partes superiores son gris-oliva opaco con rayas más oscuras, las alas y cola son oscuras, café-oliva; las partes inferiores son oliva, el vientre un poco más oscuro con oliva ligeramente veteado; pico negruzco, a menudo más pálido en la base de la mandíbula inferior. El juvenil es como la hembra, pero más opaco, con menos amarillo en la cabeza, dorso más pálido, las partes inferiores moteadas; el macho inmaduro es como la hembra, pero con un amarillo más brillante y extenso en la garganta, puede mostrar plumas negras en las partes inferiores y el dorso. La subespecie *C. i. bogotensis*, es más grande que la nominal, la hembra es más oscura, con cabeza menos amarilla, mascara sobre las lores hasta el área del oído oscuro o negruzco fuertemente contrastante, tiene un estrecho contorno desde la línea superciliar y por el área del oído y se reconoce el babero amarillo claramente definido (Fraga, 2011). El macho de esta especie es inconfundible, la hembra puede tener un patrón similar al del macho pero es mucho más opaca (Hilty & Brown, 1986). Anteriormente en el género *Agelaius*, cambio al género *Chrysomus* según la propuesta 135 al South American Classification Committee (Schulenberg, 2004). Datos moleculares y comportamentales indican que esta especie está estrechamente relacionada con *C. ruficapillus* (Fraga, 2011).

briefDescription:

references

keywords: *Actitis macularius*, *Ampelion rubrocristatus*, *Anas discors*, *Anisognathus igniventris*, *Ardea alba*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bartramia longicauda*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Catharus fuscescens*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chrysomus icterocephalus*

editors

editors: Stiles, F G

editors: Bohórquez, C I

editors: Cadena, C D

editors: De la Zerda, S

editors: Hernández, M

editors: Roselli, L

editors: Kelsey, M

editors: Valencia, I D

editors: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

authors

authors: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book
tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus
abstract:

identifiers

last_modified:
created:
group_id:
profile_id:
id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:
providerLink:

publicationStatus

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

replacementNameFor

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:

ruleConsidered:

laterHomonymOf

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

conservedAgainst

basedOn

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

basionym

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:

datatype:

identifier:

source:

datatype:

identifier:

source:

datatype:

identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:
canonicalAuthorship

authors

year

simple:
simple: (Linnaeus, 1766)

canonicalName

infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:
ref:

uninomial:
simple: Chrysomus icterocephalus
rank: SPECIES
simple: Chrysomus icterocephalus (Linnaeus, 1766)

attributes

nomenclaturalCode:
isAnamorphic:
id:

synonymsAtomized

ancillaryData

synonymStatus:
synonymName
anyTwo:

anyOne

providerLink:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:

ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typeVoucherEntity

typeOfType:
simple:
microReference:
year:
source:
datatype:
identifier:

[combinationAuthorship](#)

basionymAuthorship

simple:
authorship

simple:
simple: (Linnaeus, 1766)

epithet

infraspecificEpithet:
specificEpithet:
infragenericEpithet:

genus

linkType:
ref:
uninomial:
simple: Agelaius icterocephalus
rank: SPECIES
simple: Agelaius icterocephalus (Linnaeus, 1766)
nomenclaturalCode:
isAnamorphic:
id:

[hierarchy](#)

[ancillaryData](#)

dataObject:
additionalInformation:

[reference](#)

geoPoint:
location:
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:
rightsHolder:
rights:
license:
title:
modified: 2016-02-16T21:32:23.798Z
created: 2016-02-16T21:32:23.798Z

[agent](#)

homepage:
role:
email:
phone:
address:
position:

organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:
parentTaxon: Chrysomus
higherClassification:
infraspecificEpithet:
specificEpithet:
taxonRank: SPECIES
subGenus:
genus: Chrysomus
family: Icteridae
order: Passeriformes
classHierarchy: Aves
phylum: Chordata
kingdom: Animalia
recommended:
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:
additionalInformation:

0

keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus

abstract:

last_modified:

created:

group_id:

profile_id:

id:

1

link: <http://www.hbw.com/species/yellow-hooded-blackbird-chrysomus-icterocephalus>

keywords: Chrysomus icterocephalus

0: del Hoyo, J

1: Elliott, A

2: Sargatal, J

3: Christie, D A

4: de Juana, E

institution:

address:

publisher: Lynx Edicions

accessed:

websites:

chapter:

series:

issue:

year: 2011

0: Fraga, R

title: Yellow-hooded Blackbird (Chrysomus icterocephalus)

source: Handbook of the Birds of the World Alive

type: book_section

tags: Chrysomus icterocephalus

abstract:

last_modified:

created:

group_id:

profile_id:

id:

2

keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coccycua

pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreovertris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerini, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

pages: 836

issue:

year: 1986

0: Hilty, Steven

1: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Agelaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coccyzus pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus

cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreovertris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerinii, Oxyura jamaicensis, Pachyrhamphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

reproductionUnstructured: En algunos territorios la temporada de reproducción está altamente sincronizada con las estaciones lluviosas. La Monjita presenta poliginia, un macho tiene un pequeño harem de entre dos y cinco hembras y territorios de un promedio de 308 m². Se ha observado algunas colonias monógamas. Anidan en colonias, el tamaño de la colonia es variable, se han encontrado colonias con hasta 100 nidos en zonas de marismas o en campos de arroz, también se observan algunos nidos solitarios o dispersos en semicolonias (Fraga, 2011). El nido es construido por los machos (Fraga, 2011), es una taza tejida con hojas de pasto (Asociación Bogotana de Ornitología, 2000), con diámetro y profundidad de aproximadamente 10 cm (Fraga, 2011), se encuentra entre 30 y

50 cm por encima del agua entre los juncos u otra vegetación acuática (Asociación Bogotana de Ornitología, 2000; Hilty & Brown, 1986), rara vez en árboles o arbustos (Fraga, 2011). Los nidos se encuentran agrupados en colonias dispersas. Ponen de tres a cuatro huevos azul claro punteados y garabateados de negro (Asociación Bogotana de Ornitología, 2000; Hilty & Brown, 1986), su dimensión promedio es de 16 x 22 mm (Fraga, 2011). Los huevos son incubados por las hembras, la eclosión es asincrónica, entre los siguientes 13 días, la hembra es la encargada de alimentar los polluelos a veces con ayuda del macho, aunque el macho se encarga principalmente de proteger el nido contra depredadores y parásitos de cría. Cuando los nidos son parasitados por más de tres huevos, las hembras de Monjita suelen abandonar el nido, por lo que un bajo porcentaje de nidos parasitados producen polluelos de tal parasito (Fraga, 2011). Una vez el macho construye el nido la hembra decide si lo acepta o no, de ser así, el macho permanece con ella hasta los primeros días de incubación, sin embargo, muchos nidos se quedan sin utilizar dado que muchos machos no logran que una hembra los acepte (Fraga, 2011).

0

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

[audience](#)

rightsHolder:

rights:

license:

modified:

created:

[agent](#)

mimeType:

dataType:

identifier:

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedBy: TraitBank

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue: Sexual

measurementType: Tipo de Reproducción

measurementID:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

0

dataObject:

additionalInformation:

0

issn: 1474-919X

keywords: Chrysomus icterocephalus

institution:

address:

publisher: Wiley Online Library

accessed:

websites:

chapter:

series:

pages: 436--444

issue:

volume: 132

year: 1990

0: Cruz, Alexander

1: Manolis, Timothy D

2: Andrews, Robert W

title: Reproductive interactions of the Shiny Cowbird *Molothrus bonariensis* and the Yellow hooded Blackbird *Agelaius icterocephalus* in Trinidad

source: Ibis

type: article

tags: Chrysomus icterocephalus

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

interactionsUnstructured: La Monjita suele ser una de las especies parasitadas por el Chamón o Tordo, *Molothrus bonariensis*, sin embargo se ha observado que esta especie suele reconocer los nidos parasitados y en un muchas ocasiones abandona estos nidos (Cruz et al., 1990).

0

0

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source: https://docs.google.com/document/d/1i3mLz_QRu3Ds4Z19EaJM1iNS1XAv8gObGB17ASur8EM/edit?usp=sharing

audience

bibliographicCitation: Futuyma, D.J. 1998. Evolutionary Biology, Third Edition. Sinauer Associates, Inc. EUA
rightsHolder:
rights:
license:
modified:
created:

agent

contentType:
dataType:
identifier:
relatedTo:
measurementRemarks:
measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue: Parasitismo
measurementType: CONABIO
measurementID:
interactionSpecies: *Molothrus bonariensis*

feeding

0

dataObject:
additionalInformation:

0

keywords: *Actitis macularius*, *Ampelion rubrocristatus*, *Anas discors*, *Anisognathus igniventris*, *Ardea alba*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bartramia longicauda*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Catharus fuscescens*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chrysomus icterocephalus*

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: *Actitis macularius*, *Ampelion rubrocristatus*, *Anas discors*, *Anisognathus igniventris*, *Ardea alba*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bartramia longicauda*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Catharus fuscescens*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chrysomus icterocephalus*

abstract:

last_modified:

created:

group_id:

profile_id:

id:

1

link: <http://www.hbw.com/species/yellow-hooded-blackbird-chrysomus-icterocephalus>

keywords: *Chrysomus icterocephalus*

0: del Hoyo, J

1: Elliott, A

2: Sargatal, J

3: Christie, D A

4: de Juana, E

institution:

address:

publisher: Lynx Edicions

accessed:

websites:

chapter:
series:
issue:
year: 2011
0: Fraga, R
title: Yellow-hooded Blackbird (*Chrysomus icterocephalus*)
source: Handbook of the Birds of the World Alive
type: book_section
tags: *Chrysomus icterocephalus*
abstract:
last_modified:
created:
group_id:
profile_id:
id:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
feedingUnstructured: Se alimentan de insectos y semillas, forrajean buscando insectos entre los juncas pero también buscan por fuera de ellos una buena parte de su alimento, frecuentemente en áreas de vegetación baja como la lengua de vaca (*Rumex*) (Asociación Bogotana de Ornitología, 2000). Se le ha observado consumiendo arroz, orugas, gorgojos y saltamontes. Las hembras consiguen alimento para los polluelos fuera del territorio del macho, la subespecie *C. i. bogotensis* no sale de su territorio para buscar alimento (Fraga, 2011).

0

type: Heterótrofos - Omnívoros

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

0

keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus

abstract:

last_modified:

created:

group_id:

profile_id:

id:

1

link: <http://www.hbw.com/species/yellow-hooded-blackbird-chrysomus-icterocephalus>

keywords: Chrysomus icterocephalus

0: del Hoyo, J

1: Elliott, A

2: Sargatal, J

3: Christie, D A

4: de Juana, E

institution:

address:

publisher: Lynx Edicions
accessed:
websites:
chapter:
series:
issue:
year: 2011
0: Fraga, R
title: Yellow-hooded Blackbird (*Chrysomus icterocephalus*)
source: Handbook of the Birds of the World Alive
type: book_section
tags: *Chrysomus icterocephalus*
abstract:
last_modified:
created:
group_id:
profile_id:
id:

2

keywords: *Accipiter striatus*, *Actitis macularius*, *Adelomyia melanogenys*, *Agelaiocercus kingii*, *Amblycercus holosericeus*, *Ampelion rubrocristatus*, *Anas andium*, *Anas clypeata*, *Anas discors*, *Andigena nigristrois*, *Anisognathus igniventris*, *Ardea alba*, *Ardea herodias*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Asthenes flammulata*, *Asthenes fuliginosa*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Butorides virescens*, *Cairina moschata*, *Calidris bairdii*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Cathartes aura*, *Catharus minimus*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chaetura pelagica*, *Chalcostigma heteropogon*, *Chordeiles minor*, *Chrysomus icterocephalus*, *Cinclodes albidiventris*, *Cinclus leucocephalus*, *Cinnycerthia unirufa*, *Cistothorus platensis*, *Coccyzua pumila*, *Coccyzus americanus*, *Coccyzus melacoryphus*, *Coeligena bonapartei*, *Colibri coruscans*, *Colibri thalassinus*, *Columba livia*, *Columbina talpacoti*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cinereus*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Dendrocygna autumnalis*, *Dendrocygna viduata*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dolichonyx oryzivorus*, *Dubusia taeniata*, *Egretta thula*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Geothlypis philadelphia*, *Geranoaetus melanoleucus*, *Glaucidium jardinii*, *Grallaria quitesis*, *Grallaria ruficapilla*, *Grallaria squamigera*, *Haplospiza rustica*, *Heliangelus amethysticollis*, *Heliangelus exortis*, *Hellmayrea gularis*, *Hemispingus superciliaris*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus icterus*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Lafresnaya lafresnayi*, *Leiostyris alpestris*, *Leptasthenura andicola*, *Lesbia nuna*, *Lesbia victoriae*, *Machetornis rixosa*, *Margarornis squamiger*, *Mecocerculus leucophrys*, *Megascops choliba*, *Melanerpes rubricapillus*, *Merganetta armata*, *Metallura tyrianthina*, *Milvago chimachima*, *Mimus gilvus*, *Mniotilta varia*, *Molothrus bonariensis*, *Muscisaxicola alpinus*, *Myiarchus crinitus*, *Myiodynastes luteiventris*, *Myiotheretes fumigatus*, *Myiotheretes striaticollis*, *Myiothlypis nigrocristata*, *Nomonyx dominicus*, *Nycticorax nycticorax*, *Ochthoeca cinnamomeiventris*, *Ochthoeca diadema*, *Ochthoeca fumicolor*, *Ochthoeca rufipectoralis*, *Orochelidon murina*, *Oxygogon guerini*, *Oxyura jamaicensis*, *Pachyrhamphus versicolor*, *Pandion haliaetus*, *Pardirallus maculatus*, *Parkesia noveboracensis*, *Patagioenas fasciata*, *Petrochelidon pyrrhonota*, *Phaetusa simplex*, *Phalacrocorax brasilianus*, *Pheucticus aureoventris*, *Pheucticus ludovicianus*, *Phimosus infuscatus*, *Phrygilus unicolor*, *Phyllomyias uropygialis*, *Piaya cayana*, *Picoides fumigatus*, *Pipraeidea melanonota*, *Pipreola riefferii*, *Piranga olivacea*, *Piranga rubra*, *Pitangus sulphuratus*, *Pluvialis dominica*, *Podilymbus podiceps*, *Porphyrio martinicus*, *Porzana carolina*, *Progne tapera*, *Protonotaria citrea*, *Pseudocolopteryx acutipennis*, *Pseudoscops clamator*, *Pterophanes cyanopterus*, *Pygochelidon cyanoleuca*, *Pyrocephalus rubinus*, *Pyrrhomyias cinnamomeus*, *Quiscalus lugubris*, *Rallus semiplumbeus*, *Ramphocelus dimidiatus*, *Ramphomicron microrhynchum*, *Riparia riparia*, *Rupornis magnirostris*, *Sayornis nigricans*, *Serpophaga cinerea*, *Setophaga castanea*, *Setophaga cerulea*, *Setophaga petechia*, *Setophaga ruticilla*, *Setophaga striata*, *Sicalis flaveola*, *Sicalis luteola*, *Spiza americana*, *Sporagra*

spinescens, *Sturnella magna*, *Synallaxis subpudica*, *Systellura longirostris*, *Tangara vitriolina*, *Thraupis episcopus*, *Thraupis palmarum*, *Tringa flavipes*, *Tringa melanoleuca*, *Tringa solitaria*, *Troglodytes aedon*, *Troglodytes solstitialis*, *Turdus fuscater*, *Turdus ignobilis*, *Tyrannus melancholicus*, *Tyrannus savana*, *Tyrannus tyrannus*, *Tyto alba*, *Vanellus chilensis*, *Vireo olivaceus*, *Vultur gryphus*, *Zenaida auriculata*, *Zonotrichia capensis*

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

pages: 836

issue:

year: 1986

0: Hilty, Steven

1: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

tags: *Accipiter striatus*, *Actitis macularius*, *Adelomyia melanogenys*, *Agelaiocercus kingii*, *Amblycercus holosericeus*, *Ampelion rubrocristatus*, *Anas andium*, *Anas clypeata*, *Anas discors*, *Andigena nigrirostris*, *Anisognathus igniventris*, *Ardea alba*, *Ardea herodias*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Asthenes flammulata*, *Asthenes fuliginosa*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Butorides virescens*, *Cairina moschata*, *Calidris bairdii*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Cathartes aura*, *Catharus minimus*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chaetura pelagica*, *Chalcostigma heteropogon*, *Chordeiles minor*, *Chrysomus icterocephalus*, *Cinclodes albidiventris*, *Cinclus leucocephalus*, *Cinnycerthia unirufa*, *Cistothorus platensis*, *Coccyzus pumila*, *Coccyzus americanus*, *Coccyzus melacoryphus*, *Coeligena bonapartei*, *Colibri coruscans*, *Colibri thalassinus*, *Columba livia*, *Columbina talpacoti*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cinereus*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Dendrocygna autumnalis*, *Dendrocygna viduata*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dolichonyx oryzivorus*, *Dubusia taeniata*, *Egretta thula*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreoventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Geothlypis philadelphia*, *Geranoaetus melanoleucus*, *Glaucidium jardinii*, *Grallaria quitesis*, *Grallaria ruficapilla*, *Grallaria squamigera*, *Haplospiza rustica*, *Heliangelus amethysticollis*, *Heliangelus exortis*, *Hellmayrea gularis*, *Hemispingus supercilialis*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus icterus*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Lafresnaya lafresnayi*, *Leiothlypis peregrina*, *Leptasthenura andicola*, *Lesbia nuna*, *Lesbia victoriae*, *Machetornis rixosa*, *Margarornis squamiger*, *Mecocerculus leucophrys*, *Megascops choliba*, *Melanerpes rubricapillus*, *Merganetta armata*, *Metallura tyrianthina*, *Milvago chimachima*, *Mimus gilvus*, *Mniotilta varia*, *Molothrus bonariensis*, *Muscisaxicola alpinus*, *Myiarchus crinitus*, *Myiodynastes luteiventris*, *Myiotheretes fumigatus*, *Myiotheretes striaticollis*, *Myiothlypis nigrocristata*, *Nomonyx dominicus*, *Nycticorax nycticorax*, *Ochthoeca cinnamomeiventris*, *Ochthoeca diadema*, *Ochthoeca fumicolor*, *Ochthoeca rufipectoralis*, *Orochelidon murina*, *Oxyopogon guerinii*, *Oxyura jamaicensis*, *Pachyramphus versicolor*, *Pandion haliaetus*, *Pardirallus maculatus*, *Parkesia noveboracensis*, *Patagioenas fasciata*, *Petrochelidon pyrrhonota*, *Phaetusa simplex*, *Phalacrocorax brasilianus*, *Pheucticus aureoventris*, *Pheucticus ludovicianus*, *Phimosus infuscatus*, *Phrygilus unicolor*, *Phyllomyias uropygialis*, *Piaya cayana*, *Picoides fumigatus*, *Pipraeidea melanonota*, *Pipreola riefferii*, *Piranga olivacea*, *Piranga rubra*, *Pitangus sulphuratus*, *Pluvialis dominica*, *Podilymbus podiceps*, *Porphyrio martinicus*, *Porzana carolina*, *Progne tapera*, *Protonotaria citrea*, *Pseudocolopteryx acutipennis*, *Pseudoscops clamator*, *Pterophanes cyanopterus*, *Pygochelidon cyanoleuca*, *Pyrocephalus rubinus*, *Pyrrhomyias cinnamomeus*, *Quiscalus lugubris*, *Rallus semiplumbeus*, *Ramphocelus dimidiatus*, *Ramphomicron microrhynchum*, *Riparia riparia*, *Rupornis magnirostris*, *Sayornis nigricans*, *Serpophaga cinerea*, *Setophaga castanea*, *Setophaga cerulea*, *Setophaga petechia*, *Setophaga ruticilla*, *Setophaga striata*, *Sicalis flaveola*, *Sicalis luteola*, *Spiza americana*, *Sporagra spinescens*, *Sturnella magna*, *Synallaxis subpudica*, *Systellura longirostris*, *Tangara vitriolina*, *Thraupis episcopus*, *Thraupis palmarum*, *Tringa flavipes*, *Tringa melanoleuca*, *Tringa solitaria*, *Troglodytes aedon*, *Troglodytes*

solstitialis,Turdus fuscater,Turdus ignobilis,Tyrannus melancholicus,Tyrannus savana,Tyrannus tyrannus,Tyto alba, Vanellus chilensis,Vireo olivaceus,Vultur gryphus,Zenaida auriculata,Zonotrichia capensis

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

behaviorUnstructured: Es un ave notablemente gregaria aun durante la reproducción (Hilty & Brown, 1986); la Monjita se encuentra generalmente en grupos descansando en los juncos (Asociación Bogotana de Ornitología, 2000); fuera de la temporada reproductiva se observa en grupos aún más grandes (Fraga, 2011).

Aunque el canto de la Monjita es elaborado, no es un canto musical, más bien es fuerte y raspante (Hilty & Brown, 1986); en los machos pueden encontrarse dos tipos de cantos, uno que consiste en breves notas o silbidos seguidos por áspero zumbido, el otro una secuencia más variable y agradable de gorjeos o silbidos (Fraga, 2011), trinos secos, altos, a veces, zumbantes o chirriantes, prolongados, mezclado con silbidos y notas cortas (Asociación Bogotana de Ornitología, 2000), los dos tipos pueden ser alternados. Frecuentemente se observan grupos cantando en coros (Fraga, 2011). Generalmente los machos vocalizan desde las puntas de los juncos. El reclamo más común es un churt o chirt metálico, también produce un chasquido seco (Asociación Bogotana de Ornitología, 2000), los llamados son un chik o un silbido descendente (Fraga, 2011).

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

endemicAtomized

distribution

ancillaryData

dataObject:
additionalInformation:

reference

keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus

editors

editors: Stiles, F G
editors: Bohórquez, C I
editors: Cadena, C D
editors: De la Zerda, S
editors: Hernández, M
editors: Roselli, L
editors: Kelsey, M
editors: Valencia, I D
editors: Knapp, D
institution:
edition: Primera Ed
address:
publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca
accessed:
websites:
chapter:
series:
pages: 276
issue:
year: 2000

authors

authors: Asociación Bogotana de Ornitología
title: Aves de la Sabana de Bogotá: guía de campo
source:
type: book
tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina

canadensis,Catamenia analis,Catamenia inornata,Catharus fuscescens,Catharus ustulatus,Chaetocercus
mulsant,Chrysomus icterocephalus

abstract:

identifiers

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

agent

contentType:

dataType:

identifier:

distributionUnstructured: La Monjita se distribuye en el norte de Sudamérica, desde Colombia, Trinidad, hasta el noreste de Perú y norte de la Amazonia Brasileña (Asociación Bogotana de Ornitología, 2000). En Colombia se encuentra en el valle bajo del Atrato (donde es poco común) y hacia el este hasta la región de Santa Marta, luego hacia el sur, en el valle medio del Cauca y alto Magdalena hasta el sur del Tolima, en el este hasta Meta y Vichada y luego a lo largo del Amazonas hasta Brasil (Hilty & Brown, 1986). Actualmente se reconocen dos subespecies con diferencias en su distribución; C. i. icterocephalus (Linnaeus, 1766) se encuentra en Colombia en los departamentos de Tolima, Meta y Vichada, en Venezuela principalmente en el norte del río Orinoco, en Trinidad, Guyanas, norte de Brasil y nororiente de Perú. C. i. bogotensis (Chapman, 1914) solo se encuentra al nororiente de Colombia en los territorios cercanos a Bogotá (Fraga, 2011). En Lima (Perú) se encuentra una población introducida persistente (The Cornell Lab of Ornithology, 2010).

Generalmente la Monjita se encuentra por debajo de los 1000 msnm a excepción de la subespecie C. i. bogotensis que está cerca de los 2600 msnm en la sabana de Bogotá (Asociación Bogotana de Ornitología, 2000).

distributionAtomized

stateProvince: Córdoba

ancillaryData

locality:

municipality:

country: Colombia

temporalCoverage

endDate:

startDate:

distributionScope

type:

threatStatus

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

0

dataObject:

additionalInformation:

0

link: <http://www.birdlife.org/datazone/species/factsheet/22724200>

keywords: Chrysomus icterocephalus

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2014

0: BirdLife International

title: Species factsheet: Chrysomus icterocephalus

source:

type: miscellany

tags: Chrysomus icterocephalus

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

populationBiologyUnstructured: El tamaño global de la población no ha sido cuantificado, sin embargo se considera una especie bastante común con una tendencia estable sin evidencia o declinaciones o cambios considerables (BirdLife International, 2014).

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: La Monjita es localmente común en los ecosistemas de agua dulce, pastizales húmedos y en las islas de los ríos, a lo largo del río Amazonas y algunos afluentes, en los llanos, y en las regiones y los valles de tierras bajas de Colombia hasta el norte de Brasil. En Lima (Perú) se encuentra una población introducida persistente. Los machos son amarillo brillante en la cabeza y en el pecho, lo que contrasta elegantemente con su plumaje negro en el resto del cuerpo. Las hembras tienen esta coloración más apagada, no tan amarillo sino más café-oliva. Al igual que en otros icteridos, los machos pueden aparearse con varias hembras, sin embargo no es muy común en esta especie dado que el nido utilizado por cada hembra debe ser construido por el macho (The Cornell Lab of Ornithology, 2010). Su alimentación se basa principalmente en insectos y semillas (Asociación Bogotana de Ornitología, 2000); se le observa generalmente en grandes grupos, reposando en los juncales o buscando en ellos su alimento (Asociación Bogotana de Ornitología, 2000; Hilty & Brown, 1986).

[associatedParty](#)

role:

personnelIdentifier:
personnelDirectory:
homepage:
email: enataliagomezm@gmail.com
phone:
postalCode:
country: Colombia
state: Bogotá Distrito Capital
city: Bogotá D.C.
address:
organisation:
position:
lastName: Gómez Melendro
firstName: Edith Natalia

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:
mediaURL: <http://www.xeno-canto.org/36652>
description:

subject

source: <http://www.xeno-canto.org>

audience

bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:

agent

mimeType:
dataType:
identifier:

_id

_bsontype: ObjectID
id: VÃ~çð lg#ç5