

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56e33bc283c45700544e3db8

Fecha de creación: 2016-03-11T21:42:32.528Z

creation_date: 2016-03-11T21:42:32.528Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

fullDescriptionUnstructured: Dos venas antenodales. Origen de la vena M3 más cerca del nodo que del arculo. La puna del cuadrángulo toca el margen posterior del ala. Alas muy angostas con peciolación extensa. Abdomen muy largo y delgado, con varias bandas negras que alternan con anillos café claro con visos verdosos. Machos con cercos amarillo intenso, curvos hacia adentro y provistos de una espina cerca del ápice.

briefDescription:

[references](#)

keywords: Argia cupraurea,Argia pulla Hagen in,Cora chiribiquete,Cora inca,Erythemis vesiculosa,Erythrodiplax kimminsi,Erythrodiplax umbrata,Erythrodiplax unimaculata,Mecistogaster linearis,Orthemis discolor,Perissolestes magdalenae,Perithemis lais,Perithemis mooma,Polythore beata,Polythore concinna,Polythore derivata,Polythore mutata,Telebasis corallina,Uracis fastigiata,Uracis imbuta

[editors](#)

institution:

address:

publisher: Instituto Nacional de Biodiversidad, Ministerio del Ambiente y Energía, Universidad Nacional de Costa Rica, Banco Mundial, GEF

accessed:

websites:

chapter:

series:

issue:

year: 2006

authors

authors: Esquivel, C
title: Libélulas de Mesoamérica y el Caribe. Dragonflies and damselflies of Middle America and the Caribbean
source:
type: book
tags: Argia cupraurea,Argia pulla Hagen in,Cora chiribiquete,Cora inca,Erythemis vesiculosa,Erythrodiplax kimminsi,Erythrodiplax umbrata,Erythrodiplax unimaculata,Mecistogaster linearis,Orthemis discolor,Perissolestes magdalenae,Perithemis lais,Perithemis mooma,Polythore beata,Polythore concinna,Polythore derivata,Polythore mutata,Telebasis corallina,Uracis fastigiata,Uracis imbuta
abstract:

identifiers

last_modified:
created:
group_id:
profile_id:
id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:
providerLink:

publicationStatus

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

replacementNameFor

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

sanctioned

microReference:
source:

datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

conservedAgainst

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:

identifier:

typificacion

typeOfType:

[lectotypeMicroReferenceVoucher](#)

[lectotypePublicationVoucher](#)

[voucherReference](#)

simple:

microReference:

year:

publishedIn

simple:

source:

datatype:

identifier:

specialAuthorship

[authors](#)

[year](#)

simple:

canonicalAuthorship

[authors](#)

[year](#)

simple:

simple: Williamson & Williamson, 1924

canonicalName

infraspecificEpithet:

specificEpithet:

infragenericEpithet:

linkType:

ref:

uninomial:

simple: Perissolestes magdalenae

rank: SPECIES

simple: Perissolestes magdalenae Williamson & Williamson, 1924

attributes

nomenclaturalCode:

isAnamorphic:

id:

[synonymsAtomized](#)

hierarchy

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source: The Catalogue of Life, 3rd January 2011

audience

bibliographicCitation:
rightsHolder:
rights:
license:
title:
modified: 2016-03-11T21:42:12.524Z
created: 2016-03-11T21:42:12.524Z

agent

homepage:
role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:
parentTaxon: Perissolestes
higherClassification:
infraspecificEpithet:
specificEpithet:
taxonRank: SPECIES
subGenus:
genus: Perissolestes
family: Perilestidae
order: Odonata
classHierarchy: Insecta
phylum: Arthropoda
kingdom: Animalia
recommended:

classification:

commonNameAtomized

ancillaryData

dataObject:

additionalInformation:

reference

geoPoint:

location:

thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

agent

contentType:

dataType:

identifier:

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

ancillaryData

type:

language: Español

name: Caballito del diablo

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:

additionalInformation:

geoPoint:

location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:

reproductionUnstructured: Pueden llegar a copular dos veces seguidas con la misma hembra. Se han observado machos aterrizar en la punta del abdomen de hembras posadas, y luego subir caminando por el abdomen hasta alcanzar el tórax de la hembra para sujetarla con los cercos. Una vez en tándem, la pareja vuela a otra percha y copulan.

0

dataObject:
additionalInformation:

[reference](#)

geoPoint:
location:
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

[audience](#)

rightsHolder:
rights:
license:
modified:
created:

[agent](#)

mimeType:
dataType:
identifier:
relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedBy: TraitBank
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue: Sexual
measurementType: Tipo de Reproducción
measurementID:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

feedingUnstructured:

0

0

strategy: Depredadores

type:

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

behaviorUnstructured: Se conoce poco de ellos. Su modo de posarse es único (junto con las otras especies del género); se cuelgan de ramitas u hojas, con las alas entreabiertas y el abdomen un poco curvado hacia arriba. Generalmente se posan 1.5 o aún más alto. Su vuelo es lento y delicado, y vuelan muy poco, la mayor parte del tiempo se la pasan posados, y brevemente vuelan para cambiar de percha. Los machos no muestran territorialidad, varios de ellos pueden estar cercanos en la misma rama y no pelean, excepto por ocasionales enfrentamientos en vuelo suspendido, sin tocarse, luego del cual cada uno busca su percha.

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

annualCycle

annualCycleUnstructured:

habitat

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

habitatUnstructured: Habita bosques húmedos tropicales, arroyos y pantanos en selvas húmedas.

[endemicAtomized](#)

[distribution](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

agent

contentType:

dataFormat:

identifier:

distributionUnstructured: Esta especie de libélula se ha encontrado en zonas con altitudes que alcanzan los 400 msnm. En el país se ha reportado para la región de la Amazonia.

distributionAtomized

ancillaryData

locality:

municipality:

country: México

temporalCoverage

endDate:

startDate:

distributionScope

type:

threatStatus

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

abstract: Perissolestes magdalenae es una libélula depredadora de gran tamaño, su cuerpo es muy alargado y delgado y sus alas son angostas y cortas. Se posan con las alas semiabiertas y el abdomen un poco curvado hacia arriba.

associatedParty

role:
personnelIdentifier:
personnelDirectory:
homepage:
email: enataliagomez@gmail.com
phone:
postalCode:
country: Colombia
state: Bogotá Distrito Capital
city: Bogotá D.C.
address:
organisation:
position:
lastName: Gómez Melendro
firstName: Edith Natalia

ancillaryData

dataObject:
additionalInformation: Créditos específicos: Al Instituto Amazónico de Investigaciones Científicas Sinchi. Al Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIAT-AC. La elaboración de esta ficha fue posible gracias al apoyo del gobierno de Finlandia a través del convenio entre la Secretaria General de la CAN y el Instituto Sinchi para el proyecto "Sistema Regional Amazónico de Información Ambiental con énfasis en Biodiversidad de los países miembros de la CAN, (programa BIOCAN), componente SIAT-AC".

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder:
rights:
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

mimeType:
dataType:
identifier:

_id
_bsontype: ObjectID
id: Vã;ÄfÄW DãÜ€