

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56d4bd403c16479905cba8b9

Fecha de creación: 2016-02-29T21:50:58.925Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Clara Maria Pérez Gallego

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

habitatUnstructured: Terrestre, en áreas intervenidas abiertas. Común en pueblos, dehesas y otras áreas abiertas y secas con matorrales dispersos; también matorral árido y cactus.

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

[ancillaryData](#)

type:

language: Español

name: Centzontle tropical

creation_date: 2016-02-29T21:50:58.925Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Clara Maria Pérez Gallego
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
fullDescriptionUnstructured: Tiene un tamaño de 25 cm. Delgado y de cola larga. Encima café grisáceo pálido con estrecha mascara negruzca y superciliar prominente blanco yesoso; alas negruzcas; cobertoras marginadas de blanco (sin parches blancos en el ala); cola negruzca con amplios ápices blancos; por debajo blanco opaco. Inmaduro es similar pero más café por encima, más anteado debajo.
briefDescription:

references

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:
providerLink:

publicationStatus

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

replacementNameFor

microReference:
source:
datatype:
identifier:
source:
datatype:

identifier:
note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

conservedAgainst

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:

datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion
typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:
canonicalAuthorship

authors

year

simple:
simple: (Vieillot, 1808)

canonicalName

infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:
ref:
uninomial:
simple: Mimus gilvus
rank: SPECIES
simple: Mimus gilvus (Vieillot, 1808)

attributes

nomenclaturalCode:

isAnamorphic:

id:

[synonymsAtomized](#)

[hierarchy](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:

rightsHolder:

rights:

license:

title:

modified: 2016-02-29T21:50:38.925Z

created: 2016-02-29T21:50:38.925Z

[agent](#)

homepage:

role:

email:

phone:

address:

position:

organisation:

lastName:

firstName:

contentType:

dataType:

identifier:

parentTaxon: Mimus

higherClassification:

infraspecificEpithet:

specificEpithet:

taxonRank: SPECIES

subGenus:

genus: Mimus
family: Mimidae
order: Passeriformes
classHierarchy: Aves
phylum: Chordata
kingdom: Animalia
recommended:
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Clara Maria Pérez Gallego
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:

reproductionUnstructured: El nido es una taza de ramitas a baja altura en arbustos, por lo general ponen 3 huevos de color azul verdoso pálido con manchas orín en extremo mayor.

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:
additionalInformation:

geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Clara Maria Pérez Gallego
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
feedingUnstructured: Principalmente insectívoros pero ocasionalmente comen fruta.

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Clara Maria Pérez Gallego

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

behaviorUnstructured: Diurna y en parejas y buscando el alimento sobre el suelo. Ave conspicua de muchas áreas abiertas o urbanas, notablemente agresivo alrededor de su nido. A menudo come en el suelo y frecuentemente levanta las alas. La voz es una serie larga de notas y frases, algunas repetidas varias veces, otras no; no imita a otras aves.

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

[endemicAtomized](#)

[distribution](#)

[ancillaryData](#)

dataObject:
additionalInformation:

[reference](#)

geoPoint:
location:
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:
rightsHolder:
rights: Clara Maria Pérez Gallego
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:

created:

agent

contentType:

dataType:

identifier:

distributionUnstructured: S México hasta Honduras; introducido en Panamá; Colombia hasta Guayanas, N y SE Brasil; mayoría de islas Caribe.

En el país al W de los Andes en regiones abiertas y al E de los Andes S hasta Meta y E Vichada, en las regiones Andina y Orinoquía hasta 2600 msnm (Sabana de Bogotá).

distributionAtomized

stateProvince: Tolima

ancillaryData

locality:

municipality:

country: Colombia

temporalCoverage

endDate:

startDate:

distributionScope

type:

threatStatus

ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:

locality:

municipality:

county:

stateProvince:

country: Colombia

0: IUCN

measurementValue: En peligro (EN)

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementType: IUCN

measurementID:

territory

territoryUnstructured:
areaOfOccupancy:
extentOfOccurrence:

populationBiology
populationBiologyUnstructured:

legislation
legislationUnstructured:

directThreats
directThreatsUnstructured:

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:
abstract:

associatedParty

role:
personnelIdentifier:
personnelDirectory:
homepage:
email: enataliagomezm@gmail.com
phone:
postalCode:
country: Colombia
state: Bogotá Distrito Capital
city: Bogotá D.C.

address:
organisation:
position:
lastName: Gómez Melendro
firstName: Edith Natalia

ancillaryData

dataObject:
additionalInformation: Registros biológicos: En la Central Jaguas se registró en los rastrojos de Puente Cable y Vivero y en las áreas intervenidas de Embalse, Vertedero, Compuertas y Zona Baja. En la Central San Carlos se registró en los rastrojos de Zona Baja, Portería Presa y Central San Carlos y en las áreas intervenidas de Vertedero, Casino Viejo y en la vía hacia el Domo.
Imagen: Mimus_gilvus.jpg

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder: Clara Maria Pérez Gallego
rights:
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

mimeType:
dataType:
identifier:

_id
_bsontype: ObjectID
id: VÔ½@< G™\º,•