

Reporte de Fichas para el usuario: Edgar Herrera

Ficha ID: 56e7978083c45700544e41d2

Fecha de creación: 2016-03-15T05:03:26.469Z

creation_date: 2016-03-15T05:03:26.469Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

0

keywords: Accipiter collaris, Accipiter cooperii, Accipiter poliogaster, Accipiter superciliosus, Amazona ochrocephala, Andigena hypoglaucha, Aramus guarauna, Aratinga pertinax, Ardea alba, Ardea cocoi, Ardea herodias, Arremonops conirostris (Bonaparte 1850), Arremonops tocuyensis (Todd 1912), Atalotriccus pilaris, Attila spadiceus (Gmelin 1789), Basileuterus flaveolus (S. F. Baird 1865), Bubo virginianus (Gmelin 1788), Busarellus nigricollis, Buteogallus anthracinus, Buteogallus meridionalis, Buteogallus schistaceus, Buteogallus solitarius, Buteogallus urubitinga, Cacicus cela, Cairina moschata, Campylorhamphus trochilirostris (Lichtenstein 1820), Campylorhynchus griseus (Swainson 1838), Campylorhynchus nuchalis (Cabanis 1847), Caprimulgus cayennensis (Gmelin 1789), Cardinalis phoeniceus Bonaparte 1838, Carduelis psaltria (Say 1823), Cathartes aura, Cathartes burrovianus, Cathartes melambrotus, Certhiaxis cinnamomeus (Gmelin 1788), Ceryle torquata, Chlorostilbon gibsoni (Fraser 1840), Chondrohierax uncinatus, Chordeiles acutipennis (Hermann 1783), Chrysomus icterocephalus, Circus buffoni, Circus cinereus, Circus cyaneus, Coccyzus pumila, Columbina minuta (Linnaeus 1766), Conirostrum leucogenys (Lafresnaye 1852), Coragyps atratus, Coryphospingus pileatus (Wied-Neuwied 1821), Crax alector, Crotophaga major, Crotophaga sulcirostris Swainson 1827, Cryptoleucopteryx plumbea, Cyanocorax affinis (Pelzelin 1856), Cyclarhis gujanensis (J. F. Gmelin 1789), Dryocopus lineatus (Linnaeus 1766), Elaenia chiriquensis Lawrence 1865, Elanoides forficatus, Elanus leucurus, Falco deiroleucus, Fluvicola pica (Boddaert 1783), Forpus passerinus (Linnaeus 1758), Forpus xanthopterygius (Spix 1824), Furnarius leucopus Swainson 1838, Galbula ruficauda Cuvier 1816, Gallinula chloropus (Linnaeus 1758), Gampsonyx swainsonii, Geranospiza caerulescens, Glaucidium brasilianum (Gmelin 1788), Harpagus bidentatus, Harpia harpyja, Helicolestes hamatus, Hemitriccus margaritaceiventer (Orbigny & Lafresna, Himantopus mexicanus (Statius Muller 1776), Hylophilus flavipes Lafresnaye 1845, Hypnelus ruficollis (Wagler 1829), Icterus auricapillus Cassin 1848, Icterus galbula (Linnaeus 1758), Icterus nigrogularis (Hahn 1819), Ictinia mississippiensis, Ictinia plumbea, Inezia subflava (P. L. Sclater & Salvin 1873), Inezia tenuirostris (Cory 1913), Jabiru mycteria, Jacana jacana, Lepidopyga goudoti (Bourcier 1843), Leptotila verreauxi (Bonaparte 1855), Leucopternis melanops, Leucopternis semiplumbea, Machetornis rixosa, Megaceryle alcyon (Linnaeus 1758), Megaceryle torquata (Linnaeus 1766), Megarynchus pitangua (Linnaeus 1766), Megascops choliba, Melanerpes rubricapillus, Micrastur plumbeus, Mitu tomentosa, Molothrus bonariensis, Morphnarchus princeps, Morphnus guianensis, Myiarchus crinitus, Myiarchus venezuelensis Lawrence 1865, Myiopagis gaimardii (Orbigny 1840), Myiozetetes similis (Spix 1825), Myospiza humeralis (Bosc 1792), Nemosia pileata (Boddaert 1783), Neochen jubata, Opisthocomus hoazin, Oroaetus isidori, Ortalis ruficauda Jardine 1847, Pandion haliaetus, Parabuteo unicinctus, Patagioenas corensis (Jacquin 1784), Phaeomyias murina (Spix 1825), Pheucticus ludovicianus, Piculus chrysochloros (Vieillot 1818), Picumnus cinnamomeus Wagler 1829, Pilherodius pileatus (Boddaert 1783), Pionus sordidus (Linnaeus 1758), Pitangus lictor (Lichtenstein 1823), Pitangus sulphuratus, Polioptila plumbea (J. F. Gmelin 1788), Pseudastur albicollis, Pulsatrix perspicillata (Latham 1790), Pyrocephalus rubinus, Quiscalus lugubris, Quiscalus mexicanus (Gmelin 1788), Rostrhamus sociabilis, Sakesphorus canadensis (Linnaeus 1766), Saltator coerulescens Vieillot

1817, *Sarcoramphus papa*, *Sarkidiornis melanotos* (Pennant 1769), *Spiza americana*, *Spizaetus tyrannus*, *Sporophila bouvronides* (Lesson 1831), *Sporophila intermedia* Cabanis 1851, *Sporophila nigricollis* (Vieillot 1823), *Stelgidopteryx ruficollis* (Vieillot 1817), *Sterna superciliaris*, *Sturnella magna*, *Sublegatus arenarum* (Salvin 1863), *Synallaxis candei* Orbigny & Lafresnaye 1838, *Tachycineta bicolor* (Vieillot 1808), *Tapera naevia* (Linnaeus 1766), *Thamnophilus doliatus* (Linnaeus 1764), *Thraupis glaucocolpa* Cabanis 1850, *Thraupis palmarum*, *Thryothorus leucotis* (Lafresnaye 1845), *Tyrannus dominicensis* (Gmelin 1788), *Tyto alba*, *Vireo flavoviridis* (Cassin 1851), *Zenaida auriculata*

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

issue:

year: 1986

0: Hilty, S

1: Brown, W. L.

title: A Guide to the Birds of Colombia

source:

type: book

tags: *Accipiter collaris*, *Accipiter cooperii*, *Accipiter poliogaster*, *Accipiter superciliosus*, *Amazona ochrocephala*, *Andigena hypoglaucha*, *Aramus guarauna*, *Aratinga pertinax*, *Ardea alba*, *Ardea cocoi*, *Ardea herodias*, *Arremonops conirostris* (Bonaparte 1850), *Arremonops tocuyensis* (Todd 1912), *Atalotriccus pilaris*, *Attila spadiceus* (Gmelin 1789), *Basileuterus flaveolus* (S. F. Baird 1865), *Bubo virginianus* (Gmelin 1788), *Busarellus nigricollis*, *Buteogallus anthracinus*, *Buteogallus meridionalis*, *Buteogallus schistaceus*, *Buteogallus solitarius*, *Buteogallus urubitinga*, *Cacicus cela*, *Cairina moschata*, *Campylorhamphus trochilirostris* (Lichtenstein 1820), *Campylorhynchus griseus* (Swainson 1838), *Campylorhynchus nuchalis* (Cabanis 1847), *Caprimulgus cayennensis* (Gmelin 1789), *Cardinalis phoeniceus* Bonaparte 1838, *Carduelis psaltria* (Say 1823), *Cathartes aura*, *Cathartes burrovianus*, *Cathartes melambrotus*, *Certhiaxis cinnamomeus* (Gmelin 1788), *Ceryle torquata*, *Chlorostilbon gibsoni* (Fraser 1840), *Chondrohierax uncinatus*, *Chordeiles acutipennis* (Hermann 1783), *Chrysomus icterocephalus*, *Circus buffoni*, *Circus cinereus*, *Circus cyaneus*, *Coccyzus pumila*, *Columbina minuta* (Linnaeus 1766), *Conirostrum leucogenys* (Lafresnaye 1852), *Coragyps atratus*, *Coryphospingus pileatus* (Wied-Neuwied 1821), *Crax alector*, *Crotophaga major*, *Crotophaga sulcirostris* Swainson 1827, *Cryptoleucopteryx plumbea*, *Cyanocorax affinis* (Pelzeln 1856), *Cyclarhis gujanensis* (J. F. Gmelin 1789), *Dryocopus lineatus* (Linnaeus 1766), *Elaenia chiriquensis* Lawrence 1865, *Elanoides forficatus*, *Elanus leucurus*, *Falco deiroleucus*, *Fluvicola pica* (Boddaert 1783), *Forpus passerinus* (Linnaeus 1758), *Forpus xanthopterygius* (Spix 1824), *Furnarius leucopus* Swainson 1838, *Galbula ruficauda* Cuvier 1816, *Gallinula chloropus* (Linnaeus 1758), *Gampsonyx swainsonii*, *Geranospiza caerulescens*, *Glaucidium brasilianum* (Gmelin 1788), *Harpagus bidentatus*, *Harpia harpyja*, *Helicolestes hamatus*, *Hemitriccus margaritaceiventer* (Orbigny & Lafresna), *Himantopus mexicanus* (Statius Muller 1776), *Hylophilus flavipes* Lafresnaye 1845, *Hypnelus ruficollis* (Wagler 1829), *Icterus auricapillus* Cassin 1848, *Icterus galbula* (Linnaeus 1758), *Icterus nigrogularis* (Hahn 1819), *Ictinia mississippiensis*, *Ictinia plumbea*, *Inezia subflava* (P. L. Sclater & Salvin 1873), *Inezia tenuirostris* (Cory 1913), *Jabiru mycteria*, *Jacana jacana*, *Lepidopyga goudoti* (Bourcier 1843), *Leptotila verreauxi* (Bonaparte 1855), *Leucopternis melanops*, *Leucopternis semiplumbea*, *Machetornis rixosa*, *Megaceryle alcyon* (Linnaeus 1758), *Megaceryle torquata* (Linnaeus 1766), *Megarynchus pitangua* (Linnaeus 1766), *Megascops choliba*, *Melanerpes rubricapillus*, *Micrastur plumbeus*, *Mitu tomentosa*, *Molothrus bonariensis*, *Morphnarchus princeps*, *Morphnus guianensis*, *Myiarchus crinitus*, *Myiarchus venezuelensis* Lawrence 1865, *Myiopagis gaimardii* (Orbigny 1840), *Myiozetetes similis* (Spix 1825), *Myospiza humeralis* (Bosc 1792), *Nemosia pileata* (Boddaert 1783), *Neochen jubata*, *Opisthocomus hoazin*, *Oroaetus isidori*, *Ortalis ruficauda* Jardine 1847, *Pandion haliaetus*, *Parabuteo unicinctus*, *Patagioenas corensis* (Jacquin 1784), *Phaeomyias murina* (Spix 1825), *Pheucticus ludovicianus*, *Piculus chrysochloros* (Vieillot 1818), *Picumnus cinnamomeus* Wagler 1829, *Pilherodius pileatus* (Boddaert 1783), *Pionus sordidus* (Linnaeus 1758), *Pitangus lictor* (Lichtenstein 1823), *Pitangus sulphuratus*, *Polioptila plumbea* (J. F. Gmelin 1788), *Pseudastur albicollis*, *Pulsatrix perspicillata* (Latham 1790), *Pyrocephalus rubinus*, *Quiscalus lugubris*, *Quiscalus mexicanus* (Gmelin 1788), *Rostrhamus sociabilis*, *Sakesphorus canadensis* (Linnaeus 1766), *Saltator coerulescens* Vieillot 1817, *Sarcoramphus papa*, *Sarkidiornis melanotos* (Pennant 1769), *Spiza americana*, *Spizaetus tyrannus*, *Sporophila bouvronides* (Lesson 1831), *Sporophila intermedia* Cabanis 1851, *Sporophila nigricollis* (Vieillot 1823), *Stelgidopteryx*

ruficollis (Vieillot 1817), *Sterna superciliaris*, *Sturnella magna*, *Sublegatus arenarum* (Salvin 1863), *Synallaxis candei* Orbigny & Lafresnaye 1838, *Tachycineta bicolor* (Vieillot 1808), *Tapera naevia* (Linnaeus 1766), *Thamnophilus doliatus* (Linnaeus 1764), *Thraupis glaucocolpa* Cabanis 1850, *Thraupis palmarum*, *Thryothorus leucotis* (Lafresnaye 1845), *Tyrannus dominicensis* (Gmelin 1788), *Tyto alba*, *Vireo flavoviridis* (Cassin 1851), *Zenaida auriculata*

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

fullDescriptionUnstructured: 66 cm.; 1.1 kg. Con aspecto de garza, pero el cuello es relativamente más corto y grueso; el pico es largo comprimido lateralmente, ligeramente curvo. Los adultos tienen la cara café grisácea, la cabeza y el cuello están finamente listados con café y blanco; el cuerpo y las coberteras de las alas son café oscuro con manchas lanceoladas blancas; la remeras son de color café oscuro con un lustre entre verdoso y púrpura. El pico es de color cuerno-amarillento desvaneciéndose a gris oscuro en la punta; las patas son verdoso opaco.

JUVENILES: parecidos, pero con menos manchas blancas; la textura del plumaje es más laxa (Hilty & Brown 1986).

briefDescription:

references

keywords: *Accipiter collaris*, *Accipiter cooperii*, *Accipiter poliogaster*, *Accipiter superciliosus*, *Amazona ochrocephala*, *Andigena hypoglaucha*, *Aramus guarauna*, *Aratinga pertinax*, *Ardea alba*, *Ardea cocoi*, *Ardea herodias*, *Arremonops conirostris* (Bonaparte 1850), *Arremonops tocuyensis* (Todd 1912), *Atalotriccus pilaris*, *Attila spadiceus* (Gmelin 1789), *Basileuterus flaveolus* (S. F. Baird 1865), *Bubo virginianus* (Gmelin 1788), *Busarellus nigricollis*, *Buteogallus anthracinus*, *Buteogallus meridionalis*, *Buteogallus schistaceus*, *Buteogallus solitarius*, *Buteogallus urubitinga*, *Cacicus cela*, *Cairina moschata*, *Campylorhamphus trochilostrius* (Lichtenstein 1820), *Campylorhynchus griseus* (Swainson 1838), *Campylorhynchus nuchalis* (Cabanis 1847), *Caprimulgus cayennensis* (Gmelin 1789), *Cardinalis phoeniceus* Bonaparte 1838, *Carduelis psaltria* (Say 1823), *Cathartes aura*, *Cathartes burrovianus*, *Cathartes melambrotus*, *Certhiaxis cinnamomeus* (Gmelin 1788), *Ceryle torquata*, *Chlorostilbon gibsoni* (Fraser 1840), *Chondrohierax uncinatus*, *Chordeiles acutipennis* (Hermann 1783), *Chrysomus icterocephalus*, *Circus buffoni*, *Circus cinereus*, *Circus cyaneus*, *Coccyzus pumila*, *Columbina minuta* (Linnaeus 1766), *Conirostrum leucogenys* (Lafresnaye 1852), *Coragyps atratus*, *Coryphospingus pileatus* (Wied-Neuwied 1821), *Crax alector*, *Crotophaga major*, *Crotophaga sulcirostris* Swainson 1827, *Cryptoleucopteryx plumbea*, *Cyanocorax affinis* (Pelzelin 1856), *Cyclarhis gujanensis* (J. F. Gmelin 1789), *Dryocopus lineatus* (Linnaeus 1766), *Elaenia chiriquensis* Lawrence 1865, *Elanoides forficatus*, *Elanus leucurus*, *Falco deiroleucus*, *Fluvicola pica* (Boddaert 1783), *Forpus passerinus* (Linnaeus 1758), *Forpus xanthopterygius* (Spix 1824), *Furnarius leucopus* Swainson 1838, *Galbula ruficauda* Cuvier 1816, *Gallinula chloropus* (Linnaeus 1758), *Gampsonyx swainsonii*, *Geranospiza caerulescens*, *Glaucidium brasilianum* (Gmelin 1788), *Harpagus bidentatus*, *Harpia harpyja*, *Helicolestes hamatus*, *Hemitriccus margaritaceiventer* (Orbigny & Lafresnaye 1838), *Himantopus mexicanus* (Statius Muller 1776), *Hylophilus flavipes* Lafresnaye 1845, *Hypnelus ruficollis* (Wagler 1829), *Icterus auricapillus* Cassin 1848, *Icterus galbula* (Linnaeus 1758), *Icterus nigrogularis* (Hahn 1819), *Ictinia mississippiensis*, *Ictinia plumbea*, *Inezia subflava* (P. L. Sclater & Salvin 1873), *Inezia tenuirostris* (Cory 1913), *Jabiru mycteria*, *Jacana jacana*, *Lepidopygia*

goudoti (Bourcier 1843), *Leptotila verreauxi* (Bonaparte 1855), *Leucopternis melanops*, *Leucopternis semiplumbea*, *Machetornis rixosa*, *Megaceryle alcyon* (Linnaeus 1758), *Megaceryle torquata* (Linnaeus 1766), *Megarynchus pitangua* (Linnaeus 1766), *Megascops choliba*, *Melanerpes rubricapillus*, *Micrastur plumbeus*, *Mitu tomentosa*, *Molothrus bonariensis*, *Morphnarchus princeps*, *Morphnus guianensis*, *Myiarchus crinitus*, *Myiarchus venezuelensis* Lawrence 1865, *Myiopagis gaimardii* (Orbigny 1840), *Myiozetetes similis* (Spix 1825), *Myospiza humeralis* (Bosc 1792), *Nemosia pileata* (Boddaert 1783), *Neochen jubata*, *Opisthocomus hoazin*, *Oroaetus isidori*, *Ortalis ruficauda* Jardine 1847, *Pandion haliaetus*, *Parabuteo unicinctus*, *Patagioenas corensis* (Jacquin 1784), *Phaeomyias murina* (Spix 1825), *Pheucticus ludovicianus*, *Piculus chrysochloros* (Vieillot 1818), *Picumnus cinnamomeus* Wagler 1829, *Pilherodius pileatus* (Boddaert 1783), *Pionus sordidus* (Linnaeus 1758), *Pitangus lictor* (Lichtenstein 1823), *Pitangus sulphuratus*, *Polioptila plumbea* (J. F. Gmelin 1788), *Pseudastur albicollis*, *Pulsatrix perspicillata* (Latham 1790), *Pyrocephalus rubinus*, *Quiscalus lugubris*, *Quiscalus mexicanus* (Gmelin 1788), *Rostrhamus sociabilis*, *Sakesphorus canadensis* (Linnaeus 1766), *Saltator coerulescens* Vieillot 1817, *Sarcoramphus papa*, *Sarkidiornis melanotos* (Pennant 1769), *Spiza americana*, *Spizaetus tyrannus*, *Sporophila bouvionides* (Lesson 1831), *Sporophila intermedia* Cabanis 1851, *Sporophila nigricollis* (Vieillot 1823), *Stelgidopteryx ruficollis* (Vieillot 1817), *Sterna supercilialis*, *Sturnella magna*, *Sublegatus arenarum* (Salvin 1863), *Synallaxis candei* Orbigny & Lafresnaye 1838, *Tachycineta bicolor* (Vieillot 1808), *Tapera naevia* (Linnaeus 1766), *Thamnophilus doliatus* (Linnaeus 1764), *Thraupis glaucocolpa* Cabanis 1850, *Thraupis palmarum*, *Thryothorus leucotis* (Lafresnaye 1845), *Tyrannus dominicensis* (Gmelin 1788), *Tyto alba*, *Vireo flavoviridis* (Cassin 1851), *Zenaida auriculata*

editors

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

issue:

year: 1986

authors

authors: Hilty, S

authors: Brown, W. L.

title: A Guide to the Birds of Colombia

source:

type: book

tags: *Accipiter collaris*, *Accipiter cooperii*, *Accipiter poliogaster*, *Accipiter superciliosus*, *Amazona ochrocephala*, *Andigena hypoglaucha*, *Aramus guarauna*, *Aratinga pertinax*, *Ardea alba*, *Ardea cocoi*, *Ardea herodias*, *Arremonops conirostris* (Bonaparte 1850), *Arremonops tocuyensis* (Todd 1912), *Atalotriccus pilaris*, *Attila spadiceus* (Gmelin 1789), *Basileuterus flaveolus* (S. F. Baird 1865), *Bubo virginianus* (Gmelin 1788), *Busarellus nigricollis*, *Buteogallus anthracinus*, *Buteogallus meridionalis*, *Buteogallus schistaceus*, *Buteogallus solitarius*, *Buteogallus urubitinga*, *Cacicus cela*, *Cairina moschata*, *Campylorhamphus trochilirostris* (Lichtenstein 1820), *Campylorhynchus griseus* (Swainson 1838), *Campylorhynchus nuchalis* (Cabanis 1847), *Caprimulgus cayennensis* (Gmelin 1789), *Cardinalis phoeniceus* Bonaparte 1838, *Carduelis psaltria* (Say 1823), *Cathartes aura*, *Cathartes burrovianus*, *Cathartes melambrotus*, *Certhiaxis cinnamomeus* (Gmelin 1788), *Ceryle torquata*, *Chlorostilbon gibsoni* (Fraser 1840), *Chondrohierax uncinatus*, *Chordeiles acutipennis* (Hermann 1783), *Chrysomus icterocephalus*, *Circus buffoni*, *Circus cinereus*, *Circus cyaneus*, *Coccyzus pumila*, *Columbina minuta* (Linnaeus 1766), *Conirostrum leucogenys* (Lafresnaye 1852), *Coragyps atratus*, *Coryphospingus pileatus* (Wied-Neuwied 1821), *Crax alector*, *Crotophaga major*, *Crotophaga sulcirostris* Swainson 1827, *Cryptoleucopteryx plumbea*, *Cyanocorax affinis* (Pelzelin 1856), *Cyclarhis gujanensis* (J. F. Gmelin 1789), *Dryocopus lineatus* (Linnaeus 1766), *Elaenia chiriquensis* Lawrence 1865, *Elanoides forficatus*, *Elanus leucurus*, *Falco deiroleucus*, *Fluvicola pica* (Boddaert 1783), *Forpus passerinus* (Linnaeus 1758), *Forpus xanthopterygius* (Spix 1824), *Furnarius leucopus* Swainson 1838, *Galbula ruficauda* Cuvier 1816, *Gallinula chloropus* (Linnaeus 1758), *Gampsonyx swainsonii*, *Geranospiza caerulescens*, *Glaucidium brasilianum* (Gmelin 1788), *Harpagus bidentatus*, *Harpia harpyja*, *Helicolestes hamatus*, *Hemitriccus margaritaceiventer* (Orbigny & Lafresnaye), *Himantopus mexicanus* (Statius)

Muller 1776),Hylophilus flavipes Lafresnaye 1845,Hypnelus ruficollis (Wagler 1829),Icterus auricapillus Cassin 1848,Icterus galbula (Linnaeus 1758),Icterus nigrogularis (Hahn 1819),Ictinia mississippiensis,Ictinia plumbea,Inezia subflava (P. L. Sclater \& Salvin 1873),Inezia tenuirostris (Cory 1913),Jabiru mycteria,Jacana jacana,Lepidopyga goudoti (Bourcier 1843),Leptotila verreauxi (Bonaparte 1855),Leucopternis melanops,Leucopternis semiplumbea,Machetornis rixosa,Megaceryle alcyon (Linnaeus 1758),Megaceryle torquata (Linnaeus 1766),Megarynchus pitangua (Linnaeus 1766),Megascops choliba,Melanerpes rubricapillus,Micrastur plumbeus,Mitu tomentosa,Molothrus bonariensis,Morphnarchus princeps,Morphnus guianensis,Myiarchus crinitus,Myiarchus venezuelensis Lawrence 1865,Myiopagis gaimardii (Orbigny 1840),Myiozetetes similis (Spix 1825),Myospiza humeralis (Bosc 1792),Nemosia pileata (Boddaert 1783),Neochen jubata,Opisthocomus hoazin,Oroaetus isidori,Ortalis ruficauda Jardine 1847,Pandion haliaetus,Parabuteo unicinctus,Patagioenas corensis (Jacquin 1784),Phaeomyias murina (Spix 1825),Pheucticus ludovicianus,Piculus chrysochloros (Vieillot 1818),Picumnus cinnamomeus Wagler 1829,Pilherodius pileatus (Boddaert 1783),Pionus sordidus (Linnaeus 1758),Pitangus lictor (Lichtenstein 1823),Pitangus sulphuratus,Polioptila plumbea (J. F. Gmelin 1788),Pseudastur albicollis,Pulsatrix perspicillata (Latham 1790),Pyrocephalus rubinus,Quiscalus lugubris,Quiscalus mexicanus (Gmelin 1788),Rostrhamus sociabilis,Sakesphorus canadensis (Linnaeus 1766),Saltator coerulescens Vieillot 1817,Sarcoramphus papa,Sarkidiornis melanotos (Pennant 1769),Spiza americana,Spizaetus tyrannus,Sporophila bouvronides (Lesson 1831),Sporophila intermedia Cabanis 1851,Sporophila nigricollis (Vieillot 1823),Stelgidopteryx ruficollis (Vieillot 1817),Sterna superciliaris,Sturnella magna,Sublegatus arenarum (Salvin 1863),Synallaxis candei Orbigny \& Lafresnaye 1838,Tachycineta bicolor (Vieillot 1808),Tapera naevia (Linnaeus 1766),Thamnophilus doliatus (Linnaeus 1764),Thraupis glaucocolpa Cabanis 1850,Thraupis palmarum,Thryothorus leucotis (Lafresnaye 1845),Tyrannus dominicensis (Gmelin 1788),Tyto alba,Vireo flavoviridis (Cassin 1851),Zenaida auriculata

identifiers

last_modified:
created:
group_id:
profile_id:
id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:
providerLink:

publicationStatus

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

replacementNameFor

microReference:
source:
datatype:
identifier:

source:
datatype:
identifier:
note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

[conservedAgainst](#)

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

[spellingCorrentionOf](#)

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion

typeOfType:

[lectotypeMicroReferenceVoucher](#)

[lectotypePublicationVoucher](#)

[voucherReference](#)

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

[authors](#)

[year](#)

simple:
canonicalAuthorship

[authors](#)

[year](#)

simple:
simple: (Linnaeus, 1766)

canonicalName

infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:
ref:
uninomial:

simple: Aramus guarauna
rank: SPECIES
simple: Aramus guarauna (Linnaeus, 1766)

attributes

nomenclaturalCode:
isAnamorphic:
id:

[synonymsAtomized](#)

[hierarchy](#)

[ancillaryData](#)

dataObject:
additionalInformation:

[reference](#)

geoPoint:
location:
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:
rightsHolder:
rights:
license:
title:
modified: 2016-03-15T05:03:06.469Z
created: 2016-03-15T05:03:06.469Z

[agent](#)

homepage:
role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:
parentTaxon: Aramus
higherClassification:

infraspecificEpithet:
specificEpithet:
taxonRank: SPECIES
subGenus:
genus: Aramus
family: Aramidae
order: Gruiformes
classHierarchy: Aves
phylum: Chordata
kingdom: Animalia
recommended:
classification:

[commonNameAtomized](#)

[ancillaryData](#)

usedBy:

usedIn

distributionUnstructured:
endDate:
startDate:

[ancillaryData](#)

type:
language: Español
name: Carao

invasiveness

invasivenessUnstructured:

reproduction

reproductionUnstructured:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

dataObject:

additionalInformation:

0

keywords: *Acacia glomerosa*, *Acacia tamarindifolia*, *Acacia tortuosa*, *Alternanthera halimifolia*, *Amazona ochrocephala*, *Ameiva festiva*, *Anas discors*, *Anemopaegma orbiculatum*, *Anhinga anhinga* (Linnaeus 1766), *Anoda crenatiflora*, *Aphelandra pulcherrima*, *Aramides cajanea*, *Aramus guarauna*, *Aratinga acuticaudata* (Vieillot 1818), *Ardea herodias*, *Aristolochia anguicida*, *Arremonops conirostris* (Bonaparte 1850), *Arremonops tocuyensis* (Todd 1912), *Artemisia vulgaris*, *Aspidosperma cuspa*, *Atalotriccus pilaris*, *Ayenia magna*, *Basileuterus flaveolus* (S. F. Baird 1865), *Bastardia viscosa*, *Batis maritima*, *Bauhinia pauletia*, *Boerhavia scandens*, *Bonellia frutescens*, *Bourreria cumanensis*, *Bouteloua aristoides*, *Bromelia chrysantha*, *Bubo virginianus* (Gmelin 1788), *Burhinus bistriatus* (Wagler 1829), *Butorides virescens*, *Caesalpinia bonduc*, *Caesalpinia mollis*, *Calotropis gigantea*, *Campylorhynchus griseus* (Swainson 1838), *Campylorhynchus nuchalis* (Cabanis 1847), *Canavalia brasiliensis*, *Canavalia rosea*, *Capparis frondosa*, *Capparis linearis*, *Capparis pulcherrima*, *Caprimulgus cayennensis* (Gmelin 1789), *Capsicum baccatum*, *Cardinalis phoeniceus* Bonaparte 1838, *Cardiospermum corindum*, *Carduelis psaltria* (Say 1823), *Castela erecta*, *Celtis iguanaea*, *Cenchrus brownii*, *Cenchrus ciliaris*, *Cenchrus echinatus*, *Certhiaxis cinnamomeus* (Gmelin 1788), *Chlorostilbon gibsoni* (Fraser 1840), *Chordeiles acutipennis* (Hermann 1783), *Chrysolampis mosquitos* (Linnaeus 1758), *Chrysomus icterocephalus*, *Cissus trifoliata*, *Citrullus lanatus*, *Cleome spinosa*, *Cnidoscopus urens*, *Coccoloba microstachya*, *Columbina minuta* (Linnaeus 1766), *Columbina passerina* (Linnaeus 1758), *Combretum fruticosum*, *Conirostrum leucogenys* (Lafresnaye 1852), *Conyza canadensis*, *Conyza perijaensis*, *Cordia macuirensis*, *Coryphospingus pileatus* (Wied-Neuwied 1821), *Craniolaria annua*, *Crotalaria purdiana*, *Crotalaria retusa*, *Croton ferrugineus*, *Croton malambo*, *Croton niveus*, *Croton ovalifolius*, *Croton punctatus*, *Crotophaga major*, *Crotophaga sulcirostris* Swainson 1827, *Cyanocorax affinis* (Pelzelin 1856), *Cyclarhis gujanensis* (J. F. Gmelin 1789), *Cyperus articulatus*, *Datura innoxia*, *Dendrocygna autumnalis*, *Desmodium incanum*, *Diphysa carthagenensis*, *Ditaxis argothamnoides*, *Dolichandra unguis-cati*, *Dryocopus lineatus* (Linnaeus 1766), *Dysphania ambrosioides*, *Egletes prostrata*, *Egretta thula*, *Egretta tricolor* (Statius Muller 1776), *Elaenia chiriquensis* Lawrence 1865, *Eleocharis elegans*, *Eleocharis interstincta*, *Engystomops pustulosus*, *Eragrostis ciliaris*, *Erythrina berteroana*, *Euphorbia hirta*, *Evolvulus convolvuloides*, *Evolvulus sericeus*, *Exostema caribaeum*, *Fluvicola pica* (Boddaert 1783), *Forpus passerinus* (Linnaeus 1758), *Forpus xanthopterygius* (Spix 1824), *Fridericia chica*, *Fridericia dichotoma*, *Fridericia mollissima*, *Funastrum clausum*, *Furnarius leucopus* Swainson 1838, *Galactia striata*, *Galbula ruficauda* Cuvier 1816, *Glaucidium brasilianum* (Gmelin 1788), *Glossophaga longirostris* Miller 1898, *Gonatodes vittatus*, *Handroanthus serratifolius*, *Heliotropium angiospermum*, *Heliotropium indicum*, *Heliotropium procumbens*, *Hemitriccus margaritaceiventer* (Orbigny & Lafresna), *Hibiscus phoeniceus*, *Hillia costanensis*, *Himantopus mexicanus* (Statius Muller 1776), *Hippomane mancinella*, *Hylophilus flavipes* Lafresnaye 1845, *Hymenocallis caribaea*, *Hypnelus ruficollis* (Wagler 1829), *Icterus auricapillus* Cassin 1848, *Icterus galbula* (Linnaeus 1758), *Icterus nigrogularis* (Hahn 1819), *Inezia subflava* (P. L. Sclater & Salvin 1873), *Inezia tenuirostris* (Cory 1913), *Ipomoea incarnata*, *Jacquinia armillaris*, *Laguncularia racemosa*, *Lantana fucata*, *Lepidoblepharis sanctaemartae* Ruthven 1916, *Leptodactylus poecilochilus*, *Leptonycteris curasoae* Miller 1900, *Leptotila verreauxi* (Bonaparte 1855), *Licania cuspidata*, *Lonchocarpus atropurpureus*, *Lonchocarpus sanctae-marthae*, *Ludwigia octovalvis*, *Luffa operculata*, *Machetornis rixosa*, *Mallotus rhamnifolius*, *Manihot carthaginensis*, *Matelea maritima*, *Megaceryle alcyon* (Linnaeus 1758), *Megaceryle torquata* (Linnaeus 1766), *Megarynchus pitangua* (Linnaeus 1766), *Megascops choliba*, *Melanerpes rubricapillus*, *Melocactus intortus*, *Melochia parvifolia*, *Melochia tomentosa*, *Mentzelia aspera*, *Merremia aegyptia*, *Merremia umbellata*, *Micronycteris hirsuta* (Peters 1869), *Mimosa arenosa*, *Mimosa leiocarpa*, *Mimosa pellita*, *Mollugo verticillata*, *Molothrus bonariensis*, *Morisonia americana*, *Mormoops megalophylla* (Peters 1864), *Musa paradisiaca*, *Myiarchus crinitus*, *Myiarchus venezuelensis* Lawrence 1865, *Myiopagis gaimardii* (Orbigny 1840), *Myiozetetes similis* (Spix 1825), *Myospiza humeralis* (Bosc 1792), *Nemosia pileata* (Boddaert 1783), *Neurolaena lobata*, *Noctilio leporinus* (Linnaeus 1758), *Nycticorax nycticorax*, *Pachyrhamphus rufus* (Boddaert 1783), *Parkia velutina*, *Parkinsonia praecox*, *Passiflora foetida*, *Patagioenas corensis* (Jacquin 1784), *Paullinia cururu*, *Phaeomyias murina* (Spix 1825), *Phaseolus lunatus*, *Pheucticus ludovicianus*, *Phoradendron mucronatum*, *Phthirusa stelis*, *Phyla nodiflora*, *Piculus chrysochloros* (Vieillot 1818), *Picumnus cinnamomeus* Wagler 1829, *Pilherodius pileatus* (Boddaert 1783), *Pilosocereus lanuginosus*, *Pionus sordidus* (Linnaeus 1758), *Piptadenia viridiflora*, *Piranga flava* (Vieillot 1822), *Pitangus lictor* (Lichtenstein 1823), *Platalea ajaja* Linnaeus 1758, *Pleurodema brachyops*, *Polioptila plumbea* (J. F. Gmelin 1788), *Porophyllum ruderales*, *Pseudopaludicola pusilla*, *Pteronotus gymnonotus* (Natterer 1843), *Pteronotus parnellii* (Gray 1843), *Pteronotus personatus* (Wagner 1843), *Pulsatrix perspicillata* (Latham 1790), *Pyrocephalus rubinus*, *Quiscalus lugubris*, *Rhizophora mangle*, *Rivina*

humilis, Rosenbergiodendron formosum, Ruellia paniculata, Ruprechtia ramiflora, Sakesphorus canadensis (Linnaeus 1766), Saltator coerulescens Vieillot 1817, Sarcostemma glaucum, Scoparia dulcis, Senna atomaria, Senna bicapsularis, Senna candolleana, Senna italica, Senna occidentalis, Senna tora, Serjania mexicana, Sesuvium portulacastrum, Sida ciliaris, Solandra grandiflora, Solanum bicolor, Spiza americana, Sporobolus coromandelianus, Sporobolus pyramidalis, Sporophila bouvronides (Lesson 1831), Sporophila intermedia Cabanis 1851, Sporophila nigricollis (Vieillot 1823), Stelgidopteryx ruficollis (Vieillot 1817), Stemodia maritima, Streptoprocne zonaris (Shaw 1796), Sturnella magna, Sublegatus arenarum (Salvin 1863), Synallaxis candei Orbigny & Lafresnaye 1838, Tabernaemontana amygdalifolia, Tachycineta albiventer (Boddaert 1783), Tachycineta bicolor (Vieillot 1808), Talinum paniculatum, Tapera naevia (Linnaeus 1766), Tephrosia cinerea, Thamnophilus doliatus (Linnaeus 1764), Thraupis glaucocolpa Cabanis 1850, Thraupis palmarum, Thryothorus leucotis (Lafresnaye 1845), Tiaris bicolor (Linnaeus 1766), Tiaris obscurus (Orbigny & Lafresnaye 1837), Tillandsia flexuosa, Tityra inquisitor (Lichtenstein 1823), Tolmomyias sulphurescens (Spix 1825), Tournefortia hirsutissima, Trianthema portulacastrum, Tribulus cistoides, Tyrannus dominicensis (Gmelin 1788), Tyto alba, Vireo flavoviridis (Cassin 1851), Vitex compressa, Wissadula periplocifolia, Zenaida auriculata

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

pages: 417

issue:

year: 2010

0: Cerrejón

title: Monitoreo de Fauna en el Valle del Cerrejón

source:

type: report

tags: Acacia glomerosa, Acacia tamarindifolia, Acacia tortuosa, Alternanthera halimifolia, Amazona ochrocephala, Ameiva festiva, Anas discors, Anemopaegma orbiculatum, Anhinga anhinga (Linnaeus 1766), Anoda crenatiflora, Aphelandra pulcherrima, Aramides cajanea, Aramus guarauna, Aratinga acuticaudata (Vieillot 1818), Ardea herodias, Aristolochia anguicida, Arremonops conirostris (Bonaparte 1850), Arremonops tocuyensis (Todd 1912), Artemisia vulgaris, Aspidosperma cuspa, Atalotriccus pilaris, Ayenia magna, Basileuterus flaveolus (S. F. Baird 1865), Bastardia viscosa, Batis maritima, Bauhinia pauletia, Boerhavia scandens, Bonellia frutescens, Bourreria cumanaensis, Bouteloua aristoides, Bromelia chrysantha, Bubo virginianus (Gmelin 1788), Burhinus bistriatus (Wagler 1829), Butorides virescens, Caesalpinia bonduc, Caesalpinia mollis, Calotropis gigantea, Campylorhynchus griseus (Swainson 1838), Campylorhynchus nuchalis (Cabanis 1847), Canavalia brasiliensis, Canavalia rosea, Capparis frondosa, Capparis linearis, Capparis pulcherrima, Caprimulgus cayennensis (Gmelin 1789), Capsicum baccatum, Cardinalis phoeniceus Bonaparte 1838, Cardiospermum corindum, Carduelis psaltria (Say 1823), Castela erecta, Celtis iguanaea, Cenchrus brownii, Cenchrus ciliaris, Cenchrus echinatus, Certhiaxis cinnamomeus (Gmelin 1788), Chlorostilbon gibsoni (Fraser 1840), Chordeiles acutipennis (Hermann 1783), Chrysolampis mosquitos (Linnaeus 1758), Chrysomus icterocephalus, Cissus trifoliata, Citrullus lanatus, Cleome spinosa, Cnidoscolus urens, Coccoloba microstachya, Columbina minuta (Linnaeus 1766), Columbina passerina (Linnaeus 1758), Combretum fruticosum, Conirostrum leucogenys (Lafresnaye 1852), Conyza canadensis, Conyza perijaensis, Cordia macuirensis, Coryphospingus pileatus (Wied-Neuwied 1821), Craniolaria annua, Crotalaria purdiana, Crotalaria retusa, Croton ferrugineus, Croton malambo, Croton niveus, Croton ovalifolius, Croton punctatus, Crotophaga major, Crotophaga sulcirostris Swainson 1827, Cyanocorax affinis (Pelzeln 1856), Cyclarhis gujanensis (J. F. Gmelin 1789), Cyperus articulatus, Datura innoxia, Dendrocygna autumnalis, Desmodium incanum, Diphysa carthagenensis, Ditaxis argothamnoides, Dolichandra unguis-cati, Dryocopus lineatus (Linnaeus 1766), Dysphania ambrosioides, Egletes prostrata, Egretta thula, Egretta tricolor (Statius Muller 1776), Elaenia chiriquensis Lawrence 1865, Eleocharis elegans, Eleocharis interstincta, Engystomops pustulosus, Eragrostis ciliaris, Erythrina berteroana, Euphorbia hirta, Evolvulus convolvuloides, Evolvulus sericeus, Exostema caribaeum, Fluvicola pica (Boddaert 1783), Forpus passerinus (Linnaeus 1758), Forpus xanthopterygius (Spix 1824), Fridericia chica, Fridericia dichotoma, Fridericia mollissima, Funastrum clausum, Furnarius leucopus Swainson 1838, Galactia striata, Galbula ruficauda Cuvier 1816, Glaucidium brasilianum (Gmelin 1788), Glossophaga longirostris Miller 1898, Gonatodes vittatus, Handroanthus serratifolius, Heliotropium angiospermum, Heliotropium

indicum, Heliotropium procumbens, Hemitriccus margaritaceiventer (Orbigny \& Lafresna, Hibiscus phoeniceus, Hillia costanensis, Himantopus mexicanus (Statius Muller 1776), Hippomane mancinella, Hylophilus flavipes Lafresnaye 1845, Hymenocallis caribaea, Hypnelus ruficollis (Wagler 1829), Icterus auricapillus Cassin 1848, Icterus galbula (Linnaeus 1758), Icterus nigrogularis (Hahn 1819), Inezia subflava (P. L. Sclater \& Salvin 1873), Inezia tenuirostris (Cory 1913), Ipomoea incarnata, Jacquinia armillaris, Laguncularia racemosa, Lantana fucata, Lepidoblepharis sanctaemartae Ruthven 1916, Leptodactylus poecilochilus, Leptonycteris curasoae Miller 1900, Leptotila verreauxi (Bonaparte 1855), Licania cuspidata, Lonchocarpus atropurpureus, Lonchocarpus sanctae-marthae, Ludwigia octovalvis, Luffa operculata, Machetornis rixosa, Mallotus rhamnifolius, Manihot carthaginensis, Matelea maritima, Megaceryle alcyon (Linnaeus 1758), Megaceryle torquata (Linnaeus 1766), Megarynchus pitangua (Linnaeus 1766), Megascops choliba, Melanerpes rubricapillus, Melocactus intortus, Melochia parvifolia, Melochia tomentosa, Mentzelia aspera, Merremia aegyptia, Merremia umbellata, Micronycteris hirsuta (Peters 1869), Mimosa arenosa, Mimosa leiocarpa, Mimosa pellita, Mollugo verticillata, Molothrus bonariensis, Morisonia americana, Mormoops megalophylla (Peters 1864), Musa paradisiaca, Myiarchus crinitus, Myiarchus venezuelensis Lawrence 1865, Myiopagis gaimardii (Orbigny 1840), Myiozetetes similis (Spix 1825), Myospiza humeralis (Bosc 1792), Nemosia pileata (Boddaert 1783), Neurolaena lobata, Noctilio leporinus (Linnaeus 1758), Nycticorax nycticorax, Pachyrhamphus rufus (Boddaert 1783), Parkia velutina, Parkinsonia praecox, Passiflora foetida, Patagioenas corensis (Jacquin 1784), Paullinia cururu, Phaeomyias murina (Spix 1825), Phaseolus lunatus, Pheucticus ludovicianus, Phoradendron mucronatum, Phthirusa stelis, Phyla nodiflora, Piculus chrysochloros (Vieillot 1818), Picumnus cinnamomeus Wagler 1829, Pilherodius pileatus (Boddaert 1783), Pilosocereus lanuginosus, Pionus sordidus (Linnaeus 1758), Piptadenia viridiflora, Piranga flava (Vieillot 1822), Pitangus lictor (Lichtenstein 1823), Platalea ajaja Linnaeus 1758, Pleurodema brachyops, Polioptila plumbea (J. F. Gmelin 1788), Porophyllum ruderae, Pseudopaludicola pusilla, Pteronotus gymnotus (Natterer 1843), Pteronotus parnellii (Gray 1843), Pteronotus personatus (Wagner 1843), Pulsatrix perspicillata (Latham 1790), Pyrocephalus rubinus, Quiscalus lugubris, Rhizophora mangle, Rivina humilis, Rosenbergiodendron formosum, Ruellia paniculata, Ruprechtia ramiflora, Sakesphorus canadensis (Linnaeus 1766), Saltator coerulescens Vieillot 1817, Sarcostemma glaucum, Scoparia dulcis, Senna atomaria, Senna bicapsularis, Senna candolleana, Senna italica, Senna occidentalis, Senna tora, Serjania mexicana, Sesuvium portulacastrum, Sida ciliaris, Solandra grandiflora, Solanum bicolor, Spiza americana, Sporobolus coromandelianus, Sporobolus pyramidalis, Sporophila bouvronides (Lesson 1831), Sporophila intermedia Cabanis 1851, Sporophila nigricollis (Vieillot 1823), Stelgidopteryx ruficollis (Vieillot 1817), Stemodia maritima, Streptoprocne zonaris (Shaw 1796), Sturnella magna, Sublegatus arenarum (Salvin 1863), Synallaxis candei Orbigny \& Lafresnaye 1838, Tabernaemontana amygdalifolia, Tachycineta albiventer (Boddaert 1783), Tachycineta bicolor (Vieillot 1808), Talinum paniculatum, Tapera naevia (Linnaeus 1766), Tephrosia cinerea, Thamnophilus doliatus (Linnaeus 1764), Thraupis glaucocolpa Cabanis 1850, Thraupis palmarum, Thryothorus leucotis (Lafresnaye 1845), Tiaris bicolor (Linnaeus 1766), Tiaris obscurus (Orbigny \& Lafresnaye 1837), Tillandsia flexuosa, Tityra inquisitor (Lichtenstein 1823), Tolmomyias sulphurescens (Spix 1825), Tournefortia hirsutissima, Trianthema portulacastrum, Tribulus cistoides, Tyrannus dominicensis (Gmelin 1788), Tyto alba, Vireo flavoviridis (Cassin 1851), Vitex compressa, Wissadula periplocifolia, Zenaida auriculata

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

feedingUnstructured: En La Guajira forrajea en el suelo y el agua (Cerrejón 2009-2010), los caracoles hacen parte de su dieta (Cerrejón 2010).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

0

keywords: Accipiter collaris, Accipiter cooperii, Accipiter poliogaster, Accipiter superciliosus, Amazona ochrocephala, Andigena hypoglaucha, Aramus guarauna, Aratinga pertinax, Ardea alba, Ardea cocoi, Ardea herodias, Arremonops conirostris (Bonaparte 1850), Arremonops tocuyensis (Todd 1912), Atalotriccus pilaris, Attila spadiceus (Gmelin 1789), Basileuterus flaveolus (S. F. Baird 1865), Bubo virginianus (Gmelin 1788), Busarellus nigricollis, Buteogallus anthracinus, Buteogallus meridionalis, Buteogallus schistaceus, Buteogallus solitarius, Buteogallus urubitinga, Cacicus cela, Cairina moschata, Campylorhamphus trochilirostris (Lichtenstein 1820), Campylorhynchus griseus (Swainson 1838), Campylorhynchus nuchalis (Cabanis 1847), Caprimulgus cayennensis (Gmelin 1789), Cardinalis phoeniceus Bonaparte 1838, Carduelis psaltria (Say 1823), Cathartes aura, Cathartes burrovianus, Cathartes melambrotus, Certhiaxis cinnamomeus (Gmelin 1788), Ceryle torquata, Chlorostilbon gibsoni (Fraser 1840), Chondrohierax uncinatus, Chordeiles acutipennis (Hermann 1783), Chrysomus icterocephalus, Circus buffoni, Circus cinereus, Circus cyaneus, Coccyzus pumila, Columbina minuta (Linnaeus 1766), Conirostrum leucogenys (Lafresnaye 1852), Coragyps atratus, Coryphospingus pileatus (Wied-Neuwied 1821), Crax alector, Crotophaga major, Crotophaga sulcirostris Swainson 1827, Cryptoleucopteryx plumbea, Cyanocorax affinis (Pelzeln 1856), Cyclarhis gujanensis (J. F. Gmelin 1789), Dryocopus lineatus (Linnaeus 1766), Elaenia chiriquensis Lawrence 1865, Elanoides forficatus, Elanus leucurus, Falco deiroleucus, Fluvicola pica (Boddaert 1783), Forpus passerinus (Linnaeus 1758), Forpus xanthopterygius (Spix 1824), Furnarius leucopus Swainson 1838, Galbula ruficauda Cuvier 1816, Gallinula chloropus (Linnaeus 1758), Gampsonyx swainsonii, Geranospiza caerulescens, Glaucidium brasilianum (Gmelin 1788), Harpagus bidentatus, Harpia harpyja, Helicolestes hamatus, Hemitriccus margaritaceiventer (Orbigny & Lafresna, Himantopus mexicanus (Statius Muller 1776), Hylophilus flavipes Lafresnaye 1845, Hypnelus ruficollis (Wagler 1829), Icterus auricapillus Cassin 1848, Icterus galbula (Linnaeus 1758), Icterus nigrogularis (Hahn 1819), Ictinia mississippiensis, Ictinia plumbea, Inezia subflava (P. L. Sclater & Salvin 1873), Inezia tenuirostris (Cory 1913), Jabiru mycteria, Jacana jacana, Lepidopyga goudoti (Bourcier 1843), Leptotila verreauxi (Bonaparte 1855), Leucopternis melanops, Leucopternis semiplumbea, Machetornis rixosa, Megaceryle alcyon (Linnaeus 1758), Megaceryle torquata (Linnaeus 1766), Megarynchus pitangua (Linnaeus 1766), Megascops choliba, Melanerpes rubricapillus, Micrastur plumbeus, Mitu tomentosa, Molothrus bonariensis, Morphnarchus princeps, Morphnus guianensis, Myiarchus crinitus, Myiarchus venezuelensis Lawrence 1865, Myiopagis gaimardii (Orbigny 1840), Myiozetetes similis (Spix 1825), Myospiza humeralis (Bosc 1792), Nemosia pileata (Boddaert 1783), Neechen jubata, Opisthocomus hoazin, Oroaetus isidori, Ortalis ruficauda Jardine 1847, Pandion haliaetus, Parabuteo unicinctus, Patagioenas corensis (Jacquin 1784), Phaeomyias murina (Spix 1825), Pheucticus ludovicianus, Piculus chrysochloros (Vieillot 1818), Picumnus

cinnamomeus Wagler 1829, *Pilherodius pileatus* (Boddaert 1783), *Pionus sordidus* (Linnaeus 1758), *Pitangus lictor* (Lichtenstein 1823), *Pitangus sulphuratus*, *Polioptila plumbea* (J. F. Gmelin 1788), *Pseudastur albicollis*, *Pulsatrix perspicillata* (Latham 1790), *Pyrocephalus rubinus*, *Quiscalus lugubris*, *Quiscalus mexicanus* (Gmelin 1788), *Rostrhamus sociabilis*, *Sakesphorus canadensis* (Linnaeus 1766), *Saltator coerulescens* Vieillot 1817, *Sarcoramphus papa*, *Sarkidiornis melanotos* (Pennant 1769), *Spiza americana*, *Spizaetus tyrannus*, *Sporophila bouvronides* (Lesson 1831), *Sporophila intermedia* Cabanis 1851, *Sporophila nigricollis* (Vieillot 1823), *Stelgidopteryx ruficollis* (Vieillot 1817), *Sterna supercilialis*, *Sturnella magna*, *Sublegatus arenarum* (Salvin 1863), *Synallaxis candei* Orbigny & Lafresnaye 1838, *Tachycineta bicolor* (Vieillot 1808), *Tapera naevia* (Linnaeus 1766), *Thamnophilus doliatus* (Linnaeus 1764), *Thraupis glaucocolpa* Cabanis 1850, *Thraupis palmarum*, *Thryothorus leucotis* (Lafresnaye 1845), *Tyrannus dominicensis* (Gmelin 1788), *Tyto alba*, *Vireo flavoviridis* (Cassin 1851), *Zenaida auriculata*

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

issue:

year: 1986

0: Hilty, S

1: Brown, W. L.

title: A Guide to the Birds of Colombia

source:

type: book

tags: *Accipiter collaris*, *Accipiter cooperii*, *Accipiter poliogaster*, *Accipiter superciliosus*, *Amazona ochrocephala*, *Andigena hypoglaucha*, *Aramus guarauna*, *Aratinga pertinax*, *Ardea alba*, *Ardea cocoi*, *Ardea herodias*, *Arremonops conirostris* (Bonaparte 1850), *Arremonops tocuyensis* (Todd 1912), *Atalotriccus pilaris*, *Attila spadiceus* (Gmelin 1789), *Basileuterus flaveolus* (S. F. Baird 1865), *Bubo virginianus* (Gmelin 1788), *Busarellus nigricollis*, *Buteogallus anthracinus*, *Buteogallus meridionalis*, *Buteogallus schistaceus*, *Buteogallus solitarius*, *Buteogallus urubitinga*, *Caciclus cela*, *Cairina moschata*, *Campylorhamphus trochilrostris* (Lichtenstein 1820), *Campylorhynchus griseus* (Swainson 1838), *Campylorhynchus nuchalis* (Cabanis 1847), *Caprimulgus cayennensis* (Gmelin 1789), *Cardinalis phoeniceus* Bonaparte 1838, *Carduelis psaltria* (Say 1823), *Cathartes aura*, *Cathartes burrovianus*, *Cathartes melambrotus*, *Certhiaxis cinnamomeus* (Gmelin 1788), *Ceryle torquata*, *Chlorostilbon gibsoni* (Fraser 1840), *Chondrohierax uncinatus*, *Chordeiles acutipennis* (Hermann 1783), *Chrysomus icterocephalus*, *Circus buffoni*, *Circus cinereus*, *Circus cyaneus*, *Coccyzus pumila*, *Columbina minuta* (Linnaeus 1766), *Conirostrum leucogenys* (Lafresnaye 1852), *Coragyps atratus*, *Coryphospingus pileatus* (Wied-Neuwied 1821), *Crax alector*, *Crotophaga major*, *Crotophaga sulcirostris* Swainson 1827, *Cryptoleucopteryx plumbea*, *Cyanocorax affinis* (Pelzelin 1856), *Cyclarhis gujanensis* (J. F. Gmelin 1789), *Dryocopus lineatus* (Linnaeus 1766), *Elaenia chiriquensis* Lawrence 1865, *Elanoides forficatus*, *Elanus leucurus*, *Falco deiroleucus*, *Fluvicola pica* (Boddaert 1783), *Forpus passerinus* (Linnaeus 1758), *Forpus xanthopterygius* (Spix 1824), *Furnarius leucopus* Swainson 1838, *Galbula ruficauda* Cuvier 1816, *Gallinula chloropus* (Linnaeus 1758), *Gampsonyx swainsonii*, *Geranospiza caerulescens*, *Glaucidium brasilianum* (Gmelin 1788), *Harpagus bidentatus*, *Harpia harpyja*, *Helicolestes hamatus*, *Hemitriccus margaritaceiventer* (Orbigny & Lafresna, *Himantopus mexicanus* (Statius Muller 1776), *Hylophilus flavipes* Lafresnaye 1845, *Hypnelus ruficollis* (Wagler 1829), *Icterus auricapillus* Cassin 1848, *Icterus galbula* (Linnaeus 1758), *Icterus nigrogularis* (Hahn 1819), *Ictinia mississippiensis*, *Ictinia plumbea*, *Inezia subflava* (P. L. Sclater & Salvin 1873), *Inezia tenuirostris* (Cory 1913), *Jabiru mycteria*, *Jacana jacana*, *Lepidopyga goudoti* (Bourcier 1843), *Leptotila verreauxi* (Bonaparte 1855), *Leucopternis melanops*, *Leucopternis semiplumbea*, *Machetornis rixosa*, *Megaceryle alcyon* (Linnaeus 1758), *Megaceryle torquata* (Linnaeus 1766), *Megarynchus pitangua* (Linnaeus 1766), *Megascops choliba*, *Melanerpes rubricapillus*, *Micrastur plumbeus*, *Mitotomentosa*, *Molothrus bonariensis*, *Morphnarchus princeps*, *Morphnus guianensis*, *Myiarchus crinitus*, *Myiarchus venezuelensis* Lawrence 1865, *Myiopagis gaimardii* (Orbigny 1840), *Myiozetetes similis* (Spix 1825), *Myospiza humeralis* (Bosc 1792), *Nemosia pileata* (Boddaert 1783), *Neochen jubata*, *Opisthocomus hoazin*, *Oroaetus isidori*, *Ortalis ruficauda* Jardine 1847, *Pandion haliaetus*, *Parabuteo unicinctus*, *Patagioenas corensis* (Jacquin 1784), *Phaeomyias murina* (Spix 1825), *Pheucticus ludovicianus*, *Piculus chrysochloros* (Vieillot 1818), *Picumnus cinnamomeus* Wagler 1829, *Pilherodius pileatus* (Boddaert 1783), *Pionus sordidus* (Linnaeus 1758), *Pitangus lictor* (Lichtenstein 1823), *Pitangus sulphuratus*, *Polioptila plumbea* (J. F. Gmelin 1788), *Pseudastur albicollis*, *Pulsatrix*

perspicillata (Latham 1790),Pyrocephalus rubinus,Quiscalus lugubris,Quiscalus mexicanus (Gmelin 1788),Rostrhamus sociabilis,Sakesphorus canadensis (Linnaeus 1766),Saltator coerulescens Vieillot 1817,Sarcoramphus papa,Sarkidiornis melanotos (Pennant 1769),Spiza americana,Spizaetus tyrannus,Sporophila bouvronides (Lesson 1831),Sporophila intermedia Cabanis 1851,Sporophila nigricollis (Vieillot 1823),Stelgidopteryx ruficollis (Vieillot 1817),Sterna superciliaris,Sturnella magna,Sublegatus arenarum (Salvin 1863),Synallaxis candei Orbigny & Lafresnaye 1838,Tachycineta bicolor (Vieillot 1808),Tapera naevia (Linnaeus 1766),Thamnophilus doliatus (Linnaeus 1764),Thraupis glaucocolpa Cabanis 1850,Thraupis palmarum,Thryothorus leucotis (Lafresnaye 1845),Tyrannus dominicensis (Gmelin 1788),Tyto alba,Vireo flavoviridis (Cassin 1851),Zenaida auriculata

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

behaviorUnstructured: Parcialmente nocturno aunque normalmente activo durante el día. Explora cuerpos de agua someros o vegetación de pantano; cuando no está comiendo puede posarse en arbustos bajos. Vuelo boyante con aletazos rígidos hacia arriba. A menudo solitario, pero durante los periodos secos varios pueden concentrarse en pequeñas charcas y aun en sitios de lodo agrietado (Hilty & Brown 1986). A menudo ruidoso, especialmente al atardecer o durante la noche; lamento potente cariii-iii-ooou o criiii-ooou, repetido una y otra vez (Hilty & Brown 1986).

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source: <http://www.biologyreference.com/Ar-Bi/Behavior-Patterns.html>

bibliographicCitation: Alcock, John. Animal Behavior: An Evolutionary Approach, 6th ed. Sunderland, MA: Sinauer Associates, Inc., 1997.

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue: Solitario
measurementType: Dispersión y Comportamiento Social
measurementID:

annualCycle

annualCycleUnstructured:

habitat

0

dataObject:
additionalInformation:

0

keywords: *Acacia glomerosa*, *Acacia tamarindifolia*, *Acacia tortuosa*, *Alternanthera halimifolia*, *Amazona ochrocephala*, *Ameiva festiva*, *Anas discors*, *Anemopaegma orbiculatum*, *Anhinga anhinga* (Linnaeus 1766), *Anoda crenatiflora*, *Aphelandra pulcherrima*, *Aramides cajanea*, *Aramus guarauna*, *Aratinga acuticaudata* (Vieillot 1818), *Ardea herodias*, *Aristolochia anguicida*, *Arremonops conirostris* (Bonaparte 1850), *Arremonops tocuyensis* (Todd 1912), *Artemisia vulgaris*, *Aspidosperma cuspa*, *Atalotriccus pilaris*, *Ayenia magna*, *Basileuterus flaveolus* (S. F. Baird 1865), *Bastardia viscosa*, *Batis maritima*, *Bauhinia pauletia*, *Boerhavia scandens*, *Bonellia frutescens*, *Bourreria cumanensis*, *Bouteloua aristoides*, *Bromelia chrysantha*, *Bubo virginianus* (Gmelin 1788), *Burhinus bistriatus* (Wagler 1829), *Butorides virescens*, *Caesalpinia bonduc*, *Caesalpinia mollis*, *Calotropis gigantea*, *Campylorhynchus griseus* (Swainson 1838), *Campylorhynchus nuchalis* (Cabanis 1847), *Canavalia brasiliensis*, *Canavalia rosea*, *Capparis frondosa*, *Capparis linearis*, *Capparis pulcherrima*, *Caprimulgus cayennensis* (Gmelin 1789), *Capsicum baccatum*, *Cardinalis phoeniceus* Bonaparte 1838, *Cardiospermum corindum*, *Carduelis psaltria* (Say 1823), *Castela erecta*, *Celtis iguanaea*, *Cenchrus brownii*, *Cenchrus ciliaris*, *Cenchrus echinatus*, *Certhiaxis cinnamomeus* (Gmelin 1788), *Chlorostilbon gibsoni* (Fraser 1840), *Chordeiles acutipennis* (Hermann 1783), *Chrysolampis mosquitos* (Linnaeus 1758), *Chrysomus icterocephalus*, *Cissus trifoliata*, *Citrullus lanatus*, *Cleome spinosa*, *Cnidoscopus urens*, *Coccoloba microstachya*, *Columbina minuta* (Linnaeus 1766), *Columbina passerina* (Linnaeus 1758), *Combretum fruticosum*, *Conirostrum leucogenys* (Lafresnaye 1852), *Conyza canadensis*, *Conyza perijaensis*, *Cordia macuirensis*, *Coryphospingus pileatus* (Wied-Neuwied 1821), *Craniolaria annua*, *Crotalaria purdiana*, *Crotalaria retusa*, *Croton ferrugineus*, *Croton malambo*, *Croton niveus*, *Croton ovalifolius*, *Croton punctatus*, *Crotophaga major*, *Crotophaga sulcirostris* Swainson 1827, *Cyanocorax affinis* (Pelzelin 1856), *Cyclarhis gujanensis* (J. F. Gmelin 1789), *Cyperus articulatus*, *Datura innoxia*, *Dendrocygna autumnalis*, *Desmodium incanum*, *Diphysa carthagenensis*, *Ditaxis argothamnoides*, *Dolichandra unguis-cati*, *Dryocopus lineatus* (Linnaeus 1766), *Dysphania ambrosioides*, *Egletes prostrata*, *Egretta thula*, *Egretta tricolor* (Statius Muller 1776), *Elaenia chiriquensis* Lawrence 1865, *Eleocharis elegans*, *Eleocharis interstincta*, *Engystomops pustulosus*, *Eragrostis ciliaris*, *Erythrina berteriana*, *Euphorbia hirta*, *Evolvulus convolvuloides*, *Evolvulus sericeus*, *Exostema caribaeum*, *Fluvicola pica* (Boddaert 1783), *Forpus passerinus* (Linnaeus 1758), *Forpus xanthopterygius* (Spix 1824), *Fridericia chica*, *Fridericia dichotoma*, *Fridericia mollissima*, *Funastrum clausum*, *Furnarius leucopus* Swainson 1838, *Galactia striata*, *Galbula ruficauda* Cuvier 1816, *Glaucidium brasilianum* (Gmelin 1788), *Glossophaga longirostris* Miller 1898, *Gonatodes vittatus*, *Handroanthus serratifolius*, *Heliotropium angiospermum*, *Heliotropium indicum*, *Heliotropium procumbens*, *Hemitriccus margaritaceiventer* (Orbigny & Lafresna), *Hibiscus phoeniceus*, *Hillia costanensis*, *Himantopus mexicanus* (Statius Muller 1776), *Hippomane mancinella*, *Hylophilus flavipes* Lafresnaye 1845, *Hymenocallis caribaea*, *Hypnelus ruficollis* (Wagler 1829), *Icterus auricapillus* Cassin 1848, *Icterus galbula* (Linnaeus 1758), *Icterus nigrogularis* (Hahn 1819), *Inezia subflava* (P. L. Sclater & Salvin 1873), *Inezia tenuirostris* (Cory 1913), *Ipomoea incarnata*, *Jacquinia armillaris*, *Laguncularia racemosa*, *Lantana fucata*, *Lepidoblepharis sanctaemartae* Ruthven 1916, *Leptodactylus poecilochilus*, *Leptonycteris curasoae* Miller 1900, *Leptotila verreauxi* (Bonaparte 1855), *Licania cuspidata*, *Lonchocarpus atropurpureus*, *Lonchocarpus sanctae-marthae*, *Ludwigia*

octovalvis, Luffa operculata, Machetornis rixosa, Mallotus rhamnifolius, Manihot carthaginensis, Matelea maritima, Megaceryle alcyon (Linnaeus 1758), Megaceryle torquata (Linnaeus 1766), Megarynchus pitangua (Linnaeus 1766), Megascops choliba, Melanerpes rubricapillus, Melocactus intortus, Melochia parvifolia, Melochia tomentosa, Mentzelia aspera, Merremia aegyptia, Merremia umbellata, Micronycteris hirsuta (Peters 1869), Mimosa arenosa, Mimosa leiocarpa, Mimosa pellita, Mollugo verticillata, Molothrus bonariensis, Morisonia americana, Mormoops megalophylla (Peters 1864), Musa paradisiaca, Myiarchus crinitus, Myiarchus venezuelensis Lawrence 1865, Myiopagis gaimardii (Orbigny 1840), Myiozetetes similis (Spix 1825), Myospiza humeralis (Bosc 1792), Nemosia pileata (Boddaert 1783), Neurolaena lobata, Noctilio leporinus (Linnaeus 1758), Nycticorax nycticorax, Pachyramphus rufus (Boddaert 1783), Parkia velutina, Parkinsonia praecox, Passiflora foetida, Patagioenas corensis (Jacquin 1784), Paullinia cururu, Phaeomyias murina (Spix 1825), Phaseolus lunatus, Pheucticus ludovicianus, Phoradendron mucronatum, Phthirusa stelis, Phyla nodiflora, Piculus chrysochloros (Vieillot 1818), Picumnus cinnamomeus Wagler 1829, Pilherodius pileatus (Boddaert 1783), Pilosocereus lanuginosus, Pionus sordidus (Linnaeus 1758), Piptadenia viridiflora, Piranga flava (Vieillot 1822), Pitangus lictor (Lichtenstein 1823), Platalea ajaja Linnaeus 1758, Pleurodema brachyops, Polioptila plumbea (J. F. Gmelin 1788), Porophyllum ruderales, Pseudopaludicola pusilla, Pteronotus gymnotus (Natterer 1843), Pteronotus parnellii (Gray 1843), Pteronotus personatus (Wagner 1843), Pulsatrix perspicillata (Latham 1790), Pyrocephalus rubinus, Quiscalus lugubris, Rhizophora mangle, Rivina humilis, Rosenbergiodendron formosum, Ruellia paniculata, Ruprechtia ramiflora, Sakesphorus canadensis (Linnaeus 1766), Saltator coerulescens Vieillot 1817, Sarcostemma glaucum, Scoparia dulcis, Senna atomaria, Senna bicapsularis, Senna candolleana, Senna italica, Senna occidentalis, Senna tora, Serjania mexicana, Sesuvium portulacastrum, Sida ciliaris, Solandra grandiflora, Solanum bicolor, Spiza americana, Sporobolus coromandelianus, Sporobolus pyramidalis, Sporophila bouvronides (Lesson 1831), Sporophila intermedia Cabanis 1851, Sporophila nigricollis (Vieillot 1823), Stelgidopteryx ruficollis (Vieillot 1817), Stemodia maritima, Streptoprocne zonaris (Shaw 1796), Sturnella magna, Sublegatus arenarum (Salvin 1863), Synallaxis candei Orbigny & Lafresnaye 1838, Tabernaemontana amygdalifolia, Tachycineta albiventer (Boddaert 1783), Tachycineta bicolor (Vieillot 1808), Talinum paniculatum, Tapera naevia (Linnaeus 1766), Tephrosia cinerea, Thamnophilus doliatus (Linnaeus 1764), Thraupis glaucocolpa Cabanis 1850, Thraupis palmarum, Thryothorus leucotis (Lafresnaye 1845), Tiaris bicolor (Linnaeus 1766), Tiaris obscurus (Orbigny & Lafresnaye 1837), Tillandsia flexuosa, Tityra inquisitor (Lichtenstein 1823), Tolmomyias sulphurescens (Spix 1825), Tournefortia hirsutissima, Trianthema portulacastrum, Tribulus cistoides, Tyrannus dominicensis (Gmelin 1788), Tyto alba, Vireo flavoviridis (Cassin 1851), Vitex compressa, Wissadula periplocifolia, Zenaida auriculata

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

pages: 417

issue:

year: 2010

0: Cerrejón

title: Monitoreo de Fauna en el Valle del Cerrejón

source:

type: report

tags: Acacia glomerosa, Acacia tamarindifolia, Acacia tortuosa, Alternanthera halimifolia, Amazona ochrocephala, Ameiva festiva, Anas discors, Anemopaegma orbiculatum, Aninga aninga (Linnaeus 1766), Anoda crenatiflora, Aphelandra pulcherrima, Aramides cajanea, Aramus guarauna, Aratinga acuticaudata (Vieillot 1818), Ardea herodias, Aristolochia anguicida, Arremonops conirostris (Bonaparte 1850), Arremonops tocuyensis (Todd 1912), Artemisia vulgaris, Aspidosperma cuspa, Atalotriccus pilaris, Ayenia magna, Basileuterus flaveolus (S. F. Baird 1865), Bastardia viscosa, Batis maritima, Bauhinia pauletia, Boerhavia scandens, Bonellia frutescens, Bourreria cumanaensis, Bouteloua aristoides, Bromelia chrysantha, Bubo virginianus (Gmelin 1788), Burhinus bistriatus (Wagler 1829), Butorides virescens, Caesalpinia bonduc, Caesalpinia mollis, Calotropis gigantea, Campylorhynchus griseus (Swainson 1838), Campylorhynchus nuchalis (Cabanis 1847), Canavalia brasiliensis, Canavalia rosea, Capparis frondosa, Capparis linearis, Capparis pulcherrima, Caprimulgus cayennensis (Gmelin 1789), Capsicum baccatum, Cardinalis phoeniceus Bonaparte 1838, Cardiospermum corindum, Carduelis psaltria (Say 1823), Castela erecta, Celtis iguanaea, Cenchrus brownii, Cenchrus ciliaris, Cenchrus echinatus, Certhiaxis cinnamomeus (Gmelin

1788), *Chlorostilbon gibsoni* (Fraser 1840), *Chordeiles acutipennis* (Hermann 1783), *Chrysolampis mosquitus* (Linnaeus 1758), *Chrysomus icterocephalus*, *Cissus trifoliata*, *Citrullus lanatus*, *Cleome spinosa*, *Cnidoscopus urens*, *Coccoloba microstachya*, *Columbina minuta* (Linnaeus 1766), *Columbina passerina* (Linnaeus 1758), *Combretum fruticosum*, *Conirostrum leucogenys* (Lafresnaye 1852), *Conyza canadensis*, *Conyza perijaensis*, *Cordia macuirensis*, *Coryphospingus pileatus* (Wied-Neuwied 1821), *Craniolaria annua*, *Crotalaria purdiana*, *Crotalaria retusa*, *Croton ferrugineus*, *Croton malambo*, *Croton niveus*, *Croton ovalifolius*, *Croton punctatus*, *Crotophaga major*, *Crotophaga sulcirostris* Swainson 1827, *Cyanocorax affinis* (Pelzelin 1856), *Cyclarhis gujanensis* (J. F. Gmelin 1789), *Cyperus articulatus*, *Datura innoxia*, *Dendrocoryna autumnalis*, *Desmodium incanum*, *Diphysa carthagenensis*, *Ditaxis argothamnoides*, *Dolichandra unguis-cati*, *Dryocopus lineatus* (Linnaeus 1766), *Dysphania ambrosioides*, *Egletes prostrata*, *Egretta thula*, *Egretta tricolor* (Statius Muller 1776), *Elaenia chiriquensis* Lawrence 1865, *Eleocharis elegans*, *Eleocharis interstincta*, *Engystomops pustulosus*, *Eragrostis ciliaris*, *Erythrina berteriana*, *Euphorbia hirta*, *Evolvulus convolvuloides*, *Evolvulus sericeus*, *Exostema caribaeum*, *Fluvicola pica* (Boddaert 1783), *Forpus passerinus* (Linnaeus 1758), *Forpus xanthopterygius* (Spix 1824), *Fridericia chica*, *Fridericia dichotoma*, *Fridericia mollissima*, *Funastrum clausum*, *Furnarius leucopus* Swainson 1838, *Galactia striata*, *Galbula ruficauda* Cuvier 1816, *Glaucidium brasilianum* (Gmelin 1788), *Glossophaga longirostris* Miller 1898, *Gonatodes vittatus*, *Handroanthus serratifolius*, *Heliotropium angiospermum*, *Heliotropium indicum*, *Heliotropium procumbens*, *Hemitriccus margaritaceiventer* (Orbigny & Lafresnaye), *Hibiscus phoeniceus*, *Hillia costanensis*, *Himantopus mexicanus* (Statius Muller 1776), *Hippomane mancinella*, *Hylophilus flavipes* Lafresnaye 1845, *Hymenocallis caribaea*, *Hypnelus ruficollis* (Wagler 1829), *Icterus auricapillus* Cassin 1848, *Icterus galbula* (Linnaeus 1758), *Icterus nigrogularis* (Hahn 1819), *Inezia subflava* (P. L. Sclater & Salvin 1873), *Inezia tenuirostris* (Cory 1913), *Ipomoea incarnata*, *Jacquinia armillaris*, *Laguncularia racemosa*, *Lantana fucata*, *Lepidoblepharis sanctaemartae* Ruthven 1916, *Leptodactylus poecilochilus*, *Leptonycteris curasoae* Miller 1900, *Leptotila verreauxi* (Bonaparte 1855), *Licania cuspidata*, *Lonchocarpus atropurpureus*, *Lonchocarpus sanctae-marthae*, *Ludwigia octovalvis*, *Luffa operculata*, *Machetornis rixosa*, *Mallotus rhamnifolius*, *Manihot carthaginensis*, *Matelea maritima*, *Megaceryle alcyon* (Linnaeus 1758), *Megaceryle torquata* (Linnaeus 1766), *Megarynchus pitangua* (Linnaeus 1766), *Megascops choliba*, *Melanerpes rubricapillus*, *Melocactus intortus*, *Melochia parvifolia*, *Melochia tomentosa*, *Mentzelia aspera*, *Merremia aegyptia*, *Merremia umbellata*, *Micronycteris hirsuta* (Peters 1869), *Mimosa arenosa*, *Mimosa leiocarpa*, *Mimosa pellita*, *Mollugo verticillata*, *Molothrus bonariensis*, *Morisonia americana*, *Mormoops megalophylla* (Peters 1864), *Musa paradisiaca*, *Myiarchus crinitus*, *Myiarchus venezuelensis* Lawrence 1865, *Myiopagis gaimardii* (Orbigny 1840), *Myiozetetes similis* (Spix 1825), *Myospiza humeralis* (Bosc 1792), *Nemosia pileata* (Boddaert 1783), *Neurolaena lobata*, *Noctilio leporinus* (Linnaeus 1758), *Nycticorax nycticorax*, *Pachyramphus rufus* (Boddaert 1783), *Parkia velutina*, *Parkinsonia praecox*, *Passiflora foetida*, *Patagioenas corensis* (Jacquin 1784), *Paullinia cururu*, *Phaeomyias murina* (Spix 1825), *Phaseolus lunatus*, *Pheucticus ludovicianus*, *Phoradendron mucronatum*, *Phthirusa stelis*, *Phyla nodiflora*, *Piculus chrysochloros* (Vieillot 1818), *Picumnus cinnamomeus* Wagler 1829, *Pilherodius pileatus* (Boddaert 1783), *Pilosocereus lanuginosus*, *Pionus sordidus* (Linnaeus 1758), *Piptadenia viridiflora*, *Piranga flava* (Vieillot 1822), *Pitangus lictor* (Lichtenstein 1823), *Platalea ajaja* Linnaeus 1758, *Pleurodema brachyops*, *Polioptila plumbea* (J. F. Gmelin 1788), *Porophyllum ruderales*, *Pseudopaludicola pusilla*, *Pteronotus gymnotus* (Natterer 1843), *Pteronotus parnellii* (Gray 1843), *Pteronotus personatus* (Wagner 1843), *Pulsatrix perspicillata* (Latham 1790), *Pyrocephalus rubinus*, *Quiscalus lugubris*, *Rhizophora mangle*, *Rivina humilis*, *Rosenbergiodendron formosum*, *Ruellia paniculata*, *Ruprechtia ramiflora*, *Sakesphorus canadensis* (Linnaeus 1766), *Saltator coerulescens* Vieillot 1817, *Sarcostemma glaucum*, *Scoparia dulcis*, *Senna atomaria*, *Senna bicapsularis*, *Senna candolleana*, *Senna italica*, *Senna occidentalis*, *Senna tora*, *Serjania mexicana*, *Sesuvium portulacastrum*, *Sida ciliaris*, *Solandra grandiflora*, *Solanum bicolor*, *Spiza americana*, *Sporobolus coromandelianus*, *Sporobolus pyramidalis*, *Sporophila bouvronides* (Lesson 1831), *Sporophila intermedia* Cabanis 1851, *Sporophila nigricollis* (Vieillot 1823), *Stelgidopteryx ruficollis* (Vieillot 1817), *Stemodia maritima*, *Streptoprocne zonaris* (Shaw 1796), *Sturnella magna*, *Sublegatus arenarum* (Salvin 1863), *Synallaxis candei* Orbigny & Lafresnaye 1838, *Tabernaemontana amygdalifolia*, *Tachycineta albiventer* (Boddaert 1783), *Tachycineta bicolor* (Vieillot 1808), *Talinum paniculatum*, *Tapera naevia* (Linnaeus 1766), *Tephrosia cinerea*, *Thamnophilus doliatus* (Linnaeus 1764), *Thraupis glaucocolpa* Cabanis 1850, *Thraupis palmarum*, *Thryothorus leucotis* (Lafresnaye 1845), *Tiaris bicolor* (Linnaeus 1766), *Tiaris obscurus* (Orbigny & Lafresnaye 1837), *Tillandsia flexuosa*, *Tityra inquisitor* (Lichtenstein 1823), *Tolmomyias sulphurescens* (Spix 1825), *Tournefortia hirsutissima*, *Trianthema portulacastrum*, *Tribulus cistoides*, *Tyrannus dominicensis* (Gmelin 1788), *Tyto alba*, *Vireo flavoviridis* (Cassin 1851), *Vitex compressa*, *Wissadula periplocifolia*, *Zenaida auriculata*

abstract:

last_modified:

created:

group_id:
profile_id:
id:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Equipo Coordinador del SiB Colombia
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
habitatUnstructured: Local en pantanos, ciénagas, orillas cenagosas de ríos, playones de lodo y manglares (Hilty & Brown 1986). En La Guajira habita en ambientes acuáticos como ríos, lagunas, cuerpos de agua estacionales y bahías ribereñas (Cerrejón 2010).

endemicAtomized

distribution

ancillaryData

dataObject:
additionalInformation:

reference

keywords: Accipiter collaris, Accipiter cooperii, Accipiter poliogaster, Accipiter superciliosus, Amazona ochrocephala, Andigena hypoglaucha, Aramus guarauna, Aratinga pertinax, Ardea alba, Ardea cocoi, Ardea herodias, Arremonops conirostris (Bonaparte 1850), Arremonops tocuyensis (Todd 1912), Atalotriccus pilaris, Attila spadiceus (Gmelin 1789), Basileuterus flaveolus (S. F. Baird 1865), Bubo virginianus (Gmelin 1788), Busarellus nigricollis, Buteogallus anthracinus, Buteogallus meridionalis, Buteogallus schistaceus, Buteogallus solitarius, Buteogallus urubitinga, Cacicus cela, Cairina moschata, Campylorhamphus trochilirostris (Lichtenstein 1820), Campylorhynchus griseus (Swainson 1838), Campylorhynchus nuchalis (Cabanis 1847), Caprimulgus cayennensis (Gmelin 1789), Cardinalis phoeniceus Bonaparte 1838, Carduelis psaltria (Say 1823), Cathartes aura, Cathartes burrovianus, Cathartes melambrotus, Certhiaxis cinnamomeus (Gmelin 1788), Ceryle torquata, Chlorostilbon gibsoni (Fraser 1840), Chondrohierax uncinatus, Chordeiles acutipennis (Hermann 1783), Chrysomus icterocephalus, Circus buffoni, Circus cinereus, Circus cyaneus, Coccyzus pumila, Columbina minuta (Linnaeus 1766), Conirostrum leucogenys (Lafresnaye 1852), Coragyps atratus, Coryphospingus pileatus (Wied-Neuwied 1821), Crax alector, Crotophaga major, Crotophaga sulcirostris Swainson 1827, Cryptoleucopteryx plumbea, Cyanocorax affinis (Pelzelin 1856), Cyclarhis gujanensis (J. F. Gmelin 1789), Dryocopus lineatus (Linnaeus 1766), Elaenia chiriquensis Lawrence 1865, Elanoides forficatus, Elanus leucurus, Falco deiroleucus, Fluvicola pica (Boddaert 1783), Forpus passerinus (Linnaeus 1758), Forpus xanthopterygius (Spix 1824), Furnarius leucopus Swainson 1838, Galbula ruficauda Cuvier 1816, Gallinula chloropus (Linnaeus 1758), Gampsonyx swainsonii, Geranospiza caerulescens, Glaucidium brasilianum (Gmelin 1788), Harpagus bidentatus, Harpia harpyja, Helicolestes hamatus, Hemitriccus margaritaceiventer (Orbigny & Lafresna, Himantopus mexicanus (Statius Muller 1776), Hylophilus flavipes Lafresnaye 1845, Hypnelus ruficollis (Wagler 1829), Icterus auricapillus Cassin 1848, Icterus galbula (Linnaeus 1758), Icterus nigrogularis (Hahn 1819), Ictinia mississippiensis, Ictinia plumbea, Inezia subflava (P. L. Sclater & Salvin 1873), Inezia tenuirostris (Cory 1913), Jabiru mycteria, Jacana jacana, Lepidopyga goudoti (Bourcier 1843), Leptotila verreauxi (Bonaparte 1855), Leucopternis melanops, Leucopternis semiplumbea, Machetornis rixosa, Megaceryle alcyon (Linnaeus 1758), Megaceryle torquata (Linnaeus

1766), *Megarynchus pitangua* (Linnaeus 1766), *Megascops choliba*, *Melanerpes rubricapillus*, *Micrastur plumbeus*, *Mitu tomentosa*, *Molothrus bonariensis*, *Morphnarchus princeps*, *Morphnus guianensis*, *Myiarchus crinitus*, *Myiarchus venezuelensis* Lawrence 1865, *Myiopagis gaimardii* (Orbigny 1840), *Myiozetetes similis* (Spix 1825), *Myospiza humeralis* (Bosc 1792), *Nemosia pileata* (Boddaert 1783), *Neochen jubata*, *Opisthocomus hoazin*, *Oroaetus isidori*, *Ortalis ruficauda* Jardine 1847, *Pandion haliaetus*, *Parabuteo unicinctus*, *Patagioenas corensis* (Jacquin 1784), *Phaeomyias murina* (Spix 1825), *Pheucticus ludovicianus*, *Piculus chrysochloros* (Vieillot 1818), *Picumnus cinnamomeus* Wagler 1829, *Pilherodius pileatus* (Boddaert 1783), *Pionus sordidus* (Linnaeus 1758), *Pitangus lictor* (Lichtenstein 1823), *Pitangus sulphuratus*, *Polioptila plumbea* (J. F. Gmelin 1788), *Pseudastur albicollis*, *Pulsatrix perspicillata* (Latham 1790), *Pyrocephalus rubinus*, *Quiscalus lugubris*, *Quiscalus mexicanus* (Gmelin 1788), *Rostrhamus sociabilis*, *Sakesphorus canadensis* (Linnaeus 1766), *Saltator coerulescens* Vieillot 1817, *Sarcoramphus papa*, *Sarkidiornis melanotos* (Pennant 1769), *Spiza americana*, *Spizaetus tyrannus*, *Sporophila bouvronides* (Lesson 1831), *Sporophila intermedia* Cabanis 1851, *Sporophila nigricollis* (Vieillot 1823), *Stelgidopteryx ruficollis* (Vieillot 1817), *Sterna superciliaris*, *Sturnella magna*, *Sublegatus arenarum* (Salvin 1863), *Synallaxis candei* Orbigny & Lafresnaye 1838, *Tachycineta bicolor* (Vieillot 1808), *Tapera naevia* (Linnaeus 1766), *Thamnophilus doliatus* (Linnaeus 1764), *Thraupis glaucocolpa* Cabanis 1850, *Thraupis palmarum*, *Thryothorus leucotis* (Lafresnaye 1845), *Tyrannus dominicensis* (Gmelin 1788), *Tyto alba*, *Vireo flavoviridis* (Cassin 1851), *Zenaida auriculata*

editors

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

issue:

year: 1986

authors

authors: Hilty, S

authors: Brown, W. L.

title: A Guide to the Birds of Colombia

source:

type: book

tags: *Accipiter collaris*, *Accipiter cooperii*, *Accipiter poliogaster*, *Accipiter superciliosus*, *Amazona ochrocephala*, *Andigena hypoglaucha*, *Aramus guarauna*, *Aratinga pertinax*, *Ardea alba*, *Ardea cocoi*, *Ardea herodias*, *Arremonops conirostris* (Bonaparte 1850), *Arremonops tocuyensis* (Todd 1912), *Atalotriccus pilaris*, *Attila spadiceus* (Gmelin 1789), *Basileuterus flaveolus* (S. F. Baird 1865), *Bubo virginianus* (Gmelin 1788), *Busarellus nigricollis*, *Buteogallus anthracinus*, *Buteogallus meridionalis*, *Buteogallus schistaceus*, *Buteogallus solitarius*, *Buteogallus urubitinga*, *Cacicus cela*, *Cairina moschata*, *Campylorhamphus trochilirostris* (Lichtenstein 1820), *Campylorhynchus griseus* (Swainson 1838), *Campylorhynchus nuchalis* (Cabanis 1847), *Caprimulgus cayennensis* (Gmelin 1789), *Cardinalis phoeniceus* Bonaparte 1838, *Carduelis psaltria* (Say 1823), *Cathartes aura*, *Cathartes burrovianus*, *Cathartes melambrotus*, *Certhiaxis cinnamomeus* (Gmelin 1788), *Ceryle torquata*, *Chlorostilbon gibsoni* (Fraser 1840), *Chondrohierax uncinatus*, *Chordeiles acutipennis* (Hermann 1783), *Chrysomus icterocephalus*, *Circus buffoni*, *Circus cinereus*, *Circus cyaneus*, *Coccyzus pumila*, *Columbina minuta* (Linnaeus 1766), *Conirostrum leucogenys* (Lafresnaye 1852), *Coragyps atratus*, *Coryphospingus pileatus* (Wied-Neuwied 1821), *Crax alector*, *Crotophaga major*, *Crotophaga sulcirostris* Swainson 1827, *Cryptoleucopteryx plumbea*, *Cyanocorax affinis* (Pelzelin 1856), *Cyclarhis gujanensis* (J. F. Gmelin 1789), *Dryocopus lineatus* (Linnaeus 1766), *Elaenia chiriquensis* Lawrence 1865, *Elanoides forficatus*, *Elanus leucurus*, *Falco deiroleucus*, *Fluvicola pica* (Boddaert 1783), *Forpus passerinus* (Linnaeus 1758), *Forpus xanthopterygius* (Spix 1824), *Furnarius leucopus* Swainson 1838, *Galbula ruficauda* Cuvier 1816, *Gallinula chloropus* (Linnaeus 1758), *Gampsonyx swainsonii*, *Geranospiza caerulescens*, *Glaucidium brasilianum* (Gmelin 1788), *Harpagus bidentatus*, *Harpia harpyja*, *Helicolestes hamatus*, *Hemitriccus margaritaceiventer* (Orbigny & Lafresnaye), *Himantopus mexicanus* (Statius Muller 1776), *Hylophilus flavipes* Lafresnaye 1845, *Hypnelus ruficollis* (Wagler 1829), *Icterus auricapillus* Cassin 1848, *Icterus galbula* (Linnaeus 1758), *Icterus nigrogularis* (Hahn 1819), *Ictinia mississippiensis*, *Ictinia plumbea*, *Inezia*

subflava (P. L. Sclater & Salvin 1873), Inezia tenuirostris (Cory 1913), Jabiru mycteria, Jacana jacana, Lepidopyga goudoti (Bourcier 1843), Leptotila verreauxi (Bonaparte 1855), Leucopternis melanops, Leucopternis semiplumbea, Machetornis rixosa, Megaceryle alcyon (Linnaeus 1758), Megaceryle torquata (Linnaeus 1766), Megarynchus pitangua (Linnaeus 1766), Megascops choliba, Melanerpes rubricapillus, Micrastur plumbeus, Mitu tomentosa, Molothrus bonariensis, Morphnarchus princeps, Morphnus guianensis, Myiarchus crinitus, Myiarchus venezuelensis Lawrence 1865, Myiopagis gaimardii (Orbigny 1840), Myiozetetes similis (Spix 1825), Myospiza humeralis (Bosc 1792), Nemosia pileata (Boddaert 1783), Neochen jubata, Opisthocomus hoazin, Oroaetus isidori, Ortalis ruficauda Jardine 1847, Pandion haliaetus, Parabuteo unicinctus, Patagioenas corensis (Jacquin 1784), Phaeomyias murina (Spix 1825), Pheucticus ludovicianus, Piculus chrysochloros (Vieillot 1818), Picumnus cinnamomeus Wagler 1829, Pilherodius pileatus (Boddaert 1783), Pionus sordidus (Linnaeus 1758), Pitangus lictor (Lichtenstein 1823), Pitangus sulphuratus, Polioptila plumbea (J. F. Gmelin 1788), Pseudastur albicollis, Pulsatrix perspicillata (Latham 1790), Pyrocephalus rubinus, Quiscalus lugubris, Quiscalus mexicanus (Gmelin 1788), Rostrhamus sociabilis, Sakesphorus canadensis (Linnaeus 1766), Saltator coerulescens Vieillot 1817, Sarcoramphus papa, Sarkidiornis melanotos (Pennant 1769), Spiza americana, Spizaetus tyrannus, Sporophila bouvronides (Lesson 1831), Sporophila intermedia Cabanis 1851, Sporophila nigricollis (Vieillot 1823), Stelgidopteryx ruficollis (Vieillot 1817), Sterna supercilialis, Sturnella magna, Sublegatus arenarum (Salvin 1863), Synallaxis candei Orbigny & Lafresnaye 1838, Tachycineta bicolor (Vieillot 1808), Tapera naevia (Linnaeus 1766), Thamnophilus doliatus (Linnaeus 1764), Thraupis glaucocolpa Cabanis 1850, Thraupis palmarum, Thryothorus leucotis (Lafresnaye 1845), Tyrannus dominicensis (Gmelin 1788), Tyto alba, Vireo flavoviridis (Cassin 1851), Zenaida auriculata

abstract:

identifiers

last_modified:
created:
group_id:
profile_id:
id:
geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder:
rights: Equipo Coordinador del SiB Colombia
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

mimeType:
dataType:
identifier:
distributionUnstructured: Se encuentra en alturas hasta 400 m (Hilty & Brown 1986). En bases de datos y monitoreos de fauna se encontró que: Esta especie se distribuye en: Antioquia, Arauca, Atlántico, Bolívar, Caquetá, Cesar, La Guajira, Magdalena (Cerrejón 2010, Collections MNH 2012, Cerrejón 2010, ICN 2004). La subespecie Aramus guarauna guarauna se distribuye en: Antioquia, Arauca, Atlántico, Bolívar, Caquetá, Cesar, Magdalena (Collections MNH 2012, ICN 2004). Sus regiones naturales son la

Amazonía, Andina y Caribe de Colombia. A nivel mundial en bases de datos se encontró que:"W7F W7 V6–R 6R distribuye en: Brasil, Estados Unidos, México, (Arctos 2012)
Para las siguientes tres subespecies la distribución en el mundo es:
Aramus guarauna dolosus México (Arctos 2012), Aramus guarauna guarauna Colombia (Collections MNH 2012), Ecuador (Arctos 2012), Aramus guarauna pictus Estados Unidos (Arctos 2012)

distributionAtomized

stateProvince: Antioquia

ancillaryData

locality:
municipality:
country: Colombia

temporalCoverage

endDate:
startDate:

distributionScope

type:

threatStatus

ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:
locality:
municipality:
county:
stateProvince:
country: Global
0: IUCN
measurementValue: Preocupación menor (LC)
relatedTo:
measurementRemarks:
measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementType: IUCN
measurementID:

territory

territoryUnstructured:
areaOfOccupancy:
extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: Mide de 61 a 71 cm. Similar a un ibis, con alas anchas y redondeadas, cuello Delgado, patas largas y pico largo, ligeramente curvado. Principalmente pardo con unas pocas estrías blanquecinas e inconspicuas en la cabeza, cuello y la porción superior de la espalda; pico y patas oscuros (Hilty & Brown 1986).

[associatedParty](#)

role:

personnelIdentifier:

personnelDirectory:

homepage:

email: edgarestebanhc@gmail.com

phone:

postalCode:

country:

state:

city:

address:

organisation:

position:

lastName: Herrera Collazos

firstName: Edgar Esteban

ancillaryData

dataObject:
additionalInformation: Créditos específicos:
Ficha elaborada con el apoyo de Carbones del Cerrejón Limited.

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder:
rights: Equipo Coordinador del SiB Colombia
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

contentType:
dataType:
identifier:

_id
_bsontype: ObjectID
id: Vç—€fÄW Dä