

Title

Author

Author Affiliations

Course

Professor

Date

Title**Section****Subsection**

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tincidunt tristique, libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accumsan nec, suscipit a, ipsum. Morbi blandit ligula feugiat magna. Nunc eleifend consequat lorem. Sed lacinia nulla vitae enim. Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus. Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus eu enim. Vestibulum pellentesque felis eu massa.

Appendix

L^AT_EX Snippets

Citation & Reference Usage

- Parenthetical: This is a cited statement (Lu et al., 2023).
- Narrative: Lu et al. (2023) did research.
- Multiple: (Lu et al., 2023; Seiti et al., 2023) corroborate the findings.
- Figure: Refer to Figure A1.

Figure

Figure A1

Figure caption



Note. Adapted with permission.

Formatting

- **bold**
- *italic*
- typewriter
- underlined
- SMALL CAPS
- sans serif

Mathematics

Fubini’s theorem states that if a function $f(x, y)$ is continuous on a rectangular region $R = [a, b] \times [c, d]$, then the double integral can be evaluated as an iterated integral:

$$\iint_R f(x, y) \, dA = \int_a^b \left(\int_c^d f(x, y) \, dy \right) dx = \int_c^d \left(\int_a^b f(x, y) \, dx \right) dy$$

(A1)

Quote

This is a quote.

Table

Table A1

Table Caption

Variable	Condition A	Condition B
Value 1	0.00	0.00
Value 2	0.00	0.00

Note. Some notes.

References

- Lu, T., Ji, S., Jin, W., Yang, Q., Luo, Q., & Ren, T.-L. (2023). Biocompatible and long-term monitoring strategies of wearable, ingestible and implantable biosensors: Reform the next generation healthcare. *Sensors*, 23(6), 2991. <https://doi.org/10.3390/s23062991>
- Seiti, M., Giuri, A., Corcione, C. E., & Ferraris, E. (2023). Advancements in tailoring pedot: Pss properties for bioelectronic applications: A comprehensive review. *Biomaterials Advances*, 154, 213655. <https://doi.org/10.1016/j.bioadv.2023.213655>