

DATA 505 (Statistics Using R): Exam 1

Part 2: Open Computer

Name: Ryan Lazar

2025-09-30

Instructions

Welcome to Part 2 of the exam!

You now have access to your computer and any online resources that are “not alive” (i.e., you may freely use online resources, but you may not communicate with any other person). You have approximately 45 minutes to complete this part.

Fill in the code chunks below and provide written answers where requested. Submit this completed .qmd file on Moodle along with the rendered PDF.

Setup and Data Loading

We will analyze data about Antarctic penguins from the Palmer Station research program. The data are available in the `palmerpenguins` R package.

```
# Install and load the required packages
if(!"palmerpenguins" %in% installed.packages()){
  install.packages("palmerpenguins")
}
if(!"ggplot2" %in% installed.packages()){
  install.packages("ggplot2")
}

library(palmerpenguins)
library(ggplot2)

# Load the penguins data
data("penguins")
```

Data Exploration (4 points)

1 (2 points)

Use the `str()` function to examine the structure of the penguins dataset.

```
str(penguins)
```

```
tibble [344 x 8] (S3: tbl_df/tbl/data.frame)
 $ species      : Factor w/ 3 levels "Adelie","Chinstrap",...: 1 1 1 1 1 1 1 1 1 1 ...
 $ island       : Factor w/ 3 levels "Biscoe","Dream",...: 3 3 3 3 3 3 3 3 3 3 ...
 $ bill_length_mm : num [1:344] 39.1 39.5 40.3 NA 36.7 39.3 38.9 39.2 34.1 42 ...
 $ bill_depth_mm  : num [1:344] 18.7 17.4 18 NA 19.3 20.6 17.8 19.6 18.1 20.2 ...
 $ flipper_length_mm: int [1:344] 181 186 195 NA 193 190 181 195 193 190 ...
 $ body_mass_g    : int [1:344] 3750 3800 3250 NA 3450 3650 3625 4675 3475 4250 ...
 $ sex           : Factor w/ 2 levels "female","male": 2 1 1 NA 1 2 1 2 NA NA ...
 $ year          : int [1:344] 2007 2007 2007 2007 2007 2007 2007 2007 2007 2007 ...
```

How many observations and variables are there?

Answer: There are 344 observations, and 8 variables.

2 (2 points)

Extract the `bill_length_mm` column from the penguins dataset and store it in a new variable called `bill_lengths`. Then use the `summary()` function to examine this variable.

```
bill_lengths <- penguins$bill_length_mm
summary(bill_lengths)
```

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.	NA's
32.10	39.23	44.45	43.92	48.50	59.60	2

Research Question

We want to investigate whether the mean bill length of Adelie penguins differs significantly from 39 mm. This research question is motivated by previous studies suggesting that 39 mm might be a typical bill length for this species.

1 (4 points)

Filter the dataset to include only Adelie penguins and store the result in a new variable called `adelie_penguins`. Remove any rows with missing bill length measurements.

```
adelie_penguins <- penguins[penguins$species == "Adelie" & !is.na(penguins$bill_length_mm), ]
```

2 (3 points)

Extract the bill lengths from the Adelie penguin data and store them in a variable called `adelie_bills`. How many Adelie penguins have complete bill length measurements?

```
adelie_bills <- adelie_penguins$bill_length_mm  
length(adelie_bills)
```

```
[1] 151
```

Number of Adelie penguins with complete data: 151 entries after we dropped all Adelie penguins without complete data

3 (5 points)

Calculate the following descriptive statistics for Adelie penguin bill lengths: - Mean - Standard deviation
- Median - 25th and 75th percentiles

```
mean(adelie_bills)
```

```
[1] 38.79139
```

```
sd(adelie_bills)
```

```
[1] 2.663405
```

```
median(adelie_bills)
```

```
[1] 38.8
```


```
quantile(adelie_bills, c(.25,.75))
```

```
25% 75%  
36.75 40.75
```

4 (5 points)

Create a histogram of Adelie penguin bill lengths using ggplot2 or base R. Include appropriate axis labels and a title.

```
hist(adelie_penguins$bill_length_mm,  
     main = "Distribution of Bill Lengths in Adelie Penguins",  
     xlab = "Bill Length (mm)",  
     ylab = "Frequency",  
     col = "steelblue",  
     border = "black")
```



Based on the histogram, briefly describe the distribution of bill lengths (address shape, center, and spread):

Description: Based on the histogram, the distribution of Adelie penguin bill lengths appears approximately normal (bell-shaped) with a slight right skew, centered around 38-39 mm where

the frequency is highest. The bill lengths range from approximately 32 mm to 46 mm, showing a spread of about 14 mm, with most values concentrated between 36 mm and 42 mm. The distribution shows relatively low frequencies at both extremes, indicating that very short or very long bills are uncommon in this population.

Statistical Analysis

1 (5 points)

Now we will formally test whether the population mean bill length μ differs from 39 inches, i.e. we will test:

$$H_0 : \mu = 39$$

against

$$H_A : \mu \neq 39$$

Perform a two-sided t-test to test these hypotheses. Use $\alpha = 0.05$.

```
# Perform a two-sided t-test  
t.test(adelie_penguins$bill_length_mm, mu = 39, conf.level = 0.95)
```

One Sample t-test

```
data:  adelie_penguins$bill_length_mm  
t = -0.96246, df = 150, p-value = 0.3374  
alternative hypothesis: true mean is not equal to 39  
95 percent confidence interval:  
 38.36312 39.21966  
sample estimates:  
mean of x  
 38.79139
```

Based on your test results, what is your conclusion? Interpret the results in the context of the research question.

Conclusion: There is insufficient evidence at the 0.05 significance level to conclude that the population mean bill length of Adelie penguins differs from 39 mm. The sample mean of 38.79 mm is not statistically significantly different from 39 mm.

2 (4 points)

Define another variable (a numeric vector) `confidence_interval` and assign as its value the two bounds (lower and upper) of a 95% confidence interval for the mean bill length of Adelie penguins. Print the result. Full points are possible only if you do not “hard code” the bounds for the interval, i.e. if you create this vector directly by using R functions applied to the dataset. Half points maximum if you define the vector by typing the entries manually.

```
# Perform the t-test and store the results
test_result <- t.test(adelie_penguins$bill_length_mm, mu = 39)

# Extract the confidence interval bounds
confidence_interval <- test_result$conf.int

# Print the result
confidence_interval
```

```
[1] 38.36312 39.21966
attr("conf.level")
[1] 0.95
```
