

## CLASS 9, WEDNESDAY FEBRUARY 28TH: TENSOR PRODUCTS I

In Homework 2, I defined the tensor product of 2 modules  $M, N$ . This is one of the most powerful tools in the study of module theory, and today we will study some of the basic properties. Recall the definition:

**Definition 0.1.** The **tensor product** of two  $R$ -modules  $M, N$  is

$$M \otimes_R N = \left\{ \sum_{i=1}^l m_i \otimes n_i \mid m_i \in M, n_i \in N \right\} / \sim$$

where  $\sim$  is defined by

- 1)  $mr \otimes n \sim m \otimes rn$
- 2)  $m \otimes n + m' \otimes n \sim (m + m') \otimes n$
- 3)  $m \otimes n + m \otimes n' \sim m \otimes (n + n')$

In the homework, you are asked to prove the following 2 facts:  $M \otimes_R N$  has a natural  $R$ -module structure, and tensoring by a ring  $S$  which is also an  $R$ -module can upgrade  $M$  to an  $S$ -module (cf Homework 2, 5/6). These facts are invaluable. I now state some further basic properties:

**Proposition 0.2.** 1) If  $\phi : R \rightarrow S$  is a ring homomorphism, then there is a natural  $R$ -module homomorphism  $M \rightarrow M \otimes_R S$  given by  $m \mapsto m \otimes 1$ .  
2) There is a natural map  $\otimes : M \oplus N \rightarrow M \otimes_R N$  given by  $(m, n) \mapsto m \otimes n$ . This is not a homomorphism!  
3) If  $\varphi : M \oplus N \rightarrow P$  is a bilinear map, then  $\exists! \Phi : M \otimes_R N \rightarrow P$  such that  $\varphi = \Phi \circ \otimes$ .

*Proof.* I prove the statements in order:

- 1) If  $r \in R$ , then  $r\phi(m) = r(m \otimes 1) = (rm) \otimes 1 = \varphi(rm)$ . Similarly, relation 2) above implies  $\phi$  is additive. Therefore it is a homomorphism of  $R$ -modules.
- 2) There is almost nothing to prove. This is what is called a **balanced product**, and is key to the definition of the tensor product. Note that this is not a homomorphism of  $R$ -modules because

$$\begin{aligned} \otimes((m, n) + (m', n')) &= \otimes(m + m', n + n') = (m + m') \otimes (n + n') = \\ &= m \otimes n + m \otimes n' + m' \otimes n + m' \otimes n' \neq m \otimes n + m' \otimes n' \end{aligned}$$

- 3) First note that a map is called **bilinear** if restricting to a specific  $m \in M$  or  $n \in N$  makes  $\varphi$  into a homomorphism:

$$\begin{aligned} \varphi(rm + m', n) &= r\varphi(m, n) + \varphi(m', n) \\ \varphi(m, rn + n') &= r\varphi(m, n) + \varphi(m, n') \end{aligned}$$

Therefore, in part 2),  $\otimes$  is a natural bilinear map. As a result, we define

$$\Phi : M \oplus N \rightarrow P : \sum_i m_i \otimes n_i \mapsto \sum_i \varphi(m_i, n_i)$$

This forces  $\varphi = \Phi \circ \otimes$ , since

$$(\Phi \circ \otimes)(m, n) = \Phi(m \otimes n) = \varphi(m, n)$$

Moreover, this is a homomorphism as a result of the conditions of the definition of  $M \otimes_R N$ :

$$\Phi(rm \otimes n + m' \otimes n') = \Phi(rm \otimes n) + \Phi(m' \otimes n') = r\varphi(m, n) + \varphi(m', n')$$

□

**Note:** This gives an important way to convert a pair of homomorphism  $M \otimes P$  and  $N \otimes P$  into either a bilinear map  $M \oplus N \rightarrow P$  OR a single homomorphism  $M \otimes_R N \rightarrow P$ .

**Example 0.3** (Size of the Tensor Product). It is tempting to believe that the tensor product of two modules  $M, N$  is always bigger than  $M$  or  $N$  individually, much like the direct sum. However, the following nice proposition shows this is not the case:

**Proposition 0.4.** *Let  $R$  be a ring, and  $I$  be an ideal. Then*

$$M \otimes_R R/I \cong M/IM$$

*Proof.* I claim that the map is given by

$$\varphi : M \otimes_R R/I \rightarrow M/IM : m \otimes \bar{r} \mapsto \bar{r} \cdot m$$

This map is surjective, since if  $m + IM \in M/IM$ , we can consider  $m \otimes \bar{1} \in M \otimes_R R/I$ , whose image is the desired element. In addition, if  $\varphi(m \otimes \bar{r}) = \bar{0}$ , then  $rm \in IM$ . Therefore,  $rm = im'$  for  $i \in I$ . Therefore, the following equalities show that  $\varphi$  is injective:

$$m \otimes \bar{r} = rm \otimes \bar{1} = m' \otimes \bar{i} = m' \otimes 0 = 0$$

□

Another typical example is as follows: Consider  $R = \mathbb{Z}$ ;

$$\mathbb{Z}/n\mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}/m\mathbb{Z} \cong \mathbb{Z}/\gcd(m, n)\mathbb{Z} : a \otimes b \mapsto a \cdot b$$

$$\mathbb{Z}/n\mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Q} \cong 0$$

Another natural example we will use extensively is the localization of a module:

**Example 0.5** (Localization). We have already described  $W^{-1}R$ , the localization of  $R$  at  $W$ . This can be extended to modules in the following way: There is a natural localization map  $R \rightarrow W^{-1}R : r \mapsto (1, r)$ . Thus we can form the localization of an  $R$ -module:

$$W^{-1}M := M \otimes_R W^{-1}R$$

This naturally has the structure of a  $W^{-1}R$  module by the Homework 2 Exercise 6. It is also an  $R$ -module by virtue of the localization map.

It is useful to check that the localization of a module,  $W^{-1}M$ , is equivalent to that given by copying the conditions for a ring:

$$W^{-1}M = \{(w, m) : w \in W, m \in R\} / \sim$$

$(w, m) \sim (w', m')$  if there exists  $s \in W$  such that

$$s(wm' - w'm) = 0$$

We will see soon that a module locally satisfying some properties gives us ‘global’ information as well!