

Calculating the spawn index for Pacific Herring (*Clupea pallasii*) in British Columbia, Canada

Matthew H. Grinnell, Jake F. Schweigert, Matt
Thompson, and Jaclyn S. Cleary

Fisheries and Oceans Canada
Science Branch, Pacific Region
Pacific Biological Station
3190 Hammond Bay Road
Nanaimo, BC V9T 6N7

yyyy

Canadian Technical Report of
Fisheries and Aquatic Sciences nnnn



Fisheries and Oceans Pêches et Océans
Canada Canada

Canada 

Canadian Technical Report of Fisheries and Aquatic Sciences

Technical reports contain scientific and technical information that contributes to existing knowledge but which is not normally appropriate for primary literature. Technical reports are directed primarily toward a worldwide audience and have an international distribution. No restriction is placed on subject matter and the series reflects the broad interests and policies of Fisheries and Oceans Canada, namely, fisheries and aquatic sciences.

Technical reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report is abstracted in the data base *Aquatic Sciences and Fisheries Abstracts*.

Technical reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page. Out-of-stock reports will be supplied for a fee by commercial agents.

Numbers 1–456 in this series were issued as Technical Reports of the Fisheries Research Board of Canada. Numbers 457–714 were issued as Department of the Environment, Fisheries and Marine Service, Research and Development Directorate Technical Reports. Numbers 715–924 were issued as Department of Fisheries and Environment, Fisheries and Marine Service Technical Reports. The current series name was changed with report number 925.

Rapport technique canadien des sciences halieutiques et aquatiques

Les rapports techniques contiennent des renseignements scientifiques et techniques qui constituent une contribution aux connaissances actuelles, mais qui ne sont pas normalement appropriés pour la publication dans un journal scientifique. Les rapports techniques sont destinés essentiellement à un public international et ils sont distribués à cet échelon. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques de Pêches et Océans Canada, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports techniques peuvent être cités comme des publications à part entière. Le titre exact figure au-dessus du résumé de chaque rapport. Les rapports techniques sont résumés dans la base de données *Résumés des sciences aquatiques et halieutiques*.

Les rapports techniques sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre. Les rapports épuisés seront fournis contre rétribution par les agents commerciaux.

Les numéros 1 à 456 de cette série ont été publiés à titre de Rapports techniques de l'Office des recherches sur les pêcheries du Canada. Les numéros 457 à 714 sont parus à titre de Rapports techniques de la Direction générale de la recherche et du développement, Service des pêches et de la mer, ministère de l'Environnement. Les numéros 715 à 924 ont été publiés à titre de Rapports techniques du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 925.

Canadian Technical Report of
Fisheries and Aquatic Sciences nnnn

yyyy

CALCULATING THE SPAWN INDEX FOR PACIFIC HERRING
(*CLUPEA PALLASII*) IN BRITISH COLUMBIA, CANADA

by

Matthew H. Grinnell¹, Jake F. Schweigert², Matt Thompson¹, and Jaclyn
S. Cleary¹

¹ Pacific Biological Station, Fisheries and Oceans Canada,
3190 Hammond Bay Road, Nanaimo, British Columbia, V9T 6N7

² Emeritus, Pacific Biological Station, Fisheries and Oceans Canada,
3190 Hammond Bay Road, Nanaimo, British Columbia, V9T 6N7

© Her Majesty the Queen in Right of Canada, yyyy
Cat. No. Fs97-6/nnnn E ISBN xxx-x-xxx-xxxxx-x
Cat. No. Fs97-6/nnnn E-PDF ISSN 1488-5379

Correct citation for this publication:

Grinnell, M.H., Schweigert, J.F., Thompson, M., and Cleary, J.S. yyyy.
Calculating the spawn index for Pacific Herring (*Clupea pallasii*) in British
Columbia, Canada. Can. Tech. Rep. Fish. Aquat. Sci. nnnn: vii + 41 p.

CONTENTS

LIST OF FIGURES	v
LIST OF TABLES	v
ABSTRACT	vi
RÉSUMÉ	vii
1 INTRODUCTION	1
2 STATISTICAL FRAMEWORK	4
3 SAMPLING PROTOCOL	9
3.1 SURFACE SPAWN PROTOCOL	11
3.1.1 SPAWN INTENSITY CATEGORIES	12
3.2 MACROCYSTIS SPAWN PROTOCOL	13
3.3 UNDERSTORY SPAWN PROTOCOL	13
4 CONVERTING EGGS TO BIOMASS	14
5 SURFACE SPAWN CALCULATIONS	15
5.1 SAMPLE OBSERVATIONS AND CALCULATIONS	15
5.2 SPAWN OBSERVATIONS AND CALCULATIONS	17
6 MACROCYSTIS SPAWN CALCULATIONS	18
6.1 PLANT OBSERVATIONS	18
6.2 TRANSECT OBSERVATIONS AND CALCULATIONS	18
6.3 SPAWN OBSERVATIONS AND CALCULATIONS	18
7 UNDERSTORY SPAWN CALCULATIONS	21
7.1 QUADRAT OBSERVATIONS AND CALCULATIONS	21
7.2 TRANSECT OBSERVATIONS AND CALCULATIONS	26
7.3 SPAWN OBSERVATIONS AND CALCULATIONS	26
8 TOTAL SPAWN CALCULATIONS	27
9 SPAWN ON KELP CALCULATIONS	28

10 SOURCES OF UNCERTAINTY	29
11 FUTURE RESEARCH	31
12 CAVEATS	32
13 DOWNLOAD	32
14 ACKNOWLEDGEMENTS	33
15 REFERENCES	33
A SPAWN WIDTH ADJUSTMENTS	37
A.1 SURFACE SPAWN WIDTH	37
A.2 UNDERSTORY SPAWN WIDTH	38
B SURFACE SPAWN UPDATES	39

LIST OF FIGURES

1	Pacific Herring spawn index by Location, as well as Sections and Statistical Areas in the Strait of Georgia	3
2	Spatial boundaries for Pacific Herring stock assessment regions in British Columbia	4
3	Sequence of steps for Pacific Herring spawn index calculations	5
4	Aerial view of Pacific Herring spawn	11
5	Number of Pacific Herring egg layers	12
6	Egg density for Pacific Herring surface spawn surveys	17
7	Number of eggs per plant for Pacific Herring Macrocystis spawn surveys	22
8	Substrate egg density for Pacific Herring underwater spawn surveys	24
9	Algae egg density for Pacific Herring underwater spawn surveys	25

LIST OF TABLES

1	Index notation for Pacific Herring spawn index calculations . .	8
2	Notation for Pacific Herring spawn survey statistical framework	9
3	Spawn intensity categories and number of egg layers for Pacific Herring surface spawn surveys	14
4	Notation for converting Pacific Herring eggs to biomass	15
5	Notation for Pacific Herring surface spawn index calculations .	16
6	Notation for Pacific Herring Macrocystis spawn index calculations	19
7	Notation for Pacific Herring understory spawn index calculations	23
8	Algae types and coefficients for Pacific Herring understory spawn surveys	26
9	Notation for Pacific Herring total spawn index calculations . .	28
10	Notation for Pacific Herring spawn on kelp calculations	30
A.1	Spawn width correction factors for Pacific Herring understory spawn surveys	40
A.2	Notation for Pacific Herring understory spawn width adjustments	40

ABSTRACT

Grinnell, M.H., Schweigert, J.F., Thompson, M., and Cleary, J.S. yyyy.
Calculating the spawn index for Pacific Herring (*Clupea pallasii*) in British
Columbia, Canada. Can. Tech. Rep. Fish. Aquat. Sci. nnnn: vii + 41 p.

The spawn index time series is one component of Pacific Herring (*Clupea pallasii*) stock assessments in British Columbia, Canada. This document describes how we calculate the spawn index from spawn survey observations (e.g., spawn extent, number of egg layers, substrate type). There are three types of spawn survey observations: (1) observations of spawn taken from the surface usually at low tide; (2) underwater observations of spawn on giant kelp, *Macrocystis* (*Macrocystis* spp.); and (3) underwater observations of spawn on other types of algae and the substrate, which we refer to as ‘understory.’ We calculate the spawn index in four steps. First, we develop a sampling protocol to estimate the number of eggs in a given area. Second, we develop a conversion factor to convert Pacific Herring eggs to biomass, which is critical to calculating the spawn index. Third, we calculate the spawn index for each of the three aforementioned spawn survey types: surface, *Macrocystis*, and understory. Finally, we combine the three spawn indices, and aggregate by stock assessment region and year to produce a relative index of combined sex spawning biomass. In addition, we identify uncertainties in spawn index calculations, and we describe how users can install the **R** package to calculate the spawn index using an example database. Although we transform the spawn survey data from egg density to biomass in tonnes, the annual time series of egg density and biomass are relative indices of spawning biomass.

RÉSUMÉ

Grinnell, M.H., Schweigert, J.F., Thompson, M., and Cleary, J.S. yyyy.
Calculating the spawn index for Pacific Herring (*Clupea pallasii*) in British
Columbia, Canada. Can. Tech. Rep. Fish. Aquat. Sci. nnnn: vii + 41 p.

La série chronologique de l'indice de frai est une composante des évaluations des stocks de hareng du Pacifique (*Clupea pallasii*) en Colombie-Britannique, Canada. Ce document décrit comment nous calculons l'indice de frai à partir des observations du relevé du frai (par ex., l'étendue du frai, le nombre de couches d'œufs, le type de substrat). Il existe trois types d'observations du relevé des frayères: (1) les observations des frayères prélevées à la surface habituellement à marée basse; (2) les observations sous-marines des frayères sur varech géant, *Macrocystis* (*Macrocystis* spp.); et (3) les observations sous-marines des frayères sur les autres algues et le substrat, que nous appelons «sous-étage». Nous calculons l'indice de frai en quatre étapes. Premièrement, nous élaborons un protocole d'échantillonnage pour estimer le nombre d'œufs dans une zone donnée. Deuxièmement, nous élaborons un facteur de conversion pour convertir les œufs de hareng du Pacifique en biomasse, ce qui est essentiel pour calculer l'indice de reproduction. Troisièmement, nous calculons l'indice de frai pour chacun des trois types de relevés de frai susmentionnés: surface, *Macrocystis*, et sous-étage. Enfin, nous combinons les trois indices de frai, et les regroupons par région d'évaluation du stock et par année pour produire un indice relatif de la biomasse reproductrice combinée des sexes. De plus, nous identifions les incertitudes dans le calcul de l'indice de frai, et nous décrivons comment les utilisateurs peuvent installer le packet **R** pour calculer l'indice de frai à l'aide d'une base de données exemple. Bien que nous transformions les données du relevé de la densité des œufs en biomasse en tonnes, les séries chronologiques annuelles de la densité et de la biomasse des œufs sont des indices relatifs de la biomasse des géniteurs.

1 INTRODUCTION

Statistical age-structured stock assessment models rely on an indicator of relative population abundance to reconstruct a time series of estimated abundance. For Pacific Herring (*Clupea pallasii*) in British Columbia (BC), Canada, an index of relative population abundance is provided by monitoring the extent and intensity of spawn (i.e., egg) deposition throughout coastal BC (DFO 2019b). This document describes our calculations to convert spawn survey observations (e.g., spawn extent, number of egg layers, substrate type) to the spawn index for Pacific Herring in BC. These calculations have been described elsewhere, in either published or informal, internal documents. The objective of this document is to collate the various calculations in their order of application. Spawn index calculations have been updated over the years as more data and analyses justify improvements; we restrict this document to describing the current method.

Hart and Tester (1934) first demonstrated that an estimate of Pacific Herring abundance could be determined from a count of egg deposition in a small set of sampling quadrats. Based on their work, annual spawn surveys collect data used to calculate the spawn index. There are three types of spawn survey observations:

1. Observations of spawn taken from the surface usually at low tide,
2. Underwater observations of spawn on giant kelp, *Macrocystis* (*Macrocystis* spp.), and
3. Underwater observations of spawn on other types of algae and the substrate, which we refer to as ‘understory.’

Surface spawn surveys are believed to be the least accurate of the three survey types, but they have the greatest temporal and spatial extent (Schweigert 1993). For example, surface spawn surveys were the only survey type prior to 1988, and they are still used extensively for minor spawns, remote spawns (i.e., outside stock assessment region boundaries; see below), as well as unusually early or late spawns. *Macrocystis* and understory spawn surveys are conducted under water using SCUBA gear, and have been used for all major spawns since 1988. Thus, we describe the spawn index as having two periods based on the predominant survey type: the surface survey period from 1951 to 1987, and the dive survey period from 1988 to present.

The inclusion of dive surveys in 1988 makes it challenging to compare the

spawn index between these two periods. For example, surface surveys are less accurate than dive surveys (appendix A.1), and spawn surveys used subjective intensity categories instead of direct egg layer estimates until 1978 (subsubsection 3.1.1). In addition, Pacific Herring spawn survey effort has been inconsistent over the time series due to available resources and departmental priorities. For example, in the past, surveyors often dedicated several months each year to spawn surveys; they used small vessels to search for spawn, and surface surveys to estimate spawn biomass. Currently, surveyors use flights to search for spawn, and underwater SCUBA surveys to estimate spawn biomass. Thus, widespread effort (both spatially and temporally) in the past has been replaced with intense effort in the present. Pacific Herring spawn surveys began in 1928, but are considered incomplete for indexing purposes prior to 1937 because many potential areas were not surveyed (Hay and Kronlund 1987).

Pacific Herring spawn survey observations have a nested hierarchical structure: samples, *Macrocystis* plants, and quadrats are nested within transects, transects are nested within spawns, and spawns are nested within Locations. To develop spawn indices, Locations are nested within Sections (Figure 1a), Sections are nested within Statistical Areas (Figure 1b), and Statistical Areas are nested within stock assessment regions (SARs; Figure 1c). There are seven SARs in BC, which we categorize as either ‘major’ or ‘minor’ (Figure 2; Haist and Rosenfeld 1988). The terms ‘major’ and ‘minor’ describe relative differences in geographic and biomass scales. The major SARs are Haida Gwaii (formerly known as Queen Charlotte Islands), Prince Rupert District, Central Coast, Strait of Georgia, and West Coast of Vancouver Island. The minor SARs are Area 27 and Area 2 West; we do not develop spawn indices for minor SARs.

We calculate the spawn index in four steps. First, we develop a sampling protocol to estimate the number of eggs in a given area (section 2, section 3). Second, we develop a conversion factor to convert Pacific Herring eggs to biomass (section 4), which is critical to calculating the spawn index. Third, we calculate the spawn index for each of the three aforementioned spawn survey types: surface (section 5), *Macrocystis* (section 6), and understory (section 7). Within each section, we use subsections to separate levels of spatial aggregation (e.g., calculations at the quadrat, or transect level; Figure 3). Finally, we combine the three spawn indices, and aggregate by stock assessment region and year to produce a relative index of combined sex spawning biomass




Figure 1. Pacific Herring spawn index by Location in 2019 (a), Sections (b), and Statistical Areas (c) in the Strait of Georgia (SoG) stock assessment region (SAR; Figure 2).

73 (section 8).

74 We developed this document while converting spawn index calculations
 75 implemented in a **Microsoft Access** database to an **R** (RCT 2017) package,
 76 **SpawnIndex**. We updated the calculations from a database to an **R** package
 77 for several reasons. First, the database has been used for various purposes
 78 over the last two decades and has incidental calculations that make it overly
 79 complex. Second, the database is difficult to troubleshoot because it is hard
 80 to differentiate between input (i.e., data) and derived values. Third, the **R**
 81 package is open and transparent; researchers can view and download the
 82 package and an example spawn survey database. For example, users can
 83 run the **R** package functions to implement the calculations described in this
 84 document using a small set of actual observations. Fourth, we consider it good
 85 practice to separate data from analyses. Fifth, an **R** package will facilitate
 86 future research to quantify spawn index uncertainty. Finally, an **R** package
 87 will allow us to generate dynamic documents in the spirit of reproducible
 88 research using **knitr** (Xie 2015; Marwick et al. 2018). Essentially, we have
 89 attempted to follow ‘good enough’ practices in scientific computing (Wilson
 90 et al. 2016).




Figure 2. Spatial boundaries for Pacific Herring stock assessment regions (SARs) in British Columbia. There are five major SARs: Haida Gwaii (HG), Prince Rupert District (PRD), Central Coast (CC), Strait of Georgia (SoG), and West Coast of Vancouver Island (WCVI). There are two minor SARs: Area 27 (A27) and Area 2 West (A2W). Units: kilometres (km).

2 STATISTICAL FRAMEWORK

91

92 Historical and recent surface spawn surveys use an ad hoc sampling regimen,
 93 where surveys are often opportunistic given the state of the tide, as well as
 94 available sampling tools such as boats, rakes, and viewers. The data are
 95 analysed assuming simple random sampling, which likely generates a biased




Figure 3. Sequence of steps (e.g., S1) for Pacific Herring spawn index calculations for the three spawn survey types $x = \{\text{surface, Macrocystis, understory}\}$. Legend: rounded rectangles indicate start, rectangles indicate observations and calculations, parallelograms indicate output, arrows show order of operation, ‘§’ indicates section, and ‘Eq’ indicates equation.

96 estimate of mean egg density.

97 In contrast, underwater dive surveys using SCUBA gear instituted in
 98 1988 follow a two-stage systematic sampling design where transects are the
 99 first sampling stage, and individual quadrats within transects are the second
 100 sampling stage (Jessen 1978). Two steps are required to calculate mean
 101 understory egg density in each surveyed spawn s (Table 1). Note that we
 102 simplify index notation in this section by suppressing subscripts for spawn
 103 survey type x , Location n , SAR r , and year y . First, mean egg density for
 104 transect t in spawn s is

$$\overline{\rho}_{ts} = \frac{1}{Q} \sum_{q=1}^Q \rho_{qts} \quad (1)$$

105 where ρ_{qts} is egg density in thousands of eggs per square metre (m) for quadrat
 106 q (eggs $\cdot 10^3 \cdot \text{m}^{-2}$), Q is the number of quadrats in transect t (Table 2), and
 107 $\overline{\rho}_{ts}$ is in eggs $\cdot 10^3 \cdot \text{m}^{-2}$. Before we calculate the mean egg density for spawn
 108 s , we determine the mean number of potential quadrats in spawn s

$$\overline{Q}'_s = \frac{1}{T} \sum_{t=1}^T Q' \quad (2)$$

109 where Q' is the number of potential quadrats in transect t (i.e., a function of
 110 spawn width). Then, we calculate mean egg density in eggs per square metre
 111 for spawn s

$$\overline{\rho}_s = \frac{1}{T \overline{Q}'_s} \sum_{t=1}^T Q' \overline{\rho}_{ts} \quad (3)$$

112 where T is the number of transects in spawn s , \overline{Q}'_s is the mean number of
 113 potential quadrats in spawn s from Equation 2, Q' is the number of potential
 114 quadrats in transect t , $\overline{\rho}_{ts}$ is the mean transect egg density from Equation 1,
 115 and $\overline{\rho}_s$ is in eggs $\cdot 10^3 \cdot \text{m}^{-2}$. The egg density estimator from Equation 3 is
 116 unbiased, and the variance is

$$\sigma_s^2 = \frac{T' - T}{TT'} \sum_{t=1}^T \frac{(Q' \overline{\rho}_{ts} - \overline{Q}'_s \overline{\rho}_s)^2}{\overline{Q}'_s^2 (T - 1)} + \frac{f^t}{T^2} \sum_{t=1}^T \left(\frac{Q'}{\overline{Q}'_s} \right)^2 \frac{(1 - f^q) \sigma_{ts}^2}{Q} \quad (4)$$

117 where T' is the number of potential transects in spawn s (i.e., a function of
 118 spawn length), $\overline{\rho}_{ts}$ is the mean transect egg density from Equation 1, $\overline{\rho}_s$ is
 119 the mean spawn egg density from Equation 3, f^t is the transect sampling

120 fraction for spawn s

$$f^t = \frac{T}{T'}, \quad (5)$$

121 f^q is the quadrat sampling fraction for transect t

$$f^q = \frac{Q}{Q'}, \quad (6)$$

122 and σ_{ts}^2 is the within transect egg density variance

$$\sigma_{ts}^2 = \frac{1}{Q-1} \sum_{q=1}^Q (\rho_{qts} - \bar{\rho}_{ts})^2 \quad (7)$$

123 where ρ_{qts} is egg density for quadrat q , and $\bar{\rho}_{ts}$ is the mean transect egg
124 density from Equation 1.

125 The calculation of the mean egg density for each spawn requires estimates
126 of total spawn length, mean spawn width, length of each transect sampled,
127 and estimated egg density in each sampling quadrat. The protocol to optimize
128 sampling was determined through a series of studies conducted in the Strait
129 of Georgia in 1981 and 1983 (Schweigert et al. 1985, 1990), and on the West
130 Coast of Vancouver Island in 1982 (Schweigert et al. 1990). In the 1981
131 study, the location of transects and sampling quadrats along transects was
132 determined using random allocation (Schweigert et al. 1985). However, this
133 proved to be logistically difficult because neither the spawn length or width
134 is known a priori, and divers had difficulty making the necessary calculations
135 underwater. Nevertheless, data from these studies were used to determine a
136 sampling protocol to estimate mean egg density with a standard error of no
137 more than 25%. The results indicated that the sampling required to achieve
138 this level of precision included surveying three transects per kilometre of
139 spawn length, and sampling at least five quadrats per transect (i.e., spawn
140 width). The sampling design was tested during a 1983 survey in the Strait of
141 Georgia that applied a systematic rather than a random sampling protocol
142 to simplify the logistics; variance estimates were similar to those from the
143 1981 study. This sampling protocol was further re-evaluated after additional
144 surveys occurred in all areas of the coast during 1984 and 1985; the protocol
145 was found to be robust and has been in routine use since 1988 (Schweigert
146 et al. 1990). Although samples are collected systematically within each spawn,
147 we assume that transects and quadrats are located randomly with respect to

Table 1. Index notation for Pacific Herring spawn index calculations. Legend: stock assessment region (SAR), spawn on kelp (SOK).

Name	Description	Range
i	Substrate type	1, 2, 3, ..., I
I	Number of substrate types in sample j , type x , spawn s , Location n , SAR r , and year y	
j	Sample	1, 2, 3, ..., J
J	Number of samples in type x , spawn s , Location n , SAR r , and year y	
p	Plant	1, 2, 3, ..., P
P	Number of plants in transect t , type x , spawn s , Location n , SAR r , and year y	
t	Transect	1, 2, 3, ..., T
T	Number of transects in type x , spawn s , Location n , SAR r , and year y	
T'	Number of potential transects in type x , spawn s , Location n , SAR r , and year y	
v	Algae (aka vegetation) type	1, 2, 3, ..., V
V	Number of algae types in quadrat q , transect t , type x , spawn s , Location n , SAR r , and year y	
q	Quadrat	1, 2, 3, ..., Q
Q	Number of quadrats in transect t , type x , spawn s , Location n , SAR r , and year y	
Q'	Number of potential quadrats in transect t , type x , spawn s , Location n , SAR r , and year y	
x	Spawn survey type	1, 2, 3, ..., X
X	Number of spawn survey types in spawn s , location n , SAR r , and year y	
s	Spawn	1, 2, 3, ..., S
S	Number of spawns in Location n , SAR r , and year y	
n	Location	1, 2, 3, ..., N
N	Number of locations in SAR r and year y	
f	SOK fishery	1, 2, 3, ..., F
F	Number of SOK fisheries in SAR r and year y	
r	SAR	1, 2, 3, ..., R
R	Number of SARs	
y	Year	y_1, y_2, y_3, \dots, Y
Y	Last year of time series	

Table 2. Notation for Pacific Herring spawn survey statistical framework. Legend: metres (m).

Name	Description	Value or unit
ρ	Egg density	eggs $\cdot 10^3 \cdot \text{m}^{-2}$
$\bar{\rho}$	Mean egg density	eggs $\cdot 10^3 \cdot \text{m}^{-2}$
f^t	Transect sampling fraction	(0, 1]
f^q	Quadrat sampling fraction	(0, 1]
σ^2	Egg density variance	ρ^2

the underlying spawn distribution, and so these estimators are applicable. An analogous approach had previously been adopted in the sampling of various commercial fisheries where vessels arrive in port at random but are sampled in a systematic fashion to obtain a random sample (Quinn et al. 1983; Sen 1984).

Giant kelp, *Macrocystis* sp., requires a different sampling protocol than the aforementioned understory spawn survey protocol. Giant kelp routinely reach heights of 15 m, but once weighed down with herring eggs the plants can sink to lay flat on the bottom. After sampling dozens of giant kelp plants covered with herring eggs, it was determined that plant height, number of fronds per plant, and number egg layers per plant were key counts required to estimate the number of eggs per plant (Haegele and Schweigert 1990). The survey design employed to capture these data for each spawning bed rely on determining the average plant height, number of fronds in each plant holdfast, and number of giant kelp plants occurring within a 1 m swath on each side of the transect line. These data are used to determine the total egg deposition on *Macrocystis* sp. for each spawning bed (subsection 6.3).

3 SAMPLING PROTOCOL

The following is a brief summary of the spawn survey sampling protocol in the [Pacific Herring spawn survey manual](#). Pacific Herring in BC primarily spawn in sheltered bays and inlets, depositing their eggs on rocks and algae between depths of 1.5 m above and 18 m below the 0-tide level (Humphreys and Hourston 1978; Haegele and Schweigert 1985). We identify distinct spawns (both spatially and temporally) by a unique combi-

172 nation of year, Location, and ‘spawn number.’ Spawns are numbered
 173 $s = 1, 2, 3, \dots, S_{xnry}$ where S_{xnry} is the number of spawns in spawn sur-
 174 vey type x (i.e., $x = \{\text{surface, Macrocystis, understory}\}$), Location n , SAR
 175 r , and year y . A distinct spawn is a continuous stretch of shoreline with no
 176 detectable break in egg deposition; this is the finest scale at which we calculate
 177 the spawn index. A break in egg deposition is determined by the absence of
 178 Pacific Herring spawn on two consecutive transects, or by a temporal gap in
 179 spawning. Most spawns are also characterized by longitude and latitude, as
 180 well as start and end dates of spawning. Surveyors usually collect longitude
 181 and latitude at the start and end of each transect; for surface spawn surveys
 182 that don’t use transects (subsection 3.1), surveyors collect longitude and
 183 latitude at the start and ends of the spawn (i.e., overall length and width).

184 Pacific Herring spawns typically extend along the shore; from above,
 185 spawns are identified by a milky or turquoise discolouration of the ocean
 186 caused by the release of milt, and often appear as bands running parallel to
 187 the shore (Figure 4). Thus, spawn ‘length’ refers to distance parallel to the
 188 shore, and ‘width’ refers to distance perpendicular to the shore. Similarly,
 189 Macrocystis bed length L^m and algae (aka vegetation) bed length L^v refer
 190 to distances that Macrocystis and algae beds extend parallel to the shore,
 191 respectively.

192 Most areas of the BC coast have ‘permanent transect’ locations recorded
 193 on charts which enable surveyors to place transects in the same place each
 194 year. When permanent transects are unavailable for a given spawn, surveyors
 195 set new transects perpendicular to the shore, beginning 200 m in from one
 196 end of the spawn, and spaced 350 m apart along the length of the spawn.
 197 The end of the spawn is determined by the absence of eggs. We digitize
 198 new transects to make them available as permanent transects in subsequent
 199 surveys. Transects generally go from the deep edge of the spawn towards
 200 shore until divers reach the near-shore edge of the spawn; the near-shore edge
 201 can be out of the water depending on the tide height.

202 Pacific Herring spawn surveyors first determine the spatial extent of the
 203 spawn in terms of length of shoreline to survey (Figure 3, steps S1, M1, &
 204 U1); this is done by raking (subsection 3.1) or brief dives to determine the
 205 presence or absence of spawn. Surveyors place the first transect in from one
 206 end (i.e., at the first permanent transect, or 200 m if there are no permanent
 207 transects) to avoid surveying areas with patchy and sparse egg layers. Within
 208 the spawn area, surveyors use transects to determine spawn width, quadrat



Figure 4. Aerial view of Pacific Herring spawn taken during a spawn reconnaissance flight in the Strait of Georgia. The spawn is identified by the band of discoloured water parallel to the shore.

209 placement, and which *Macrocystis* plants to survey. In some cases, we adjust
210 spawn width to improve the accuracy of spawn index estimates (appendix A).

211 After determining the spatial extent, surveyors determine the number of
212 egg layers on substrate and algae according to sampling protocols described
213 in subsection 3.1, subsection 3.2, & subsection 3.3. For eggs on substrate,
214 one egg layer is a layer of eggs one egg thick over the entire spawned surface
215 (Figure 5a). For eggs on algae, surveyors count egg layers one of two ways
216 depending on whether the algae is flat or round in cross-section. Egg layers
217 on flat algae are counted on both sides of the algae (Figure 5b); egg layers on
218 round algae are counted across the diameter of the algae (Figure 5c).

219 **3.1 SURFACE SPAWN PROTOCOL**

220 Surface spawn surveyors use the aforementioned transect interval when pos-
221 sible, but the sampling interval relies on surveyor judgement and available
222 resources. If the spawn area is sufficiently large, surface surveyors usually
223 sample along permanent transects. Small spawns can still be mapped as they
224 were historically, with surveyors deciding how to sample the spawn. To sample,
225 surveyors deploy specialized rakes throughout the spawn to determine algae




Figure 5. Cross-sections showing the number of Pacific Herring egg layers on substrate (a), flat algae (b), and round algae (c). Diagrams copied with permission from the [Pacific Herring spawn survey manual](#).

226 (aka vegetation) type, number of egg layers, and percent cover. Surveyors
 227 may deploy a viewing box in shallow water, and at low tide a portion of the
 228 spawn may be visible for direct observation. We refer to these surface spawn
 229 observations as ‘samples.’

230 Recall that there are two cases of surface spawn surveys: all surveys prior
 231 to 1988, and surveys since 1988 when dive surveys are not possible. Data from
 232 surface surveys are combined with data from dive surveys (i.e., *Macrocystis*,
 233 understory) to produce the total spawn index (section 8).

234 3.1.1 SPAWN INTENSITY CATEGORIES

235 From 1928 to 1978, surface spawn surveyors categorized spawn by subjective
 236 ‘intensity’ categories instead of directly estimating the number of egg layers
 237 (Table 3). From 1928 to 1968 there were five intensity categories described as
 238 very light, light, medium, heavy, and very heavy (numbered 1 to 5, respec-
 239 tively). Starting in 1969 there were nine intensity categories; the change from
 240 five to nine intensity categories was probably to accommodate the practice
 241 of reporting intermediate categories such as 3.5 (Hay and Kronlund 1987).
 242 Starting in 1979, spawn surveyors estimated the number of egg layers directly,
 243 and they continued to record intensity categories until 1981 to provide overlap
 244 between the two methods. In addition to recording the number of egg layers,
 245 surveyors sometimes recorded intensity after it was officially discontinued in
 246 1981. We have converted spawn intensity observations in the Pacific Herring

spawn survey database from five to nine categories for spawns that used the five category scale between 1951 and 1968. Thus, spawn data used for stock assessments is represented either by a nine category intensity scale, or a direct estimate of the number of egg layers.

3.2 MACROCYSTIS SPAWN PROTOCOL

Macrocystis spawn surveyors take a census of Macrocystis plants within 1 m of the transect line, on both the left- and right-hand sides. We refer to the swath of substrate along Macrocystis transects as the transect swath, $\chi_{ts} = 2$ m in transect t and spawn s . Divers categorize Macrocystis plants as either ‘mature’ or ‘immature’ based on stipe height; mature plants have stipes ≥ 1 m high, and are the only plants used for Macrocystis spawn index calculations. Immature plants are excluded because Pacific Herring spawn on Macrocystis fronds, not stipes; immature plants have limited fronds and slimy stipes that prevent egg adhesion. In addition, Pacific Herring typically deposit spawn higher up Macrocystis plants. For each mature plant, divers record the number of stalks. For each transect, divers record the average number of egg layers, and average plant height. Haegele and Schweigert (1990) provide a description of the sampling technique, and the basis for estimating the total number of eggs per plant.

3.3 UNDERSTORY SPAWN PROTOCOL

Understory spawn surveyors place quadrats along transects, with a target frequency of ≥ 5 quadrats per transect, given a minimum spacing of 2 m and a maximum spacing of 40 m. Similar to how the first transect is moved in from one end of the spawn, the first quadrat is moved in from the edge of the spawn to the first 5 m mark on the transect line to avoid surveying areas with patchy and sparse egg layers. Note that transect line position along permanent transects varies year to year: spawn location causes transects to shift seaward or shoreward, and spawn width causes transects to be shorter or longer. Understory spawn surveys use 0.5 m^2 quadrats; other sizes (e.g., 0.25 and 1.0 m^2) have been used for research, but are not used to calculate the spawn index (Schweigert 1993). Within each quadrat, divers record the dominant (i.e., most heavily spawned) substrate type, percentage of the quadrat covered by spawn, and number of egg layers. In addition, divers

Table 3. Spawn intensity categories, description, and number of egg layers for Pacific Herring surface spawn surveys for the periods 1928 to 1968, and 1969 to 1978 (Hay and Kronlund 1987; Schweigert and Stocker 1988).

Intensity category		Description	Number of egg layers
1928 to 1968	1969 to 1978		
1	1	Very light	0.5529
	2		0.9444
2	3	Light	1.3360
	4		2.1496
3	5	Medium	2.9633
	6		4.1318
4	7	Heavy	5.3002
	8		6.5647
5	9	Very heavy	7.8291

identify the three most abundant algae types that have spawn. For each of these algae types, divers record the percentage of the quadrat covered by the algae and number of egg layers.

4 CONVERTING EGGS TO BIOMASS

After estimating the number of eggs in a spawn, the next step is to estimate the biomass of Pacific Herring that spawned. Female Pacific Herring produce an average of approximately 200 000 eggs per kilogram (kg) of total body weight (Hay 1985; Hay and Brett 1988). We assume that females account for 50% of spawners, and we convert eggs to tonnes (t) of spawners using

$$\theta = \omega \phi^f \frac{10^3 \text{ kg}}{\text{t}} \quad (8)$$

where ω is the number of eggs per kilogram of total female body weight, ϕ^f is the proportion of spawners that are female, and θ is the egg conversion factor (Table 4). Thus we convert eggs to biomass in tonnes of both sexes combined by dividing the number of eggs by θ . Although Pacific Herring egg production is affected by environmental variability and other factors (Tanasichuk and Ware 1987; Hay and Brett 1988), we assume that bias to the spawn index

from Equation 8 is insignificant in most areas and years (Schweigert 1993; Ware 1985).

5 SURFACE SPAWN CALCULATIONS

This section describes steps S2 to S4 in Figure 3. As in section 2, we simplify index notation in this section by suppressing subscripts for spawn survey type x , Location n , SAR r , and year y . Surface spawn surveyors sample along transects or using their judgement. Surveyors collect data for at the substrate type i , sample j , and spawn s levels; we calculate metrics at the substrate type i , sample j , and spawn s levels (Table 5). Recall that surface spawn ‘samples’ include observations collected using specialized rakes and viewing boxes (subsection 3.1). Occasionally, we use field data sheets to fill-in missing egg layer information for surface survey data (appendix B).

5.1 SAMPLE OBSERVATIONS AND CALCULATIONS

Each sample j (Figure 3, step S2) can have one or more substrate types i . The number of egg layers in substrate i , sample j , and spawn s is

$$E_{ijs} = E'_{ijs} \phi_{ijs}^b \quad (9)$$

where E'_{ijs} is the number of egg layers on substrate i , and ϕ_{ijs}^b is the proportion of substrate i covered with spawn. The total number of egg layers in sample

Table 4. Notation for converting the number of Pacific Herring eggs to biomass. Legend: kilograms (kg), tonnes (t).

Name	Description	Value or unit	Reference
ω	Female fecundity	200,000 eggs · kg ⁻¹	Hay (1985); Hay and Brett (1988)
ϕ^f	Proportion female	0.5	Hay (1985); Hay and Brett (1988)
θ	Egg conversion factor	eggs · 10 ⁸ · t ⁻¹	

Table 5. Notation for Pacific Herring surface spawn index calculations. Legend: metres (m), tonnes (t).

Name	Description	Value or unit	Reference
E'	Number of egg layers	> 0	
ϕ^b	Proportion of substrate covered in eggs	$(0, 1]$	
E	Number of egg layers	> 0	
α	Regression intercept	$14.698 \text{ eggs} \cdot 10^3 \cdot \text{m}^{-2}$	Schweigert et al. (1997)
β	Regression slope	$212.218 \text{ eggs} \cdot 10^3 \cdot \text{m}^{-2}$	Schweigert et al. (1997)
ρ	Egg density	$\text{eggs} \cdot 10^3 \cdot \text{m}^{-2}$	
$\bar{\rho}$	Mean egg density	$\text{eggs} \cdot 10^3 \cdot \text{m}^{-2}$	
L	Spawn length	m	
W	Spawn width	m	
B	Surface spawn index (i.e., biomass)	t	

312 j and spawn s is

$$E_{js} = \sum_{i=1}^I E_{ijs} \quad (10)$$

313 where E_{ijs} is the number of egg layers in substrate i from Equation 9. For
314 the time period when surveyors recorded spawn ‘intensity’ categories instead
315 of direct egg layer estimates, we convert intensity to number of egg layers E_{js}
316 (Table 3). Schweigert et al. (1997) developed a predictive model of surface
317 egg density as a function of number of egg layers using a linear regression
318 model¹

$$\rho_{js} = \alpha + \beta E_{js} \quad (11)$$

319 where α is the regression intercept, β is the regression slope, E_{js} is the total
320 number of egg layers in sample j from Equation 10, and ρ_{js} is in $\text{eggs} \cdot 10^3 \cdot \text{m}^{-2}$
321 (Figure 6). Note that we only calculate ρ_{js} if $E_{js} > 0$.

¹There is an error in Schweigert et al. (1997); surface egg density is in thousands per square metre ($\text{eggs} \cdot 10^3 \cdot \text{m}^{-2}$). Likewise, we report eggs in thousands (i.e., $\text{eggs} \cdot 10^3$) in this document and in the **R** package script.




Figure 6. Egg density in thousands of eggs per square metre (m) as a function of number of egg layers for Pacific Herring surface spawn surveys (Equation 11; Schweigert et al. 1997). Note that number of egg layers can exceed those shown in this figure; values shown are for demonstration only.

322 5.2 SPAWN OBSERVATIONS AND CALCULATIONS

323 The mean egg density in spawn s (Figure 3, step S3) is

$$\bar{\rho}_s = \frac{1}{J} \sum_{j=1}^J \rho_{js} \quad (12)$$

324 where ρ_{js} is egg density in sample j from Equation 11, J is the number of
 325 samples, and $\bar{\rho}_s$ is in eggs · 10³ · m⁻². Two other metrics are required at the
 326 spawn level: spawn length L_s , and spawn width W_s , both in metres. We set
 327 W_s to the first non-missing value of median pool width, median Section width,
 328 median SAR width, or observed width W' (in that order; subsection A.1).
 329 The surface spawn index in spawn s is

$$B_s = \frac{\bar{\rho}_s L_s W_s 10^3}{\theta} \quad (13)$$

330 where $\bar{\rho}_s$ is mean egg density in spawn s from Equation 12, L_s is length
 331 of spawn s , W_s is width of spawn s , θ is the egg conversion factor from
 332 Equation 8, and B_s is in tonnes (Figure 3, step S4).

6 MACROCYSTIS SPAWN CALCULATIONS

This section describes steps M2 to M5 in Figure 3. As with the previous section, we simplify index notation in this section by suppressing subscripts for spawn survey type x , Location n , SAR r , and year y . Macrocystis spawn surveyors use SCUBA gear to collect underwater data for individual plants p , transects t , and spawns s ; we calculate metrics at the transect t , and spawn s levels (Table 6).

6.1 PLANT OBSERVATIONS

For each mature plant p in transect t and spawn s (Figure 3, step M2), surveyors count the number of stalks K_{pts} .

6.2 TRANSECT OBSERVATIONS AND CALCULATIONS

At the transect t level (Figure 3, step M3), spawn width is W_{ts} , and transect swath is $\chi_{ts} = 2$ m, both in metres. We calculate the area in transect t and spawn s as

$$A_{ts} = W_{ts}\chi_{ts} \quad (14)$$

in square metres. In addition, divers estimate summary statistics for mature Macrocystis plants along transect t in spawn s : mean height \overline{H}_{ts} in metres, and mean number of egg layers \overline{E}_{ts} . The total number of plants in transect t and spawn s is P . We calculate the total number of stalks in transect t and spawn s

$$K_{ts} = \sum_{p=1}^P K_{pts} \quad (15)$$

where K_{pts} is the number of stalks on plant p .

6.3 SPAWN OBSERVATIONS AND CALCULATIONS

At the spawn s level (Figure 3, step M4), we determine the length of the Macrocystis bed L_s^m in metres. If L_s^m is inadvertently not recorded, we set L_s^m to the spawn length L_s . The mean width of spawn s is

$$\overline{W}_s = \frac{1}{T} \sum_{t=1}^T W_{ts} \quad (16)$$

Table 6. Notation for Pacific Herring *Macrocystis* spawn index calculations. Legend: metres (m), tonnes (t).

Name	Description	Value or unit	Reference
K	Number of stalks	> 0	
W	Spawn width	m	
χ	Transect swath	m	
A	Area	m^2	
\bar{W}	Mean spawn width	m	
\bar{H}	Mean plant height	m	
\bar{E}	Mean number of egg layers	> 0	
L^m	Length of the <i>Macrocystis</i> bed	m	
L	Spawn length	m	
P	Number of plants	> 0	
$\bar{\kappa}$	Mean number of stalks per plant	> 0	
β	Regression slope	$0.073 \text{ eggs} \cdot 10^3 \cdot \text{plant}^{-1}$	Haegle and Schweigert (1990)
γ	Regression exponent on \bar{E}	0.673	Haegle and Schweigert (1990)
δ	Regression exponent on \bar{H}	0.932	Haegle and Schweigert (1990)
ϵ	Regression exponent on $\bar{\kappa}$	0.703	Haegle and Schweigert (1990)
$\bar{\psi}$	Mean number of eggs per plant	$\text{eggs} \cdot 10^3 \cdot \text{plant}^{-1}$	
$\bar{\rho}$	Mean egg density	$\text{eggs} \cdot 10^3 \cdot \text{m}^{-2}$	
B	<i>Macrocystis</i> spawn index (i.e., biomass)	t	

where W_{ts} is the spawn width at transect t , T is the number of transects in spawn s , and \bar{W}_s is in metres. The total area of transects in spawn s is

$$A_s = \sum_{t=1}^T A_{ts} \quad (17)$$

359 where A_{ts} is the transect area from Equation 14, and A_s is in square metres.
 360 The total number of stalks in spawn s is

$$K_s = \sum_{t=1}^T K_{ts} \quad (18)$$

361 where K_{ts} is the number of stalks in transect t from Equation 15. The total
 362 number of plants in spawn s is

$$P_s = \sum_{t=1}^T P_t \quad (19)$$

363 where P_t is the number of plants in transect t . The mean plant height in
 364 spawn s is

$$\overline{H}_s = \frac{1}{T} \sum_{t=1}^T \overline{H}_{ts} \quad (20)$$

365 where \overline{H}_{ts} is the mean plant height in transect t , T is the number of transects
 366 in spawn s , and \overline{H}_s is in metres. The mean number of egg layers in spawn s is

$$\overline{E}_s = \frac{1}{T} \sum_{t=1}^T \overline{E}_{ts} \quad (21)$$

367 where \overline{E}_{ts} is the mean number of egg layers in transect t , and T is the number
 368 of transects in spawn s . The mean number of stalks per plant in spawn s is

$$\overline{\kappa}_s = \frac{K_s}{P_s} \quad (22)$$

369 where K_s is the number of stalks in spawn s from Equation 18, and P_s is the
 370 number of plants in spawn s from Equation 19.

371 Haegle and Schweigert (1990) developed a predictive model of number of
 372 eggs per plant as a function of number of egg layers on plants, plant height,
 373 and number of stalks per plant using a nonlinear multiple regression model

$$\overline{\psi}_s = \beta \overline{E}_s^\gamma \overline{H}_s^\delta \overline{\kappa}_s^\epsilon 10^3 \quad (23)$$

374 where β is the regression slope, \overline{E}_s is the mean number of egg layers in spawn
 375 s from Equation 21, γ is the regression exponent on \overline{E}_s , \overline{H}_s is the mean plant
 376 height in spawn s from Equation 20, δ is the regression exponent on \overline{H}_s , $\overline{\kappa}_s$ is

the mean number of stalks per plant in spawn s from Equation 22, ϵ is the regression exponent on $\bar{\kappa}_s$, and $\bar{\psi}_s$ is in eggs $\cdot 10^3 \cdot \text{plant}^{-1}$ (Figure 7). Mean macrocystis egg density in spawn s is

$$\bar{\rho}_s = \frac{\bar{\psi}_s P_s}{A_s} \quad (24)$$

where $\bar{\psi}_s$ is the mean number of eggs per plant in spawn s from Equation 23, P_s is the number of plants in spawn s from Equation 19, A_s is the total area of transects in spawn s from Equation 17, and $\bar{\rho}_s$ is in eggs $\cdot 10^3 \cdot \text{m}^{-2}$. The Macrocystis spawn index in spawn s is

$$B_s = \frac{\bar{\rho}_s L_s^m \bar{W}_s 10^3}{\theta} \quad (25)$$

where $\bar{\rho}_s$ is the mean egg density in spawn s from Equation 24, L_s^m is the length of the Macrocystis bed in spawn s , \bar{W}_s is the mean width of spawn s from Equation 16, θ is the egg conversion factor from Equation 8, and B_s is in tonnes (Figure 3, step M5).

7 UNDERSTORY SPAWN CALCULATIONS

This section describes steps U2 to U5 in Figure 3. As with the previous two sections, we simplify index notation in this section by suppressing subscripts for spawn survey type x , Location n , SAR r , and year y . Understory spawn surveyors use SCUBA gear to collect underwater data for algae (aka vegetation) types v , quadrats q , transects t , and spawns s ; we calculate metrics at the algae type v , quadrat q , transect t , and spawn s levels (Table 7).

7.1 QUADRAT OBSERVATIONS AND CALCULATIONS

We calculate two separate estimates of egg density at the quadrat level (Figure 3, step U2): spawn on substrate, and spawn on algae v . Haegele et al. (1979) developed a predictive model of substrate egg density in quadrat q , transect t , and spawn s from egg layers using a linear regression model

$$\rho_{qts}^b = \alpha E_{qts}^b \phi_{qts}^b \quad (26)$$

where α is the regression slope, E_{qts}^b is the number of egg layers on substrate in quadrat q , ϕ_{qts}^b is the proportion of substrate in quadrat q covered by spawn, and ρ_{qts}^b is substrate egg density in eggs $\cdot 10^3 \cdot \text{m}^{-2}$ (Figure 8).




Figure 7. Number of eggs in thousands per *Macrocystis* plant as a function of number of egg layers on plants, plant height in metres (m), and number of stalks per plant for Pacific Herring *Macrocystis* spawn surveys (Equation 23; Haegele and Schweigert 1990). Note that number of egg layers, plant height, and number of stalks per plant can exceed those shown in this figure; values shown are for demonstration only.

403 Although quadrats have only one substrate type, they can have up to
 404 three algae types v (subsection 3.3). Schweigert (2005) developed a predictive
 405 model of algae egg density from egg layers, proportion of the quadrat covered
 406 by algae, and an algae coefficient using a generalized linear model. Algae
 407 coefficients account for the effect of algae morphology on Pacific Herring egg

Table 7. Notation for Pacific Herring understory spawn index calculations. Legend: metres (m), tonnes (t).

Name	Description	Value or unit	Reference
E^b	Number of substrate egg layers	> 0	
ϕ^b	Proportion of substrate covered in eggs	$(0, 1]$	
α	Regression slope for substrate	$340 \text{ eggs} \cdot 10^3 \cdot \text{m}^{-2}$	Haegele et al. (1979)
ρ^b	Substrate egg density	$\text{eggs} \cdot 10^3 \cdot \text{m}^{-2}$	
β	Regression slope for algae	$600.567 \text{ eggs} \cdot 10^3 \cdot \text{m}^{-2}$	Schweigert (2005)
E^v	Number of algae egg layers	> 0	
γ	Regression exponent on E^v	0.6355	Schweigert (2005)
ϕ^v	Proportion of algae covered in eggs	$(0, 1]$	
δ	Regression exponent on ϕ^v	1.413	Schweigert (2005)
C	Algae coefficient	see Table 8	
ρ^v	Algae egg density	$\text{eggs} \cdot 10^3 \cdot \text{m}^{-2}$	
ρ	Egg density	$\text{eggs} \cdot 10^3 \cdot \text{m}^{-2}$	
W	Spawn width	m	
$\bar{\rho}$	Mean egg density	$\text{eggs} \cdot 10^3 \cdot \text{m}^{-2}$	
\bar{W}	Mean spawn width	m	
L^v	Length of the algae bed	m	
L	Spawn length	m	
B	Understory spawn index (i.e., biomass)	t	

density (Table 8). Egg density on algae v in quadrat q , transect t , and spawn s (Schweigert 2005) is

$$\rho_{vqts}^v = \beta E_{vqts}^{v\gamma} \phi_{vqts}^{v\delta} C_v \quad (27)$$

where β is the regression slope, E_{vqts}^v is the number of egg layers on algae v , γ is the regression exponent on E_{vqts}^v , ϕ_{vqts}^v is the proportion of quadrat q




Figure 8. Substrate egg density in thousands of eggs per square metre (m) as a function of number of egg layers on substrate and proportion of substrate covered in spawn for Pacific Herring underwater spawn surveys (Equation 26; Haegele et al. 1979). Note that number of egg layers can exceed those shown in this figure; values shown are for demonstration only.

412 covered by algae v , δ is the regression exponent on ϕ_{vqts}^v , C_v is the coefficient
 413 for algae v , and ρ_{vqts}^v is in eggs $\cdot 10^3 \cdot \text{m}^{-2}$ (Figure 9). The total algae egg
 414 density for quadrat q in transect t and spawn s is

$$\rho_{qts}^v = \sum_{v=1}^V \rho_{vqts}^v \quad (28)$$

415 where ρ_{vqts}^v is egg density on algae v from Equation 27, and ρ_{qts}^v is in eggs \cdot
 416 $10^3 \cdot \text{m}^{-2}$.

417 The total understory egg density for quadrat q in transect t and spawn s
 418 is

$$\rho_{qts} = \rho_{qts}^b + \rho_{qts}^v \quad (29)$$

419 where ρ_{qts}^b is substrate egg density from Equation 26, ρ_{qts}^v is algae egg density
 420 from Equation 28, and ρ_{qts} is in eggs $\cdot 10^3 \cdot \text{m}^{-2}$. Thus, we assume that eggs
 421 on substrate and algae are independent, and can be added without bias.




Figure 9. Algae egg density in thousands of eggs per square metre (m) as a function of number of egg layers on algae, proportion of substrate covered in algae, and algae coefficient (Table 8) for Pacific Herring underwater spawn surveys (Equation 27; Schweigert 2005). Note that number of egg layers can exceed those shown in this figure; values shown are for demonstration only.

Table 8. Algae (aka vegetation) types v and coefficients C for Pacific Herring understory spawn surveys (Schweigert 2005).

Algae type v	Coefficient C
Grasses	0.9715
Grunge	1.0000
Kelp (flat)	0.9119
Kelp (standing)	1.1766
Leafy algae	0.6553
Rockweed	0.7793
Sargassum	1.1766
Stringy algae	1.0000

7.2 TRANSECT OBSERVATIONS AND CALCULATIONS

At the transect level (Figure 3, step U3), the mean understory egg density in transect t and spawn s is

$$\overline{\rho}_{ts} = \frac{1}{Q} \sum_{q=1}^Q \rho_{qts} \quad (30)$$

where Q is the number of quadrats in transect t , ρ_{qts} is total understory egg density in quadrat q from Equation 29, and $\overline{\rho}_{ts}$ is in $\text{eggs} \cdot 10^3 \cdot \text{m}^{-2}$. Note that we update spawn width to correct for errors regarding the assumed accuracy of transect lines used to measure spawn width for understory surveys between 2003 and 2014 (subsection A.2).

7.3 SPAWN OBSERVATIONS AND CALCULATIONS

At the spawn level (Figure 3, step U4), the mean width of spawn s is

$$\overline{W}_s = \frac{1}{T} \sum_{t=1}^T W_{ts} \quad (31)$$

where W_{ts} is the spawn width at transect t , T is the number of transects in spawn s , and \overline{W}_s is in metres. The length of the algae bed in spawn s is L_s^v , also in metres. As with *Macrocystis* spawn calculations, if L_s^v is inadvertently not recorded, we set L_s^v to the spawn length L_s . Thus, we assume that eggs on

436 substrate and eggs on algae are represented by the same length measurement.
 437 Next, we calculate the weighted mean egg density in spawn s , where transect
 438 egg density is weighted by spawn width at transect t , W_{ts} . We calculate a
 439 weighted mean because spawn width varies along the spawn length; a weighted
 440 mean ensures that transects contribute proportionally to their area. The
 441 mean egg density in spawn s is

$$\bar{\rho}_s = \frac{\sum_{t=1}^T \bar{\rho}_{ts} W_{ts}}{\sum_{t=1}^T W_{ts}} \quad (32)$$

442 where $\bar{\rho}_{ts}$ is the mean understory egg density in transect t from Equation 30,
 443 W_{ts} is the spawn width for transect t in metres, and $\bar{\rho}_s$ is in eggs $\cdot 10^3 \cdot \text{m}^{-2}$.
 444 The understory spawn index in spawn s is

$$B_s = \frac{\bar{\rho}_s L_s^v \bar{W}_s 10^3}{\theta} \quad (33)$$

445 where $\bar{\rho}_s$ is the mean understory egg density from Equation 32, L_s^v is the
 446 length of the algae bed, \bar{W}_s is the mean spawn width from Equation 31, θ
 447 is the egg conversion factor from Equation 8, and B_s is in tonnes (Figure 3,
 448 step U5).

8 TOTAL SPAWN CALCULATIONS

449
 450 This section describes step T1 in Figure 3. Unlike the previous three sections,
 451 we include subscripts for spawn survey type x , Location n , SAR r , and year
 452 y in the equations in this section (Table 9). The total spawn index in spawn
 453 s , Location n , region r , and year y is

$$B_{snry} = \sum_{x=1}^X B_{xsnry} \quad (34)$$

454 where B_{xsnry} is spawn index for surface, Macrocystis, and understory spawn
 455 surveys from Equation 13, Equation 25, and Equation 33, respectively, and
 456 B_{snry} is in tonnes (Figure 3, step T1). Finally, we aggregate the total spawn
 457 index by SAR r and year y

$$B_{ry} = \sum_{n=1}^N \sum_{s=1}^S B_{snry} \quad (35)$$

where B_{snry} is the total spawn index from Equation 34, and B_{ry} is a relative index of combined sex spawning biomass for SAR r and year y in tonnes. We use B_{ry} as an indicator of Pacific Herring relative population abundance (i.e., biomass) in stock assessment models.

9 SPAWN ON KELP CALCULATIONS

Spawn on kelp (SOK) fisheries collect Pacific Herring roe that adhere to algae such as *Macrocystis* after spawning. Other similar fisheries include spawn on bough, in which operators collect roe that adhere to tree boughs; we refer to these fisheries collectively as SOK in this document. There are two types of SOK fisheries in BC: ‘open-pond’ in which operators provide algae to spawning Pacific Herring, and ‘closed-pond’ in which operators impound spawning Pacific Herring in floating nets that contain algae (Shields et al. 1985). Although SOK fisheries do not directly remove spawning Pacific Herring, they do remove eggs that could otherwise have contributed to recruitment. Note that closed-pond operations also cause incidental mortality to spawning Pacific Herring (Shields et al. 1985), but we do not address this issue here. Thus, SOK fisheries present an issue in terms of their impact to the population, and accounting in stock assessment and monitoring (Schweigert et al. 2018). Although Pacific Herring stock assessments do not account for eggs removed by SOK fisheries at this time, there are a few options to account for the impact of SOK harvest. The most direct is to estimate the quantity of eggs removed from the population, and treat them as though they would have spawned and contributed to spawning biomass.

Shields et al. (1985) collected information on the relationship between the number of egg layers in SOK product, and proportion of product weight that consists of eggs and kelp. They determined that kelp represents an average of 12% of the total product weight. Since SOK product is universally brined at the time of harvest, it is necessary to also consider the uptake of salt

Table 9. Notation for Pacific Herring total spawn index calculations. Legend: tonnes (t).

Name	Description	Value or unit
B	Spawn index (i.e., biomass)	t

by the eggs, which increases the overall product weight. However, there is uncertainty in the degree of brining that occurs prior to weighing the product. Nevertheless, Whyte and Englar (1977) determined that following a 24-hour brining period, wet product weight increases by about 13% due to salt uptake. By osmosis, brining would also draw some water from the eggs; unfortunately we are unable to account for osmosis at this time. The last factor to consider is the mean fertilized egg weight, which was determined by Hay and Miller (1982) to be $2.38 \cdot 10^{-6}$ kg.

We estimate spawning biomass removed from the population by SOK fishery f in SAR r and year y as

$$B_{fry} = \frac{H_{fry}\nu v}{M\theta} \quad (36)$$

where H_{fry} is the weight in kilograms of Pacific Herring SOK harvest in fishery f , SAR r , and year y , ν is the proportion of SOK product that is eggs, not kelp, v is the proportion of SOK product that is eggs after brining, M is the average mass in kilograms of a fertilized egg, and B_{fry} is spawning biomass in tonnes for SOK fishery f (Table 10). Then we aggregate spawning biomass by stock assessment region (SAR) r and year y

$$B_{ry} = \sum_{f=1}^F B_{fry} \quad (37)$$

where B_{fry} is spawning biomass by fishery f from Equation 36, and B_{ry} is estimated spawning biomass removed by SOK fisheries in SAR r and year y in tonnes.

10 SOURCES OF UNCERTAINTY

Like all biological models, spawn index calculations are affected by various potential sources of uncertainty including natural variability, observation error (e.g., bias, precision), measurement error, and model structural complexity (Link et al. 2012). Some examples illustrate these sources of uncertainty:

1. Natural variability could affect Pacific Herring fecundity, and the sex ratio of spawning Pacific Herring (Equation 8). Fecundity could be influenced by biological processes such as the observed non-stationarity of weight-at-age, or a truncated age distribution caused by selective fishing or high natural mortality.

Table 10. Notation for Pacific Herring spawn on kelp (SOK) calculations. Legend: kilograms (kg), tonnes (t).

Name	Description	Value or unit	Reference
H	Weight of SOK harvest	kg	
ν	Proportion of SOK product that is eggs, not kelp	0.88	Shields et al. (1985)
v	Proportion of SOK product that is eggs after brining	0.885	Whyte and Englar (1977)
M	Average mass of a fertilized egg	2.38×10^{-6} kg	Hay and Miller (1982)
B	Spawning biomass	t	

515 2. Measurement error could affect input data such as the number of
516 egg layers, while model structural complexity could affect estimated
517 prediction model parameters, or the form of their relationship, or both
518 (e.g., Equation 11). In addition, spawn index prediction models are
519 dated, and our understanding of these processes could have changed in
520 the intervening years.

521 3. Uncertainty in fixed parameters that are used as data without error
522 (e.g., Equation 11); uncertainty in spawn index parameters is currently
523 unknown.

524 4. Uncertainty in the number of egg layers for spawn intensity categories
525 (Table 3), and algae coefficients (Table 8). Again, uncertainty in these
526 values is currently unknown.

527 Despite these assumptions and potential sources of uncertainty, the spawn
528 index has typically been reported without quantifying uncertainty (but see
529 Schweigert et al. 1993). Reporting the spawn index without uncertainty may
530 perpetuate the misconception that the spawn index is observed data, whereas
531 it is derived data with assumptions and uncertainties.

532 There are several potential benefits to addressing spawn index uncertainty.
533 First, quantifying uncertainty could identify parameters to target with future
534 research. Potential analyses to quantify spawn index uncertainty include:

535 1. Investigate factors that influence fecundity and sex ratios (Equation 8);

- 536 2. Quantify and report variability in estimated prediction model parameters and equations (e.g., Equation 11);
- 537
- 538 3. Bootstrap observed input data (see Schweigert 1993); and
- 539 4. Conduct sensitivity analyses.

540 Second, acknowledging uncertainty can reduce another source of uncertainty: inadequate communication among scientists, managers, and stakeholders, which can lead to misapplication of scientific advice (Link et al. 2012). Finally, 542 acknowledging uncertainty will increase transparency, and enable users to 543 assess potential impacts to Pacific Herring stock assessments in a management strategy evaluation (MSE) approach (e.g., DFO 2019a). Addressing data and 544 model uncertainty is a required component of MSE approaches (Punt et al. 545 2016).

548 Quantifying uncertainty may also identify options to increase survey program efficiency, in terms of data precision and accuracy. Sampling surveys 549 trade off precision of estimated quantities versus survey effort or cost. Ideally, 550 reducing survey effort does not result in biased target variable estimates. 551 Therefore, understanding this trade-off is important if, for example, budget 552 reductions cause reduced survey effort. Potential strategies to improve spawn 553 survey efficiency include:

- 555 1. Conduct underwater surveys for major spawns in core areas, and surface 556 surveys for other spawns;
- 557 2. Quantify the precision and accuracy of spawn width estimates (e.g., 558 appendix A.1);
- 559 3. Review transect and quadrat spacing (section 2; see Schweigert 1993);
- 560 4. Review egg prediction model accuracy (Equation 8);
- 561 5. Review temporal stability of egg layer estimates; and
- 562 6. Conduct periodic versus annual surveys.

563 Even with a stable budget, there is a trade-off between high survey effort in 564 some areas, versus low survey effort or no information in other areas.

565 11 FUTURE RESEARCH

566 Many of the parameters and prediction models used to calculate the spawn 567 index are dated; these analyses could be checked with new information, and

568 updated if required. Parameters include ω , ϕ^f , ν , v , and M . Prediction
 569 models include Equation 11, Equation 23, Equation 26, and Equation 27. In
 570 addition, parameter and prediction model uncertainty should be propagated
 571 through the calculations to quantify spawn index uncertainty (section 10).
 572 One approach to account for prediction model uncertainty is to incorporate the
 573 underlying data that informs these equations into spawn index calculations.
 574 In addition, future work could review the assumed statistical framework.

575 12 CAVEATS

576 In addition to issues regarding uncertainty (section 10), there are a few caveats
 577 to keep in mind when interpreting the Pacific Herring spawn index, and using
 578 the spawn index in analyses:

- 579 1. The spawn index a relative index of spawning biomass,
- 580 2. The spawn index is derived data with uncertainty and assumptions, as
 581 opposed to observed data (section 10),
- 582 3. There are two different survey periods with substantial differences in
 583 survey effort and method (surface surveys from 1951 to 1987, and dive
 584 surveys from 1988 to present),
- 585 4. Surface surveys use two different methods to estimate the number of
 586 egg layers:
 - 587 (a) spawn intensity categories (five categories from 1951 to 1968, and
 588 nine categories from 1969 to 1978), and
 - 589 (b) direct estimates (from 1979 to present), and
- 590 5. The spawn index is presence only and does not indicate the absence of
 591 spawn.

592 13 DOWNLOAD

593 The **R** package to calculate the Pacific Herring spawn index **SpawnIndex**, is
 594 publicly accessible on the [Pacific Herring spawn index repository](#). The package
 595 includes an example database of Pacific Herring spawn survey observations,
 596 and a vignette with examples. The **R** package contains functions to import
 597 tables from the database, and calculate the spawn index. This document
 598 is meant to accompany the **R** package, which implements the calculations

described here. The **R** package is documented and has examples to promote accessibility and transparency.

14 ACKNOWLEDGEMENTS

The authors thank Ashleen Benson for translating the calculations in the **Microsoft Access** database to an **R** script, which they referenced when developing the **SpawnIndex** package. They also thank Paul Starr and Rob Kronlund for their thorough reviews that improved this document.

15 REFERENCES

- Cleary, J.S., Taylor, N.G., and Haist, V. 2017. Status of BC Pacific Herring (*Clupea pallasii*) in 2013 and forecasts for 2014. Research Document 2017/014, Canadian Science Advisory Secretariat, Fisheries and Oceans Canada. URL <http://cat.fsl-bsf.scitech.gc.ca/record=b4061206~S1>
- DFO (Fisheries and Oceans Canada). 2019a. Evaluation of management procedures for Pacific Herring (*Clupea pallasii*) in the Strait of Georgia and the West Coast of Vancouver Island management areas of British Columbia. Science Advisory Report 2019/001, Canadian Science Advisory Secretariat, Fisheries and Oceans Canada. URL http://www.dfo-mpo.gc.ca/csas-sccs/Publications/SAR-AS/2019/2019_001-eng.html
- DFO (Fisheries and Oceans Canada). 2019b. Status of Pacific Herring (*Clupea pallasii*) in 2018 and forecast for 2019. Science Response 2019/001, Canadian Science Advisory Secretariat, Fisheries and Oceans Canada. URL http://www.dfo-mpo.gc.ca/csas-sccs/Publications/SCR-RS/2019/2019_001-eng.html
- Haegle, C.W., Hourston, A.S., Humphreys, R.D., and Miller, D.C. 1979. Eggs per unit area in British Columbia herring spawn depositions. Fisheries and Marine Service Technical Report 894, Department of Fisheries and Oceans. URL <http://cat.fsl-bsf.scitech.gc.ca/record=b3858115~S1>
- Haegle, C.W., and Schweigert, J.F. 1985. Distribution and characteristics of herring spawning grounds and description of spawning behavior. *Canadian Journal of Fisheries and Aquatic Sciences* **42**(S1): 39–55. doi:[10.1139/f85-261](https://doi.org/10.1139/f85-261)

- 631 Haegele, C.W., and Schweigert, J.F. 1990. A model which predicts Pacific
632 Herring (*Clupea harengus pallasii*) egg deposition on giant kelp (*Macrocystis*
633 sp.) plants from underwater observations. Canadian Manuscript Report
634 of Fisheries and Aquatic Sciences 2056, Fisheries and Oceans Canada.
635 URL <http://cat.fsl-bsf.scitech.gc.ca/record=b3898101~S1>
- 636 Haist, V., and Rosenfeld, L. 1988. Definitions and codings of localities,
637 sections, and assessment regions for British Columbia herring data. Cana-
638 dian Manuscript Report of Fisheries and Aquatic Science 1994, Fisheries
639 and Oceans Canada. URL [http://cat.cisti-icist.nrc-cnrc.gc.ca/
640 record=b3927024~S1](http://cat.cisti-icist.nrc-cnrc.gc.ca/record=b3927024~S1)
- 641 Hart, J.L., and Tester, A.L. 1934. Quantitative studies on herring spawn-
642 ing. *Transactions of the American Fisheries Society* **64**(1): 307–312.
643 doi:[10.1577/1548-8659\(1934\)64\[307:qsohs\]2.0.co;2](https://doi.org/10.1577/1548-8659(1934)64[307:qsohs]2.0.co;2)
- 644 Hay, D.E. 1985. Reproductive biology of Pacific Herring (*Clupea harengus*
645 *pallasii*). *Canadian Journal of Fisheries and Aquatic Sciences* **42**(S1):
646 111–126. doi:[10.1139/f85-267](https://doi.org/10.1139/f85-267)
- 647 Hay, D.E., and Brett, J.R. 1988. Maturation and fecundity of Pacific Herring
648 (*Clupea harengus pallasii*): An experimental study with comparisons to
649 natural populations. *Canadian Journal of Fisheries and Aquatic Sciences*
650 **45**(3): 399–406. doi:[10.1139/f88-048](https://doi.org/10.1139/f88-048)
- 651 Hay, D.E., and Kronlund, A.R. 1987. Factors affecting the distribution,
652 abundance, and measurement of Pacific Herring (*Clupea harengus pal-*
653 *lasi*) spawn. *Canadian Journal of Fisheries and Aquatic Sciences* **44**(6).
654 doi:[10.1139/f87-141](https://doi.org/10.1139/f87-141)
- 655 Hay, D.E., and Miller, D.C. 1982. A quantitative assessment of herring spawn
656 lost by storm action in French Creek, 1980. Canadian Manuscript Report of
657 Fisheries and Aquatic Sciences 1636, Department of Fisheries and Oceans.
658 URL <http://cat.fsl-bsf.scitech.gc.ca/record=b3849753~S1>
- 659 Humphreys, R.D., and Hourston, A.S. 1978. British Columbia herring spawn
660 deposition manual. Miscellaneous Special Publication 38, Department of
661 Fisheries and the Environment, Fisheries and Marine Service. URL [http:
662 //cat.fsl-bsf.scitech.gc.ca/record=b3686454~S1](http://cat.fsl-bsf.scitech.gc.ca/record=b3686454~S1)
- 663 Jessen, R.J. 1978. Statistical survey techniques. John Wi-
664 ley & Sons. URL [https://openlibrary.org/books/OL4552019M/
665 Statistical_survey_techniques](https://openlibrary.org/books/OL4552019M/Statistical_survey_techniques)
- 666 Link, J.S., Ihde, T.F., Harvey, C.J., Gaichas, S.K., Field, J.C., Brodziak,

- 667 J.K.T., Townsend, H.M., and Peterman, R.M. 2012. Dealing with
668 uncertainty in ecosystem models: The paradox of use for living ma-
669 rine resource management. *Progress in Oceanography* **102**: 102–114.
670 doi:[10.1016/j.pocean.2012.03.008](https://doi.org/10.1016/j.pocean.2012.03.008)
- 671 Marwick, B., Boettiger, C., and Mullen, L. 2018. Packaging data analytical
672 work reproducibly with R (and friends). *PeerJ Preprints (not peer reviewed)*
673 doi:[10.7287/peerj.preprints.3192v2](https://doi.org/10.7287/peerj.preprints.3192v2)
- 674 Punt, A.E., Butterworth, D.S., de Moor, C.L., De Oliveira, J.A.A., and
675 Haddon, M. 2016. Management strategy evaluation: Best practices. *Fish*
676 *and Fisheries* **17**(2): 303–334. doi:[10.1111/faf.12104](https://doi.org/10.1111/faf.12104)
- 677 Quinn, T.J., Best, E.A., Bijsterveld, L., and McGregor, I.R. 1983. Sampling
678 Pacific halibut (*Hippoglossus stenolepis*) landings for age composition:
679 History, evaluation, and estimation. Scientific Report 68, International
680 Pacific Halibut Commission
- 681 RCT (R Core Team). 2017. R: A language and environment for statistical com-
682 puting. URL <http://www.R-project.org>. R Foundation for Statistical
683 Computing. Vienna, Austria. Version 3.4.1 64 bit
- 684 Schweigert, J., Cleary, J., and Midgley, P. 2018. Synopsis of the Pacific Herring
685 spawn-on-kelp fishery in British Columbia. Canadian Manuscript Report
686 of Fisheries and Aquatic Sciences 3148, Fisheries and Oceans Canada.
687 URL <http://cat.fsl-bsf.scitech.gc.ca/record=b4068121~S1>
- 688 Schweigert, J.F. 1993. A review and evaluation of methodology for
689 estimating Pacific Herring egg deposition. *Bulletin of Marine Sci-*
690 *ence* **53**(2). URL [http://www.ingentaconnect.com/content/umrsmas/](http://www.ingentaconnect.com/content/umrsmas/bullmar/1993/00000053/00000002/art00019)
691 [bullmar/1993/00000053/00000002/art00019](http://www.ingentaconnect.com/content/umrsmas/bullmar/1993/00000053/00000002/art00019)
- 692 Schweigert, J.F. 2005. An assessment framework for Pacific Herring (*Clupea*
693 *pallasi*) stocks in British Columbia. Research Document 2005/083, Fisheries
694 and Oceans Canada. URL [http://cat.cisti-icist.nrc-cnrc.gc.ca/](http://cat.cisti-icist.nrc-cnrc.gc.ca/record=4025049)
695 [record=4025049](http://cat.cisti-icist.nrc-cnrc.gc.ca/record=4025049)
- 696 Schweigert, J.F., Fort, C., and Hamer, L. 1997. Stock assessment for British
697 Columbia herring in 1996 and forecasts of the potential catch in 1997. Cana-
698 dian Technical Report of Fisheries and Aquatic Sciences 2173, Department
699 of Fisheries and Oceans. URL [http://cat.cisti-icist.nrc-cnrc.gc.](http://cat.cisti-icist.nrc-cnrc.gc.ca/record=b4020685~S1)
700 [ca/record=b4020685~S1](http://cat.cisti-icist.nrc-cnrc.gc.ca/record=b4020685~S1)
- 701 Schweigert, J.F., Haegle, C.W., and Stocker, M. 1985. Optimizing sam-
702 pling design for herring spawn surveys in the Strait of Georgia, B.C.

- 703 *Canadian Journal of Fisheries and Aquatic Sciences* **42**(11): 1806–1814.
704 doi:[10.1139/f85-226](https://doi.org/10.1139/f85-226)
- 705 Schweigert, J.F., Haegele, C.W., and Stocker, M. 1990. Evaluation of sam-
706 pling strategies for scuba surveys to assess spawn deposition by Pacific
707 Herring. *North American Journal of Fisheries Management* **10**(2): 185–195.
708 doi:[10.1577/1548-8675\(1990\)010<0185:eossfs>2.3.co;2](https://doi.org/10.1577/1548-8675(1990)010<0185:eossfs>2.3.co;2)
- 709 Schweigert, J.F., Hay, D.E., and Fort, C. 1993. Herring spawn index analysis.
710 PSARC H93-02, Department of Fisheries and Oceans. URL <http://cat.fsl-bsf.scitech.gc.ca/record=b4018577~S1>
- 712 Schweigert, J.F., and Stocker, M. 1988. Escapement model for estimat-
713 ing Pacific Herring stock size from spawn survey data and its manage-
714 ment implications. *North American Journal of Fisheries Management* **8**.
715 doi:[10.1577/1548-8675\(1988\)008<0063:EMFEPH>2.3.CO;2](https://doi.org/10.1577/1548-8675(1988)008<0063:EMFEPH>2.3.CO;2)
- 716 Sen, A.R. 1984. Sampling commercial rockfish landings in California. NOAA
717 Technical Memorandum 45, National Marine Fisheries Service. URL <https://repository.library.noaa.gov/view/noaa/6024>
- 719 Shields, T.L., Jamieson, G.S., and Sprout, P.E. 1985. Spawn-on-kelp fisheries
720 in the Queen Charlotte Islands and northern British Columbia coast - 1982
721 and 1983. Canadian Technical Report of Fisheries and Aquatic Sciences
722 1372, Department of Fisheries and Oceans. URL <http://cat.fsl-bsf.scitech.gc.ca/record=b1319605~S1>
- 724 Tanasichuk, R.W., and Ware, D.M. 1987. Influence of interannual variations
725 in winter sea temperature on fecundity and egg size in Pacific Herring
726 (*Clupea harengus pallasii*). *Canadian Journal of Fisheries and Aquatic*
727 *Sciences* **44**(8): 1485–1495. doi:[10.1139/f87-178](https://doi.org/10.1139/f87-178)
- 728 Ware, D.M. 1985. Life history characteristics, reproductive value, and re-
729 siliance of Pacific Herring (*Clupea harengus pallasii*). *Canadian Journal of*
730 *Fisheries and Aquatic Sciences* **42**: 127–137. doi:[10.1139/f85-268](https://doi.org/10.1139/f85-268)
- 731 Whyte, J.N.C., and Englar, J.R. 1977. Aspects of the production of herring roe
732 on *Macrocystis integrifolia* in Georgia Strait locations. Fisheries and Marine
733 Service Technical Report 751, Fisheries and Marine Service. URL <http://cat.fsl-bsf.scitech.gc.ca/record=b1115904~S1>
- 735 Wilson, G., Bryan, J., Cranston, K., Kitzes, J., Nederbragt, L., and Teal, T.
736 2016. Good Enough Practices in Scientific Computing. *PLOS* URL <https://arxiv.org/abs/1609.00037v2>
- 737

738 Xie, Y. 2015. Dynamic documents with R and knitr. The R series. Chapman
739 and Hall/CRC, Florida, USA, 2nd ed. URL [https://github.com/yihui/
740 knitr-book](https://github.com/yihui/knitr-book). ISBN 978-1498716963

741 APPENDIX A SPAWN WIDTH ADJUSTMENTS

742 Spawn width is a critical component in spawn index calculations. There are
743 two cases where we adjust spawn width estimates to improve spawn index
744 accuracy: surface surveys in all years from 1951 to present, and understory
745 dive surveys from 2003 to 2014.

746 A.1 SURFACE SPAWN WIDTH

747 Surface surveys were the only survey type prior to 1988, while the majority
748 of spawns since 1988 have been surveyed using SCUBA gear. Recall that we
749 describe the spawn index as having two periods based on the predominant
750 survey type: the surface survey period from 1951 to 1987, and the dive survey
751 period from 1988 to present.

752 One issue with comparing these two partly overlapping protocols is that
753 surface surveyors tend to underestimate spawn width (Hay and Kronlund
754 1987). To improve the consistency of spawn index estimates throughout the
755 time period from 1951 to present, we adjust surface spawn width estimates
756 using underwater estimates when available (Schweigert et al. 1993). Our
757 preferred width is the median width from all dive surveys within a ‘pool.’ A
758 pool is a group of Locations within a Section that are often adjacent, contain
759 similar algae and substrate, and can be treated as a group with likely similar
760 widths. We summarise spawn width by the median because widths are not
761 normally distributed (Schweigert et al. 1993). If there are no dive data that
762 meet these criteria, we use the median width from all dives within the Section,
763 or within the SAR if there are no dives within the Section. If there are
764 still no dive data that meet these criteria, we use the observed width W'
765 from the surface survey. We update the aforementioned median width values
766 periodically, not annually.

767 **A.2 UNDERSTORY SPAWN WIDTH**

768 In 2013, Fisheries and Oceans Canada (DFO) staff realized that they were
769 inadvertently underestimating spawn width for Pacific Herring understory
770 dive surveys (Cleary et al. 2017). The issue was caused by the assumed
771 accuracy of transect lines used by spawn surveyors to measure spawn width.
772 Spawn surveyors determine spawn width by placing transects perpendicular
773 to the shore. Surveyors use weighted lead lines to ensure that lines rest on
774 the substrate; these lines are marked in 1 m increments, and are standardized
775 to 20 m segments. Segments refer to individual sections of line, which may be
776 linked together to make complete transects.

777 Sometime in the mid- to late-1990s, spawn surveyors observed that the
778 20 m segments shrank by approximately 1 m during the first season of use,
779 and continued to shrink over time. DFO staff noticed that this issue was
780 occurring coast wide, and began re-measuring lead lines each season. They
781 also modified the lead line marking protocol to account for shrinkage by
782 marking 1.15 m increments; thus, segments were extended to 23 m. DFO staff
783 derived this 15% increase by measuring and re-marking lead lines each year.
784 Lead lines are made of a mix of polypropylene and nylon; nylon tightens up
785 under repeated use, which is thought to explain the shrinkage. DFO staff
786 re-measured lead line increments in about 2005, and found that they still
787 shrank from 1.15 m to 1.0 m, and continued to use the modified protocol.

788 In 2013, spawn surveyors observed that lead line increments were con-
789 sistently 1.15 m, and no longer appeared to be shrinking. Following this
790 observation, DFO staff re-measured additional lead lines and found that lead
791 lines were made up of a combination of 1.0 m and 1.15 m increments. The
792 combination of observed increment lengths is explained by the lifespan of lead
793 lines: lead lines are replaced every 5 to 10 years, with some segments being
794 replaced more frequently (i.e., inner segments are replaced more frequently
795 than seaward segments, and segments in some SARs are replaced more fre-
796 quently than in other SARs). DFO staff believe that a change in lead line
797 manufacturing prevents new lead lines from shrinking.

798 The earliest written instructions that describe the modified protocol of
799 marking 1.15 m increments is from 2003, and this protocol was used until
800 2013. Note that some SARs continued to use old lead lines in 2014. The
801 practice of annually re-measuring lead line increments ceased around 2005;
802 thus we are unable to determine when lead lines ceased shrinking. Given

the observations summarized above, we adjust spawn width estimates based on written instructions for the marking protocol in 2003. Accordingly, our best estimate of years impacted by marking lead lines at 1.15 m increments (when shrinking no longer occurred) is from 2003 to 2014. However, not all SARs r and years y are impacted equally by this issue (Table A.1): some SARs and years had all 1.0 m increment lengths (no correction factor needed; $\tau_{ry} = 1.0$), others had all 1.15 m increment lengths ($\tau_{ry} = 1.15$), and others had a combination of 1.0 m and 1.15 m increment lengths which we assume to be in equal proportion ($\tau_{ry} = 1.075$). We correct understory spawn widths by multiplying the observed width by the correction factor

$$W_{txsnry} = W'_{txsnry} \tau_{ry} \quad (\text{A.1})$$

where W'_{txsnry} is the observed spawn width for transect t in spawn survey type x (i.e., understory), spawn s , Location n , SAR r , and year y , τ_{ry} is the spawn width correction factor for SAR r and year y (Table A.1), and W_{txsnry} is the corrected understory spawn width in metres (Table A.2). Instead of updating the database permanently, we adjust spawn width in the **R** package script to be transparent, and to prevent mismatches between the original data sheets and databases.

APPENDIX B SURFACE SPAWN UPDATES

One record in the surface spawn database since 1951 requires an update to fill-in missing egg layer information. As with understory spawn width updates, we make this update in the **R** package script. This affects the following record:

1. Update ‘intensity’ from 0 to 1 for the 1 record in the year 1962, Statistical Area 14, Section 142, Location code 820, and with intensity = 0. We update intensity from 0 to 1 because spawn was surveyed but not reported.

Spawn survey records prior to 1951 have additional missing or inaccurate egg layer information, and are unreliable for indexing purposes. Therefore, we do not include spawn data prior to 1951 in stock assessments.

While reviewing the spawn index calculations and translating them from the **Microsoft Access** database to **R**, we found several cases where index data were being over-written with no documented reason. These updates have been omitted, and affected the following records:

Table A.1. Spawn width correction factors τ_{ry} for Pacific Herring understory spawn surveys by stock assessment region (SAR, r) and year y . Legend: Haida Gwaii (HG), Prince Rupert District (PRD), Central Coast (CC), Strait of Georgia (SoG), West Coast of Vancouver Island (WCVI), Area 27 (A27), and Area 2 West (A2W).

Year	SAR						
	HG	PRD	CC	SoG	WCVI	A27	A2W
2003	1.000	1.075	1.075	1.075	1.075	1.075	1.000
2004	1.000	1.075	1.075	1.075	1.075	1.075	1.000
2005	1.000	1.075	1.075	1.075	1.075	1.075	1.000
2006	1.000	1.075	1.075	1.075	1.075	1.075	1.000
2007	1.000	1.075	1.075	1.075	1.075	1.075	1.000
2008	1.000	1.075	1.075	1.075	1.075	1.075	1.000
2009	1.150	1.075	1.075	1.075	1.075	1.075	1.150
2010	1.150	1.075	1.075	1.075	1.075	1.075	1.150
2011	1.150	1.075	1.075	1.075	1.075	1.075	1.150
2012	1.150	1.075	1.075	1.075	1.075	1.075	1.150
2013	1.150	1.150	1.075	1.075	1.075	1.000	1.150
2014	1.150	1.150	1.000	1.000	1.000	1.000	1.150

Table A.2. Notation for Pacific Herring understory spawn width adjustments. Legend: metres (m).

Name	Description	Value or unit
W'	Observed spawn width	m
τ	Spawn width correction factor	see Table A.1
W	Corrected spawn width	m

1. Update E_{js} (i.e., the total number of egg layers) to 2.1496 for the 15 records in the year 1979, Statistical Area 2, and with intensity 4;
2. Update E_{js} to 0.5529 for the 4 records in the year 1981, Statistical Area 24, and with $E_{js} = 0.0$;
3. Update E_{js} to 1.3360 for the 7 records in the year 1982, Statistical Area 23, and with intensity 3;

- 842 4. Update E_{js} to 2.3300 for 41 records in the year 1984, Statistical Area
843 24, and with intensity 0; and
- 844 5. Update E_{js} to 2.9800 for 14 records in the year 1982, Statistical Area
845 27, and with $E_{js} = 0.0$.
- 846 In the first three cases, E_{js} was updated using intensity categories (Table 3);
847 in the last two cases, E_{js} was updated using historical averages. These changes
848 had negligible effects on spawn index values.