

# Chapter 3 Solutions

Andrew Wu  
Wasserman: All of Statistics

February 17, 2025

**Problem 3.1.** Suppose we play a game where you start with  $c$  dollars. On each turn of the game you either halve or double your money, with equal probability. What is your expected fortune after  $n$  turns?

*Solution.* Let  $X$  be a random variable denoting how much money you have after one turn of the game if you originally have  $a$  dollars.

Then  $\mathbb{E}(X) = \sum x f(x)$ , where  $f(x)$  is the PMF of  $X$ . We have  $f(\frac{a}{2}) = f(2a) = \frac{1}{2}$ , and  $f(x) = 0$  for all other  $x$ .

So

$$\begin{aligned}\mathbb{E}(X) &= 2a \cdot f(2a) + \frac{a}{2} \cdot f\left(\frac{a}{2}\right) \\ &= a + \frac{a}{4} \\ &= \frac{5a}{4}.\end{aligned}$$

Note this is true for any  $a$ , so your expected fortune after  $n$  turns is  $c \cdot (\frac{5}{4})^n$ . □

**Problem 3.2.** Show that  $\mathbb{V}(X) = 0$  if and only if there is a constant  $c$  such that  $\mathbb{P}(X = c) = 1$ .

*Solution.* By definition, we have

$$\mathbb{V}(X) = \mathbb{E}((X - \mathbb{E}(X))^2).$$

If there is a constant  $c$  such that  $\mathbb{P}(X = c) = 1$ , then  $X = c$  and

$$\begin{aligned}\mathbb{V}(X) &= \mathbb{E}((X - c)^2) \\ &= \mathbb{E}(0) = 0.\end{aligned}$$

On the other hand, if  $\mathbb{V}(X) = 0$ , then  $\mathbb{E}((X - \mathbb{E}(X))^2) = 0$ . Note that by the Trivial Inequality,  $(X - \mathbb{E}(X))^2$  is nonnegative, so for the equation  $\mathbb{E}((X - \mathbb{E}(X))^2) = 0$  to hold, we must have  $X - \mathbb{E}(X) = 0$  for all possible values of  $X$ . Thus  $X = \mathbb{E}(X)$ , so  $X$  must be equal to some constant, or equivalently,  $\mathbb{P}(X = c) = 1$  for some constant  $c$ . □

**Problem 3.3.** Let  $X_1, \dots, X_n \sim \text{Uniform}(0, 1)$  and let  $Y_n = \max(X_1, \dots, X_n)$ . Find  $\mathbb{E}(Y_n)$ .

*Solution.* We have

$$\mathbb{E}(Y_n) = \int y f_Y(y) dy,$$

where  $f_Y(y)$  is the PDF of  $Y_n$ .

We begin by finding  $F_Y(y)$ , the CDF of  $Y_n$ . We have

$$\begin{aligned}F_Y(y) &= \mathbb{P}(Y_n \leq y) \\ &= \mathbb{P}(X_1, \dots, X_n \leq y) \\ &= \mathbb{P}(X_1 \leq y) \cdots \mathbb{P}(X_n \leq y) \\ &= y^n.\end{aligned}$$

Thus  $f_Y(y) = ny^{n-1}$ , and

$$\begin{aligned}\mathbb{E}(Y_n) &= \int y f_Y(y) dy \\ &= \int_0^1 ny^n dy \\ &= \frac{n}{n-1}.\end{aligned}$$

□

**Problem 3.4.** A particle stands at the origin of the real line and moves along the line in jumps of one unit. For each jump the probability is  $p$  that the particle will jump one unit to the left and the probability is  $1-p$  that the particle will jump one unit to the right. Let  $X_n$  be the position of the particle after  $n$  jumps. Find  $\mathbb{E}(X_n)$  and  $\mathbb{V}(X_n)$ . (This is known as a random walk.)

*Solution.* Note that  $X_n$  can be written as the sum  $Y_1 + Y_2 + \dots + Y_n$ , where each  $Y_i$  is a random variable equal to  $-1$  with probability  $p$  and equal to  $1$  with probability  $1-p$ , and where all the  $Y_i$ s are independent.

We have

$$\begin{aligned}\mathbb{E}(Y_i) &= (-1) \cdot p + 1 \cdot (1-p) \\ &= 1 - 2p\end{aligned}$$

so therefore

$$\begin{aligned}\mathbb{E}(X_n) &= \mathbb{E}(Y_1 + \dots + Y_n) \\ &= \mathbb{E}(Y_1) + \dots + \mathbb{E}(Y_n) \\ &= n(1 - 2p).\end{aligned}$$

Next, observe that as the  $Y_i$  are independent, we have

$$\begin{aligned}\mathbb{V}(X_n) &= \mathbb{V}(Y_1 + \dots + Y_n) \\ &= \mathbb{V}(Y_1) + \dots + \mathbb{V}(Y_n) \\ &= n\mathbb{V}(Y_i).\end{aligned}$$

Thus we need only compute  $\mathbb{V}(Y_i)$ . We have

$$\begin{aligned}\mathbb{V}(Y_i) &= \mathbb{E}(Y_i^2) - \mathbb{E}(Y_i)^2 \\ &= 1 - (1 - 2p)^2 \\ &= 4p - 4p^2\end{aligned}$$

where we used that  $Y_i^2 = 1$  because  $Y_i = 1$  or  $Y_i = -1$ .

Thus our answer is  $\mathbb{V}(X_n) = n(4p - 4p^2)$ .

□

**Problem 3.5.** A fair coin is tossed until a head is obtained. What is the expected number of tosses that will be required?

*Solution.* Let  $X$  denote the number of tosses necessary to obtain a head, with PMF  $f(x)$ .

Observe that  $f(n) = (\frac{1}{2})^n$ , as one must toss  $n-1$  tails in a row, with probability  $(\frac{1}{2})^{n-1}$ , and then a head, with probability  $\frac{1}{2}$ ; the product of these probabilities is  $(\frac{1}{2})^n$ .

Then

$$\begin{aligned}
\mathbb{E}(X) &= \sum_{i=1}^{\infty} x f(x) \\
&= 1 \cdot \frac{1}{2} + 2 \cdot \frac{1}{4} + 3 \cdot \frac{1}{8} + \dots \\
&= \left( \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots \right) + \left( \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \dots \right) + \left( \frac{1}{8} + \frac{1}{16} + \frac{1}{32} + \dots \right) + \dots \\
&= 1 + \frac{1}{2} + \frac{1}{4} + \dots \\
&= 2.
\end{aligned}$$

□

**Problem 3.6.** Let  $Y = r(X)$  for discrete random variables  $X$  and  $Y$ . Prove that  $\mathbb{E}(Y) = \mathbb{E}(r(X)) = \sum r(x)f_X(x)$ .

*Solution.* Let  $A_y = \{x : r(x) = y\}$ .

We have  $f_Y(y) = \sum_{A_y} f_X(x)$ , for all  $y$ . Then

$$\begin{aligned}
\mathbb{E}(Y) &= \sum_y y f_Y(y) = \sum_y \left( y \sum_{A_y} f_X(x) \right) \\
&= \sum_y \left( y \sum_{x_1, \dots, x_k} f_X(x) \right) \\
&= \sum_x r(x) f_X(x),
\end{aligned}$$

where we used the fact that  $r(x_1) = \dots = r(x_k) = y$ .

□

**Problem 3.7.** Let  $X$  be a continuous random variable with CDF  $F$ . Suppose that  $\mathbb{P}(X > 0) = 1$  and that  $\mathbb{E}(X)$  exists. Show that  $\mathbb{E}(X) = \int_0^{\infty} \mathbb{P}(X > x) dx$ .

*Solution.* We have  $\mathbb{E}(X) = \int x f(x) dx$  by definition, and as  $\mathbb{P}(X > 0) = 1$ , we have

$$\mathbb{E}(X) = \int_0^{\infty} x f(x) dx.$$

Now we integrate by parts. We have

$$\begin{aligned}
\int_0^{\infty} x f(x) dx &= [x F(x)]_0^{\infty} - \int_0^{\infty} F(x) dx \\
&= \lim_{x \rightarrow \infty} x F(x) - \int_0^{\infty} F(x) dx \\
&= \lim_{x \rightarrow \infty} x - \int_0^{\infty} F(x) dx \\
&= \int_0^{\infty} 1 dx - \int_0^{\infty} F(x) dx \\
&= \int_0^{\infty} 1 - F(x) dx \\
&= \int_0^{\infty} \mathbb{P}(X > x) dx,
\end{aligned}$$

where we have used the fact that as  $\mathbb{E}(X)$  exists,  $\lim_{x \rightarrow \infty} [x F(x) - x] = 0$ .

□

**Problem 3.8.** Let  $X_1, \dots, X_n$  be IID and let  $\mu = \mathbb{E}(X_i), \sigma^2 = \mathbb{V}(X_i)$ . Prove that

$$\mathbb{E}(\bar{X}_n) = \mu, \mathbb{V}(\bar{X}_n) = \frac{\sigma^2}{n}, \quad \text{and} \quad \mathbb{E}(S_n^2) = \sigma^2.$$

*Solution.* We have

$$\begin{aligned} \mathbb{E}(\bar{X}_n) &= \mathbb{E}\left(\frac{X_1 + \dots + X_n}{n}\right) \\ &= \frac{1}{n} (\mathbb{E}(X_1) + \dots + \mathbb{E}(X_n)) \\ &= \frac{1}{n} (n \cdot \mu) = \mu. \end{aligned}$$

Next,

$$\begin{aligned} \mathbb{V}(\bar{X}_n) &= \mathbb{V}\left(\frac{X_1 + \dots + X_n}{n}\right) \\ &= \frac{1}{n^2} (\mathbb{V}(X_1) + \dots + \mathbb{V}(X_n)) \\ &= \frac{1}{n^2} (n\sigma^2) = \frac{\sigma^2}{n}. \end{aligned}$$

To solve the last part, we expand the summation:

$$\begin{aligned} \mathbb{E}(S_n^2) &= \mathbb{E}\left(\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X}_n)^2\right) \\ &= \frac{1}{n-1} \mathbb{E}\left(\sum_{i=1}^n (X_i - \bar{X}_n)^2\right) \\ &= \frac{1}{n-1} \mathbb{E}\left(\sum_{i=1}^n (X_i^2 + \bar{X}_n^2 - 2X_i\bar{X}_n)\right) \\ &= \frac{1}{n-1} \mathbb{E}\left(\sum_{i=1}^n X_i^2 + \sum_{i=1}^n \bar{X}_n^2 - 2 \sum_{i=1}^n (X_i\bar{X}_n)\right) \\ &= \frac{1}{n-1} \mathbb{E}\left(\sum_{i=1}^n X_i^2 + n \cdot \bar{X}_n^2 - \frac{2}{n} \sum_{i=1}^n (X_i(X_1 + \dots + X_n))\right) \\ &= \frac{1}{n-1} \mathbb{E}\left(\sum_{i=1}^n X_i^2 + \frac{1}{n} (X_1 + \dots + X_n)^2 - \frac{2}{n} (X_1 + \dots + X_n)^2\right) \\ &= \frac{1}{n-1} \mathbb{E}\left(\sum_{i=1}^n X_i^2 - \frac{1}{n} (X_1 + \dots + X_n)^2\right) \\ &= \frac{1}{n-1} \mathbb{E}\left(\frac{n-1}{n} (X_1^2 + \dots + X_n^2) - \frac{1}{n} \sum_{1 \leq i < j \leq n} 2X_i X_j\right). \end{aligned}$$

Now we can use linearity of expectation. Begin by noting that

$$\begin{aligned} \frac{1}{n-1} \mathbb{E}\left(\frac{n-1}{n} (X_1^2 + \dots + X_n^2)\right) &= \frac{1}{n} \mathbb{E}(X_1^2 + \dots + X_n^2) \\ &= \frac{1}{n} (n \cdot \mathbb{E}(X_i^2)) = \mathbb{E}(X_i^2). \end{aligned}$$

Next, observe that

$$\begin{aligned}\frac{1}{n-1}\mathbb{E}\left(\frac{1}{n}\sum_{1\leq i<j\leq n}2X_iX_j\right) &= \frac{2}{n(n-1)}\mathbb{E}\left(\sum_{1\leq i<j\leq n}X_iX_j\right) \\ &= \frac{2}{(n(n-1))}\cdot\frac{n(n-1)}{2}\cdot\mathbb{E}(X_iX_j) \\ &= \mu^2,\end{aligned}$$

using the fact that there are  $\binom{n}{2}$  pairs  $(i, j)$  with  $1 \leq i < j \leq n$ .

Putting it all together, we obtain

$$\mathbb{E}(S_n^2) = \mathbb{E}(X_i^2) - \mu^2 = \sigma^2,$$

as desired. □

**Problem 3.10.** Let  $X \sim N(0, 1)$  and let  $Y \sim e^X$ . Find  $\mathbb{E}(Y)$  and  $\mathbb{V}(Y)$ .

*Solution.* Note that  $e^x$  is a strictly monotone increasing function. Therefore,

$$f_Y(y) = f_X(s(y)) \left| \frac{ds(y)}{dy} \right|$$

where  $s$  is the inverse of  $e^x$ .

Thus

$$\begin{aligned}f_Y(y) &= \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{1}{2}s(y)^2\right) \cdot \left| \frac{ds(y)}{dy} \right| \\ &= \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{1}{2}\ln(y)^2\right) \cdot \left| \frac{1}{y} \right| \\ &= \frac{1}{y\sqrt{2\pi}} \exp\left(-\frac{1}{2}\ln(y)^2\right)\end{aligned}$$

when  $y > 0$ , using the fact that  $Y$  must be positive.

Now we want to compute

$$\begin{aligned}\mathbb{E}(Y) &= \int y f_Y(y) dy \\ &= \int_0^\infty \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{1}{2}\ln(y)^2\right) dy \\ &= \frac{1}{\sqrt{2\pi}} \int_0^\infty \exp\left(-\frac{1}{2}\ln(y)^2\right) dy.\end{aligned}$$

A  $u$ -substitution with  $\ln y = u$  yields

$$\frac{1}{\sqrt{2\pi}} \int_0^\infty \exp\left(-\frac{1}{2}\ln(y)^2\right) dy = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty \exp\left(-\frac{1}{2}u^2 - u\right) du.$$

(Note that one can obtain the right-hand-side directly through the Law of the Unconscious Statistician.) We'll evaluate the integral by itself and multiply back by  $\frac{1}{\sqrt{2\pi}}$  later, using the method of completing the square. Observe that

$$\begin{aligned}\int_{-\infty}^\infty \exp\left(-\frac{1}{2}u^2 - u\right) du &= e^{\frac{1}{2}} \int_{-\infty}^\infty \exp\left(-\frac{1}{2}u^2 - u - \frac{1}{2}\right) du \\ &= e^{\frac{1}{2}} \int_{-\infty}^\infty \exp\left(-\left(\frac{1}{\sqrt{2}}u + \frac{1}{\sqrt{2}}\right)^2\right) du \\ &= \sqrt{2e} \int_{-\infty}^\infty \exp(-v^2) dv\end{aligned}$$

where the last step follows from substituting  $v = \frac{1}{\sqrt{2}}u + \frac{1}{\sqrt{2}}$ ,  $du = \sqrt{2}dv$ .

We know that  $\int_{-\infty}^{\infty} \exp(-v^2)dv = \sqrt{\pi}$ , due to a well-established result. Putting everything together, it follows that

$$\begin{aligned}\mathbb{E}(Y) &= \frac{1}{\sqrt{2\pi}} \cdot \sqrt{2e} \cdot \sqrt{\pi} \\ &= \sqrt{e}.\end{aligned}$$

Next, to find  $\mathbb{V}(Y)$ , we begin by computing  $\mathbb{E}(Y^2)$ . By the Law of the Unconscious Statistician and our previous computation of  $f_Y(y)$ , we obtain

$$\begin{aligned}\mathbb{E}(Y^2) &= \int_0^{\infty} y^2 \cdot \frac{1}{y\sqrt{2\pi}} \exp\left(-\frac{1}{2}\ln(y)^2\right) dy \\ &= \frac{1}{\sqrt{2\pi}} \int_0^{\infty} y \exp\left(-\frac{1}{2}\ln(y)^2\right) dy.\end{aligned}$$

Let  $u = \ln y$ , such that  $y = e^u$  and  $dy = e^u du$ . Again, we deal with the integral separately from the constant factor, and multiply that back later. We have

$$\begin{aligned}\int_0^{\infty} y \exp\left(-\frac{1}{2}\ln(y)^2\right) dy &= \int_{-\infty}^{\infty} \exp\left(2u - \frac{1}{2}u^2\right) du \\ &= e^2 \int_{-\infty}^{\infty} \exp\left(-\left(\frac{1}{\sqrt{2}}u - \sqrt{2}\right)^2\right) du \\ &= e^2 \sqrt{2} \int_{-\infty}^{\infty} \exp(-v^2) dv\end{aligned}$$

where like before, we substitute  $v = \frac{1}{\sqrt{2}}u - \sqrt{2}$ .

Again, using  $\int_{-\infty}^{\infty} \exp(-v^2)dv = \sqrt{\pi}$ , and putting everything together, we obtain

$$\begin{aligned}\mathbb{E}(Y^2) &= \frac{1}{\sqrt{2\pi}} e^2 \cdot \sqrt{2} \cdot \sqrt{\pi} \\ &= e^2.\end{aligned}$$

Thus

$$\begin{aligned}\mathbb{V}(Y) &= \mathbb{E}(Y^2) - \mathbb{E}(Y)^2 \\ &= e^2 - e\end{aligned}$$

and we are done. □

**Problem 3.12.** Prove the formulas given at the beginning of section 3.4, regarding mean and variance, for the following distributions: Bernoulli, Poisson, Uniform, Exponential, Gamma, Beta.

*Solution.* For each part of this solution, suppose  $X$  is a random variable distributed according to the distribution we are investigating.

For the **Bernoulli** distribution: we have  $\mathbb{P}(X = 1) = p$  and  $\mathbb{P}(X = 0) = 1 - p$ . Thus  $\mathbb{E}(X) = \sum xf(x) = 1 \cdot p + 0 \cdot (1 - p) = p$ . Then,  $\mathbb{V}(X) = \mathbb{E}(X^2) - \mathbb{E}(X)^2$ , and as  $X^2 = X$  when  $X = 0$  or  $X = 1$ , we obtain  $\mathbb{V}(X) = \mathbb{E}(X) - \mathbb{E}(X)^2 = p - p^2$ .

For the **Poisson** distribution: we have  $\mathbb{P}(X = x) = e^{-\lambda} \frac{\lambda^x}{x!}$ . Thus we compute

$$\begin{aligned}\mathbb{E}(X) &= \sum_{x=0}^{\infty} x \cdot e^{-\lambda} \frac{\lambda^x}{x!} \\ &= e^{-\lambda} \cdot \lambda \cdot \sum_{x=1}^{\infty} \frac{\lambda^{x-1}}{(x-1)!} \\ &= e^{-\lambda} \cdot \lambda \cdot e^{\lambda} = \lambda.\end{aligned}$$

To find the variance, we first compute

$$\begin{aligned}\mathbb{E}(X(X-1)) &= \sum_{x=0}^{\infty} x(x-1)e^{-\lambda} \frac{\lambda^x}{x!} \\ &= e^{-\lambda} \lambda^2 \sum_{x=2}^{\infty} \frac{\lambda^{x-2}}{(x-2)!} \\ &= \lambda^2.\end{aligned}$$

Thus  $\mathbb{E}(X^2 - X) = \lambda^2$ , so using linearity of expectation and our previous work, we obtain  $\mathbb{E}(X^2) = \lambda^2 + \lambda$ . Finally, we have  $\mathbb{V}(X) = \mathbb{E}(X^2) - \mathbb{E}(X)^2 = \lambda^2 + \lambda - \lambda^2 = \lambda$ .

For the **Uniform** distribution: we have  $f(x) = \frac{1}{b-a}$  for all  $x \in (a, b)$ . Thus

$$\begin{aligned}\mathbb{E}(X) &= \int_a^b x \cdot \frac{1}{b-a} dx \\ &= \frac{1}{b-a} \int_a^b x dx \\ &= \frac{1}{b-a} \left( \frac{b^2}{2} - \frac{a^2}{2} \right) \\ &= \frac{a+b}{2}.\end{aligned}$$

Then,

$$\begin{aligned}\mathbb{E}(X^2) &= \int_a^b x^2 \cdot \frac{1}{b-a} dx \\ &= \frac{1}{b-a} \int_a^b x^2 dx \\ &= \frac{1}{b-a} \left( \frac{b^3}{3} - \frac{a^3}{3} \right) \\ &= \frac{1}{3}(a^2 + ab + b^2).\end{aligned}$$

Thus

$$\begin{aligned}\mathbb{V}(X) &= \mathbb{E}(X^2) - \mathbb{E}(X)^2 \\ &= \frac{1}{3}(a^2 + ab + b^2) - \left( \frac{a+b}{2} \right)^2 \\ &= \frac{1}{12}(a^2 + b^2) - \frac{1}{6}ab \\ &= \frac{1}{12}(b-a)^2.\end{aligned}$$

For the **Exponential** distribution: we have  $f(x) = \frac{1}{\beta}e^{-x/\beta}$  when  $x > 0$ . Thus

$$\begin{aligned}\mathbb{E}(X) &= \int_0^{\infty} x \cdot \frac{1}{\beta} e^{-x/\beta} dx \\ &= \frac{1}{\beta} \int_0^{\infty} x e^{-x/\beta} dx \\ &= \frac{1}{\beta} \left[ \left[ x \cdot (-\beta) e^{-x/\beta} \right]_0^{\infty} - \int_0^{\infty} (-\beta) e^{-x/\beta} dx \right] \\ &= \frac{1}{\beta} \left[ \lim_{x \rightarrow \infty} (-\beta \cdot x e^{-x/\beta}) + \beta^2 \int_0^{\infty} f(x) dx \right] \\ &= - \lim_{x \rightarrow \infty} (x e^{-x/\beta}) + \beta\end{aligned}$$

where we used integration by parts. Now we must simply compute this limit. Here we can use L'Hopital's rule:

$$\begin{aligned}\lim_{x \rightarrow \infty} x e^{-x\beta} &= \lim_{x \rightarrow \infty} \frac{x}{e^{x\beta}} \\ &= \lim_{x \rightarrow \infty} \frac{1}{\beta e^{x\beta}} \\ &= 0.\end{aligned}$$

Thus we obtain  $\mathbb{E}(X) = \beta$ . Next, we have

$$\begin{aligned}\mathbb{V}(X) &= \mathbb{E}(X^2) - \mathbb{E}(X)^2 \\ &= \int_0^\infty x^2 \cdot \frac{1}{\beta} e^{-x/\beta} dx - \beta^2.\end{aligned}$$

We handle the integral separately, again with integration by parts:

$$\begin{aligned}\int_0^\infty x^2 e^{-x/\beta} dx &= \left[ x^2 (-\beta) e^{-x/\beta} \right]_0^\infty - \int_0^\infty (-\beta) e^{-x/\beta} \cdot 2x dx \\ &= -\beta \lim_{x \rightarrow \infty} \frac{x^2}{e^{x/\beta}} + 2\beta^2 \int_0^\infty \frac{1}{\beta} e^{-x/\beta} \cdot x dx \\ &= 2\beta^2 \cdot \mathbb{E}(X) = 2\beta^3\end{aligned}$$

where we use L'Hopital to take care of the limit and our work on  $\mathbb{E}(X)$  to simplify the integral.

Thus

$$\begin{aligned}\mathbb{V}(X) &= \frac{1}{\beta} \cdot 2\beta^3 - \beta^2 \\ &= \beta^2\end{aligned}$$

and we are done.

For the **Gamma** distribution: we have  $f(x) = \frac{1}{\beta^\alpha \Gamma(\alpha)} x^{\alpha-1} e^{-x/\beta}$  for  $x > 0$ . Then

$$\begin{aligned}\mathbb{E}(X) &= \int_0^\infty x \cdot \frac{1}{\beta^\alpha \Gamma(\alpha)} x^{\alpha-1} e^{-x/\beta} dx \\ &= \int_0^\infty \frac{1}{\beta^\alpha \Gamma(\alpha)} x^\alpha e^{-x/\beta} dx \\ &= \beta \cdot \frac{\Gamma(\alpha+1)}{\Gamma(\alpha)} \int_0^\infty \frac{1}{\beta^{\alpha+1} \Gamma(\alpha+1)} x^\alpha e^{-x/\beta} dx\end{aligned}$$

where we use the fact that  $\alpha, \beta$  are constants to move terms outside the integral. Note that what remains inside the integral is the PDF of a random variable distributed according to  $\text{Gamma}(\alpha+1, \beta)$ , so the integral itself evaluates to 1.

Thus, further simplifying the result requires evaluating  $\frac{\Gamma(\alpha+1)}{\Gamma(\alpha)}$ . We proceed now by integration by parts. Observe that

$$\begin{aligned}\Gamma(\alpha+1) &= \int_0^\infty y^\alpha e^{-y} dy \\ &= [y^\alpha (-e^{-y})]_0^\infty + \int_0^\infty e^{-y} y^{\alpha-1} \alpha dy \\ &= \lim_{y \rightarrow \infty} \left( -\frac{y^\alpha}{e^y} \right) + \alpha \int_0^\infty e^{-y} y^{\alpha-1} dy \\ &= \alpha \Gamma(\alpha)\end{aligned}$$

where we used L'Hopital to evaluate the limit. Thus  $\mathbb{E}(X) = \alpha\beta$ .



Evaluating the variance is similar. We have

$$\begin{aligned}\mathbb{E}(X^2) &= \int_0^\infty x^2 \frac{1}{\beta^\alpha \Gamma(\alpha)} x^{\alpha-1} e^{-x/\beta} dx \\ &= \beta^2 \cdot \frac{\Gamma(\alpha+2)}{\Gamma(\alpha)} \int_0^\infty \frac{1}{\beta^{\alpha+2} \Gamma(\alpha+2)} x^{\alpha+1} e^{-x/\beta} dx \\ &= \alpha(\alpha+1)\beta^2\end{aligned}$$

where we used our work previously to conclude that  $\frac{\Gamma(\alpha+2)}{\Gamma(\alpha)} = \frac{\Gamma(\alpha+1)}{\Gamma(\alpha)} \cdot \frac{\Gamma(\alpha+2)}{\Gamma(\alpha+1)} = \alpha \cdot (\alpha+1)$ .

Thus  $\mathbb{V}(X) = \mathbb{E}(X^2) - \mathbb{E}(X)^2 = \alpha(\alpha+1)\beta^2 - \alpha^2\beta^2 = \alpha\beta^2$ .

Finally, for the **Beta** distribution: we have  $f(x) = \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{\alpha-1}(1-x)^{\beta-1}$ .

Thus

$$\begin{aligned}\mathbb{E}(X) &= \int_0^1 x \cdot \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{\alpha-1}(1-x)^{\beta-1} dx \\ &= \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha+\beta+1)} \cdot \frac{\Gamma(\alpha+1)}{\Gamma(\alpha)} \int_0^1 \frac{\Gamma(\alpha+\beta+1)}{\Gamma(\alpha+1)\Gamma(\beta)} x^\alpha (1-x)^{\beta-1} dx \\ &= \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha+\beta+1)} \cdot \frac{\Gamma(\alpha+1)}{\Gamma(\alpha)} \\ &= \frac{\alpha}{\alpha+\beta}.\end{aligned}$$

using our previous work to simplify the result after eliminating the integral.

Also,

$$\begin{aligned}\mathbb{E}(X^2) &= \int_0^1 x^2 \cdot \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{\alpha-1}(1-x)^{\beta-1} dx \\ &= \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha+\beta+2)} \cdot \frac{\Gamma(\alpha+2)}{\Gamma(\alpha)} \int_0^1 \frac{\Gamma(\alpha+\beta+2)}{\Gamma(\alpha+2)\Gamma(\beta)} x^{\alpha+1}(1-x)^{\beta-1} dx \\ &= \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha+\beta+2)} \cdot \frac{\Gamma(\alpha+2)}{\Gamma(\alpha)} \\ &= \frac{1}{(\alpha+\beta)(\alpha+\beta+1)} \cdot \alpha \cdot (\alpha+1).\end{aligned}$$

Thus

$$\begin{aligned}\mathbb{V}(X) &= \mathbb{E}(X^2) - \mathbb{E}(X)^2 \\ &= \frac{\alpha(\alpha+1)}{(\alpha+\beta)(\alpha+\beta+1)} - \frac{\alpha^2}{(\alpha+\beta)^2} \\ &= \frac{\alpha(\alpha+1)(\alpha+\beta) - \alpha^2(\alpha+\beta+1)}{(\alpha+\beta)^2(\alpha+\beta+1)} \\ &= \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}.\end{aligned}$$

Having derived all the desired means and variances, we are finally done.  $\square$

**Problem 3.13.** Suppose we generate a random variable  $X$  in the following way. First, we flip a fair coin. If the coin is heads, take  $X \sim \text{Uniform}(0, 1)$ . If the coin is tails, take  $X \sim \text{Uniform}(3, 4)$ . Find the mean and standard deviation of  $X$ .

*Solution.* Let  $Y \sim \text{Bernoulli}(\frac{1}{2})$  and  $Z \sim \text{Uniform}(0, 1)$  with  $Y$  and  $Z$  independent, such that  $X = 3Y + Z$ . It is easy to see this is an equivalent way of generating  $X$ .

Then

$$\begin{aligned}
\mathbb{E}(X) &= \mathbb{E}(3Y + Z) \\
&= 3\mathbb{E}(Y) + \mathbb{E}(Z) \\
&= 3 \cdot \frac{1}{2} + \frac{1}{2} \\
&= 2.
\end{aligned}$$

Next,

$$\begin{aligned}
\mathbb{V}(X) &= \mathbb{V}(3Y + Z) \\
&= 9\mathbb{V}(Y) + \mathbb{V}(Z) \\
&= 9 \cdot \frac{1}{4} + \frac{1}{12} \\
&= \frac{7}{3}.
\end{aligned}$$

Thus the standard deviation of  $X$  is  $\sqrt{\frac{7}{3}}$ . □

**Problem 3.14.** Let  $X_1, \dots, X_m$  and  $Y_1, \dots, Y_n$  be random variables and let  $a_1, \dots, a_m$  and  $b_1, \dots, b_n$  be constants. Show that

$$\text{Cov} \left( \sum_{i=1}^m a_i X_i, \sum_{j=1}^n b_j Y_j \right) = \sum_{i=1}^m \sum_{j=1}^n a_i b_j \text{Cov}(X_i, Y_j).$$

*Solution.* We have

$$\begin{aligned}
\text{Cov} \left( \sum_{i=1}^m a_i X_i, \sum_{j=1}^n b_j Y_j \right) &= \mathbb{E} \left( \sum_{i=1}^m a_i X_i \cdot \sum_{j=1}^n b_j Y_j \right) - \mathbb{E} \left( \sum_{i=1}^m a_i X_i \right) \mathbb{E} \left( \sum_{j=1}^n b_j Y_j \right) \\
&= \mathbb{E} \left( \sum_{i=1}^m \sum_{j=1}^n a_i b_j X_i Y_j \right) - \mathbb{E} \left( \sum_{i=1}^m a_i X_i \right) \mathbb{E} \left( \sum_{j=1}^n b_j Y_j \right) \\
&= \sum_{i=1}^m \sum_{j=1}^n a_i b_j \mathbb{E}(X_i Y_j) - \left( \sum_{i=1}^m a_i \mathbb{E}(X_i) \right) \left( \sum_{j=1}^n b_j \mathbb{E}(Y_j) \right) \\
&= \sum_{i=1}^m \sum_{j=1}^n [a_i b_j \mathbb{E}(X_i Y_j) - a_i \mathbb{E}(X_i) b_j \mathbb{E}(Y_j)].
\end{aligned}$$

Note now that  $a_i b_j \mathbb{E}(X_i Y_j) - a_i \mathbb{E}(X_i) b_j \mathbb{E}(Y_j) = a_i b_j \text{Cov}(X_i, Y_j)$ , so indeed,

$$\text{Cov} \left( \sum_{i=1}^m a_i X_i, \sum_{j=1}^n b_j Y_j \right) = \sum_{i=1}^m \sum_{j=1}^n a_i b_j \text{Cov}(X_i, Y_j).$$

□

**Problem 3.15.** Let

$$f_{X,Y}(x, y) = \begin{cases} \frac{1}{3}(x + y), & 0 \leq x \leq 1, 0 \leq y \leq 2 \\ 0, & \text{otherwise.} \end{cases}$$

Find  $\mathbb{V}(2X - 3Y + 8)$ .

*Solution.* Begin by noting that  $\mathbb{V}(2X - 3Y + 8) = \mathbb{E}(4X^2 - 12XY + 9Y^2) - \mathbb{E}(2X - 3Y)^2$ , as the constant term is irrelevant.

Let us compute  $f_X(x)$  and  $f_Y(y)$ . We have

$$\begin{aligned} f_X(x) &= \int f_{X,Y}(x,y) dy \\ &= \int_0^2 \frac{1}{3}(x+y) dy \\ &= \frac{2}{3}x + \frac{2}{3} \end{aligned}$$

and through similar methods we obtain  $f_Y(y) = \frac{1}{3}y + \frac{1}{6}$ .

Now, we will compute the expected values  $\mathbb{E}(X)$ ,  $\mathbb{E}(X^2)$ ,  $\mathbb{E}(Y)$ ,  $\mathbb{E}(Y^2)$ ,  $\mathbb{E}(XY)$ . That will suffice to obtain the desired variance.

$$\begin{aligned} \mathbb{E}(X) &= \int_0^1 x \left( \frac{2}{3}x + \frac{2}{3} \right) dx \\ &= \left[ \frac{2}{9}x^3 + \frac{1}{3}x^2 \right]_0^1 = \frac{5}{9}. \\ \mathbb{E}(X^2) &= \int_0^1 x^2 \left( \frac{2}{3}x + \frac{2}{3} \right) dx \\ &= \left[ \frac{1}{6}x^4 + \frac{2}{9}x^3 \right]_0^1 = \frac{7}{18}. \\ \mathbb{E}(Y) &= \int_0^2 y \left( \frac{1}{3}y + \frac{1}{6} \right) dy \\ &= \left[ \frac{1}{9}y^3 + \frac{1}{12}y^2 \right]_0^2 = \frac{11}{9}. \\ \mathbb{E}(Y^2) &= \int_0^2 y^2 \left( \frac{1}{3}y + \frac{1}{6} \right) dy \\ &= \left[ \frac{1}{12}y^4 + \frac{1}{18}y^3 \right]_0^2 = \frac{16}{9}. \\ \mathbb{E}(XY) &= \int \int xy f_{X,Y}(x,y) dy dx \\ &= \int_0^1 \int_0^2 \frac{1}{3}xy(x+y) dy dx \\ &= \int_0^1 \left( \frac{2}{3}x^2 + \frac{8}{9}x \right) dx = \frac{2}{3}. \end{aligned}$$

Thus

$$\begin{aligned} \mathbb{E}(4X^2 - 12XY + 9Y^2) - \mathbb{E}(2X - 3Y)^2 &= 4\mathbb{E}(X^2) - 12\mathbb{E}(XY) + 9\mathbb{E}(Y^2) - (2\mathbb{E}(X) - 3\mathbb{E}(Y))^2 \\ &= 4 \cdot \frac{7}{18} - 12 \cdot \frac{2}{3} + 9 \cdot \frac{16}{9} - \left( 2 \cdot \frac{5}{9} - 3 \cdot \frac{11}{9} \right)^2 \\ &= \frac{86}{9} - \frac{529}{81} = \frac{245}{81}. \end{aligned}$$

□

**Problem 3.16.** Let  $r(x)$  be a function of  $x$  and let  $s(y)$  be a function of  $y$ . Show that

$$\mathbb{E}(r(X)s(Y)|X) = r(X)\mathbb{E}(s(Y)|X).$$

Also, show that  $\mathbb{E}(r(X)|X) = r(X)$ .

*Solution.* We have  $\mathbb{E}(r(X)s(Y)|X) = \int r(x)s(y)f_{Y|X}(y|x)dy$  and  $r(X)\mathbb{E}(s(Y)|X) = r(X) \int s(y)f_{Y|X}(y|x)dy$ . But as the integral is taken  $dy$ , the  $r(x)$  term can be moved outside the integral in the first equation, so we are done.

As for the second part, take  $s(Y) = 1$  in the equation we just proved. Then  $\mathbb{E}(r(X) \cdot 1|X) = r(X)\mathbb{E}(1|X)$ . But that reduces to  $\mathbb{E}(r(X)|X) = r(X)$ , as  $\mathbb{E}(1|X) = 1$ .  $\square$

**Problem 3.17.** Prove that  $\mathbb{V}(Y) = \mathbb{E}(\mathbb{V}(Y|X)) + \mathbb{V}(\mathbb{E}(Y|X))$ .

*Solution.* Note that  $\mathbb{V}(Y) = \mathbb{E}(Y^2) - \mathbb{E}(Y)^2$ . Now we will use the iteration of expectation property.

We have

$$\begin{aligned}\mathbb{E}(Y^2) - \mathbb{E}(Y)^2 &= \mathbb{E}(\mathbb{E}(Y^2|X)) - \mathbb{E}(\mathbb{E}(Y|X))^2 \\ &= \mathbb{E}(\mathbb{V}(Y|X) - \mathbb{E}(Y|X)^2) - \mathbb{E}(\mathbb{E}(Y|X))^2\end{aligned}$$

where we use the fact that

$$\begin{aligned}\mathbb{V}(Y|X) &= \mathbb{E}((Y - \mathbb{E}(Y|X))^2|X) \\ &= \mathbb{E}(Y^2 - 2Y\mathbb{E}(Y|X) + \mathbb{E}(Y|X)^2|X) \\ &= \mathbb{E}(Y^2|X) - 2\mathbb{E}(Y\mathbb{E}(Y|X)) + \mathbb{E}(\mathbb{E}(Y|X)^2|X) \\ &= \mathbb{E}(Y^2|X) - 2\mathbb{E}(Y|X)^2 + \mathbb{E}(Y|X)^2 = \mathbb{E}(Y^2|X) - \mathbb{E}(Y|X)^2.\end{aligned}$$

(Note that the final line follows from our work in the previous problem: indeed,  $2\mathbb{E}(Y\mathbb{E}(Y|X)) = 2 \cdot \mathbb{E}(Y|X)\mathbb{E}(Y|X) = 2\mathbb{E}(Y|X)^2$ , as  $\mathbb{E}(r(X)s(Y)|X) = r(X)\mathbb{E}(s(Y)|X)$  and  $\mathbb{E}(Y|X)$  is a function of  $X$ . Similarly,  $\mathbb{E}(\mathbb{E}(Y|X)^2|X) = \mathbb{E}(Y|X)^2$ , as  $\mathbb{E}(r(X)|X) = r(X)$  and again,  $\mathbb{E}(Y|X)^2$  is a function of  $X$ .)

Finally, observe that

$$\begin{aligned}\mathbb{E}(\mathbb{V}(Y|X) - \mathbb{E}(Y|X)^2) - \mathbb{E}(\mathbb{E}(Y|X))^2 &= \mathbb{E}(\mathbb{V}(Y|X)) - \mathbb{E}(\mathbb{E}(Y|X)^2) - \mathbb{E}(\mathbb{E}(Y|X))^2 \\ &= \mathbb{E}(\mathbb{V}(Y|X)) - \mathbb{V}(\mathbb{E}(Y|X))\end{aligned}$$

where we used linearity of expectation and the definition of  $\mathbb{V}(\mathbb{E}(Y|X))$ .  $\square$

**Problem 3.18.** Show that if  $\mathbb{E}(X|Y = y) = c$  for some constant  $c$ , then  $X$  and  $Y$  are uncorrelated.

*Solution.* Note that if  $\mathbb{E}(X|Y = y) = c$  for some constant  $c$ , then the random variable  $\mathbb{E}(X|Y)$  always takes on the value  $c$ .

Therefore, by iterating expectations,  $\mathbb{E}(X) = \mathbb{E}(\mathbb{E}(X|Y)) = \mathbb{E}(c) = c$ , and  $\mathbb{E}(XY) = \mathbb{E}(\mathbb{E}(XY|Y)) = \mathbb{E}(Y\mathbb{E}(X|Y)) = \mathbb{E}(cY) = c\mathbb{E}(Y)$ .

Thus we have

$$\text{Cov}(X, Y) = \mathbb{E}(XY) - \mathbb{E}(X)\mathbb{E}(Y) = c\mathbb{E}(Y) - c \cdot \mathbb{E}(Y) = 0$$

and it follows that  $X$  and  $Y$  are uncorrelated.  $\square$

**Problem 3.20.** Prove that if  $a$  is a vector and  $X$  is a random vector with mean  $\mu$  and variance  $\Sigma$ , then  $\mathbb{E}(a^T X) = a^T \mu$  and  $\mathbb{V}(a^T X) = a^T \Sigma a$ .

Prove that if  $A$  is a matrix, then  $\mathbb{E}(AX) = A\mu$  and  $\mathbb{V}(AX) = A\Sigma A^T$ .

*Solution.* Note that if

$$X = \begin{pmatrix} X_1 \\ X_2 \\ \vdots \\ X_k \end{pmatrix}$$

then its mean  $\mu$  satisfies

$$\mu = \begin{pmatrix} \mathbb{E}(X_1) \\ \mathbb{E}(X_2) \\ \vdots \\ \mathbb{E}(X_k) \end{pmatrix}$$

and its variance-covariance matrix  $\Sigma$  satisfies

$$\Sigma = \begin{pmatrix} \mathbb{V}(X_1) & \text{Cov}(X_1, X_2) & \cdots & \text{Cov}(X_1, X_k) \\ \text{Cov}(X_2, X_1) & \mathbb{V}(X_2) & \cdots & \text{Cov}(X_2, X_k) \\ \vdots & \vdots & \ddots & \vdots \\ \text{Cov}(X_k, X_1) & \text{Cov}(X_k, X_2) & \cdots & \mathbb{V}(X_k) \end{pmatrix}.$$

Suppose that  $a = \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_k \end{pmatrix}$ . Then

$$\begin{aligned} a^T X &= (a_1 \quad a_2 \quad \cdots \quad a_k) \begin{pmatrix} X_1 \\ X_2 \\ \vdots \\ X_k \end{pmatrix} \\ &= a_1 X_1 + a_2 X_2 + \cdots + a_k X_k \end{aligned}$$

so therefore by linearity of expectation,

$$\begin{aligned} \mathbb{E}(a^T X) &= \mathbb{E}(a_1 X_1 + \cdots + a_k X_k) \\ &= \mathbb{E}(a_1 X_1) + \cdots + \mathbb{E}(a_k X_k) \\ &= a_1 \mathbb{E}(X_1) + \cdots + a_k \mathbb{E}(X_k) \\ &= a^T \mu. \end{aligned}$$

Now, note that

$$\mathbb{V}(a^T X) = \mathbb{V}(a_1 X_1 + \cdots + a_k X_k)$$

and that

$$\begin{aligned} a^T \Sigma a &= (a_1 \quad a_2 \quad \cdots \quad a_k) \begin{pmatrix} \mathbb{V}(X_1) & \text{Cov}(X_1, X_2) & \cdots & \text{Cov}(X_1, X_k) \\ \text{Cov}(X_2, X_1) & \mathbb{V}(X_2) & \cdots & \text{Cov}(X_2, X_k) \\ \vdots & \vdots & \ddots & \vdots \\ \text{Cov}(X_k, X_1) & \text{Cov}(X_k, X_2) & \cdots & \mathbb{V}(X_k) \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_k \end{pmatrix} \\ &= \left( \sum_{i=1}^k a_i \text{Cov}(X_i, X_1) \quad \cdots \quad \sum_{i=1}^k a_i \text{Cov}(X_i, X_k) \right) \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_k \end{pmatrix} \\ &= \sum_{j=1}^k \sum_{i=1}^k a_i a_j \text{Cov}(X_i, X_j) = \sum_{i=1}^k a_i^2 \mathbb{V}(X_i) + 2 \sum_{1 \leq i < j \leq k} a_i a_j \text{Cov}(X_i, X_j). \end{aligned}$$

But that means that  $\mathbb{V}(A^T X) = a^T \Sigma a$ , by Theorem 3.20.

For the second part of the problem, suppose that  $A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1k} \\ a_{21} & a_{22} & \cdots & a_{2k} \\ \vdots & \vdots & \ddots & \vdots \\ a_{k1} & a_{k2} & \cdots & a_{kk} \end{pmatrix}$ . Then

$$AX = \begin{pmatrix} \sum_{i=1}^k a_{1i} X_i \\ \sum_{i=1}^k a_{2i} X_i \\ \vdots \\ \sum_{i=1}^k a_{ki} X_i \end{pmatrix}$$

so thus

$$\begin{aligned}
\mathbb{E}(AX) &= \begin{pmatrix} \mathbb{E}\left(\sum_{i=1}^k a_{1i}X_i\right) \\ \mathbb{E}\left(\sum_{i=1}^k a_{2i}X_i\right) \\ \vdots \\ \mathbb{E}\left(\sum_{i=1}^k a_{ki}X_i\right) \end{pmatrix} \\
&= \begin{pmatrix} \sum_{i=1}^k a_{1i}\mathbb{E}(X_i) \\ \sum_{i=1}^k a_{2i}\mathbb{E}(X_i) \\ \vdots \\ \sum_{i=1}^k a_{ki}\mathbb{E}(X_i) \end{pmatrix} \\
&= A\mu
\end{aligned}$$

by linearity of expectation.

To show that  $\mathbb{V}(AX) = A\Sigma A^T$ , we note first that

$$\begin{aligned}
(X - \mathbb{E}(X))(X - \mathbb{E}(X))^T &= \begin{pmatrix} X_1 - \mathbb{E}(X_1) \\ \vdots \\ X_k - \mathbb{E}(X_k) \end{pmatrix} \begin{pmatrix} X_1 - \mathbb{E}(X_1) & \cdots & X_k - \mathbb{E}(X_k) \end{pmatrix} \\
&= \begin{pmatrix} (X_1 - \mathbb{E}(X_1))(X_1 - \mathbb{E}(X_1)) & \cdots & (X_1 - \mathbb{E}(X_1))(X_k - \mathbb{E}(X_k)) \\ \vdots & \ddots & \vdots \\ (X_k - \mathbb{E}(X_k))(X_1 - \mathbb{E}(X_1)) & \cdots & (X_k - \mathbb{E}(X_k))(X_k - \mathbb{E}(X_k)) \end{pmatrix}
\end{aligned}$$

which shows that, as  $\text{Cov}(A, B) = \mathbb{E}[(A - \mathbb{E}(A))(B - \mathbb{E}(B))]$  for random variables  $A$  and  $B$ , we have  $\mathbb{E}[(X - \mathbb{E}(X))(X - \mathbb{E}(X))^T] = \mathbb{V}(X)$ .

Thus

$$\begin{aligned}
\mathbb{V}(AX) &= \mathbb{E}[(AX - \mathbb{E}(AX))(AX - \mathbb{E}(AX))^T] \\
&= \mathbb{E}[(AX - A\mathbb{E}(X))(AX - A\mathbb{E}(X))^T] \\
&= \mathbb{E}[A(X - \mathbb{E}(X))(X - \mathbb{E}(X))^T A^T] \\
&= A\mathbb{V}(X)A^T
\end{aligned}$$

where we used the properties  $A\mathbb{E}(X) = \mathbb{E}(AX)$  (the previous result) and  $(AB)^T = B^T A^T$ .  $\square$

**Problem 3.21.** Let  $X$  and  $Y$  be random variables. Suppose that  $\mathbb{E}(Y|X) = X$ . Show that  $\text{Cov}(X, Y) = \mathbb{V}(X)$ .

*Solution.* As  $\mathbb{E}(Y|X) = X$ , we know that  $Y = \mathbb{E}(\mathbb{E}(Y|X)) = \mathbb{E}(X)$ .

That means that

$$\begin{aligned}
\text{Cov}(X, Y) &= \mathbb{E}(XY) - \mathbb{E}(X)\mathbb{E}(Y) \\
&= \mathbb{E}(\mathbb{E}(XY|X)) - \mathbb{E}(X)\mathbb{E}(\mathbb{E}(X)) \\
&= \mathbb{E}(X\mathbb{E}(Y|X)) - \mathbb{E}(X)^2 \\
&= \mathbb{E}(X^2) - \mathbb{E}(X)^2 = \mathbb{V}(X)
\end{aligned}$$

as desired. Note that we used the fact that  $\mathbb{E}(\mathbb{E}(X)) = \mathbb{E}(X)$ ; this follows by taking  $Z$  to be a random variable independent of  $X$ , and then noting that  $\mathbb{E}(X) = \mathbb{E}(\mathbb{E}(X|Z)) = \mathbb{E}(\mathbb{E}(X))$  through the law of iterated expectations and the fact that  $\mathbb{E}(X|Z) = \mathbb{E}(X)$ .  $\square$

**Problem 3.22.** Let  $X \sim \text{Uniform}(0, 1)$ . Let  $0 < a < b < 1$ . Let

$$Y = \begin{cases} 1, & 0 < x < b \\ 0 & \text{otherwise} \end{cases}$$

and let

$$Z = \begin{cases} 1, & a < x < 1 \\ 0, & \text{otherwise.} \end{cases}$$

Are  $Y$  and  $Z$  independent? Also, find  $\mathbb{E}(Y|Z)$ .

*Solution.* We have  $\mathbb{P}(Y = 1) = b$  and  $\mathbb{P}(Z = 1) = 1 - a$ . But  $\mathbb{P}(Y = 1, Z = 1) = \mathbb{P}(a < X < b) = b - a$ . Thus  $\mathbb{P}(Y = 1)\mathbb{P}(Z = 1) = b(1 - a) \neq b - a = \mathbb{P}(Y = 1, Z = 1)$ , so  $Y$  and  $Z$  are not independent.

Note that we have  $\mathbb{P}(Y = 1|Z = 1) = \mathbb{P}(0 < x < b|a < x < 1) = \frac{b-a}{1-a}$ , and  $\mathbb{P}(Y = 0|Z = 1) = 1 - \frac{b-a}{1-a} = \frac{1-b}{1-a}$ . Thus  $\mathbb{E}(Y|Z = 1) = 1 \cdot \frac{b-a}{1-a} + 0 \cdot \frac{1-b}{1-a} = \frac{b-a}{1-a}$ .

Moreover, we have  $\mathbb{P}(Y = 1|Z = 0) = 1$ , so  $\mathbb{E}(Y|Z = 0) = 1$ .

Thus it follows that

$$\mathbb{E}(Y|Z) = \begin{cases} 1, & Z = 0 \\ \frac{b-a}{1-a}, & Z = 1. \end{cases}$$

□

**Problem 3.23.** Find the moment generating function for the Poisson, Normal, and Gamma distributions.

*Solution.* For each part of this solution, suppose  $X$  is a random variable distributed according to the distribution we are investigating.

For the **Poisson** distribution: we have  $\mathbb{P}(X = x) = e^{-\lambda} \frac{\lambda^x}{x!}$ . Thus we wish to compute  $\psi_X(t) = \sum_{x=0}^{\infty} e^{tx} e^{-\lambda} \frac{\lambda^x}{x!}$ . The key observation is that  $e^{tx} = (e^t)^x$ , and with this in mind, we see that

$$\begin{aligned} \phi_X(t) &= \sum_{x=0}^{\infty} e^{tx} e^{-\lambda} \frac{\lambda^x}{x!} = e^{-\lambda} \sum_{x=0}^{\infty} \frac{(e^t)^x \lambda^x}{x!} \\ &= e^{-\lambda} \sum_{x=0}^{\infty} \frac{(e^t \lambda)^x}{x!} \\ &= e^{-\lambda} e^{e^t \lambda} \\ &= e^{\lambda(e^t - 1)}. \end{aligned}$$

For the **Normal** distribution: we have  $f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{1}{2\sigma^2}(x - \mu)^2\right)$ . Thus we wish to compute  $\psi_X(t) = \int_{-\infty}^{\infty} e^{tx} \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{1}{2\sigma^2}(x - \mu)^2\right) dx$ .

We have

$$\begin{aligned} \int_{-\infty}^{\infty} \exp\left(-\frac{1}{2\sigma^2}(x - \mu)^2 + tx\right) dx &= \int_{-\infty}^{\infty} \exp\left(-\frac{1}{2\sigma^2}x^2 + \left(\frac{\mu}{\sigma^2} + t\right)x - \frac{\mu^2}{2\sigma^2}\right) dx \\ &= \int_{-\infty}^{\infty} \exp\left(-\left(\frac{1}{\sigma\sqrt{2}}x - \frac{(\frac{\mu}{\sigma^2} + t)\sigma\sqrt{2}}{2}\right)^2 + \left(t\mu + \frac{t^2\sigma^2}{2}\right)\right) dx \\ &= e^{\mu t + \frac{t^2\sigma^2}{2}} \int_{-\infty}^{\infty} \exp\left(-\left(\frac{1}{\sigma\sqrt{2}}x - \frac{(\frac{\mu}{\sigma^2} + t)\sigma\sqrt{2}}{2}\right)^2\right) dx \\ &= e^{\mu t + \frac{t^2\sigma^2}{2}} \cdot \sigma\sqrt{2} \int_{-\infty}^{\infty} \exp(-u^2) du = e^{\mu t + \frac{t^2\sigma^2}{2}} \cdot \sigma\sqrt{2\pi} \end{aligned}$$

where we completed the square and  $u$ -substituted.

Thus we obtain  $\phi_X(t) = \frac{1}{\sigma\sqrt{2\pi}} \cdot e^{\mu t + \frac{t^2 \sigma^2}{2}} \cdot \sigma\sqrt{2\pi} = e^{\mu t + \frac{t^2 \sigma^2}{2}}$ .

For the **Gamma** distribution: we have  $f_X(x) = \frac{1}{\beta^\alpha \Gamma(\alpha)} x^{\alpha-1} e^{-x/\beta}$ . Thus we wish to compute  $\psi_X(t) = \int_0^\infty e^{tx} \frac{1}{\beta^\alpha \Gamma(\alpha)} x^{\alpha-1} e^{-x/\beta} dx$ . In attempting to evaluate the integral, we will make use of the substitution  $-x/\beta + tx = -x/k$ , or equivalently,  $k = \frac{1}{\frac{1}{\beta} - t}$  or  $\frac{k}{1+tk} = \beta$ .

We have

$$\begin{aligned} \int_0^\infty e^{tx} \frac{1}{\beta^\alpha \Gamma(\alpha)} x^{\alpha-1} e^{-x/\beta} dx &= \int_0^\infty \frac{1}{\beta^\alpha \Gamma(\alpha)} x^{\alpha-1} e^{-x/\beta + tx} \\ &= \int_0^\infty \frac{1}{\left(\frac{k}{1+tk}\right)^\alpha \Gamma(\alpha)} x^{\alpha-1} e^{-x/k} dx \\ &= (1+tk)^\alpha \end{aligned}$$

where we used that the integral, after extracting  $(1+tk)^\alpha$ , is just the integral of a  $\text{Gamma}(\alpha, k)$  distribution. (Note that this is valid only when  $\frac{1}{\beta} > t$ , otherwise  $k$  is negative.)

Then, substituting back in for  $k$  yields

$$\begin{aligned} \psi_X(t) &= (1+tk)^\alpha = \left(1 + \frac{t}{\frac{1}{\beta} - t}\right)^\alpha \\ &= \left(\frac{1/\beta}{1/\beta - t}\right)^\alpha \\ &= \left(\frac{1}{1 - \beta t}\right)^\alpha \end{aligned}$$

and we are done. □

**Problem 3.24.** Let  $X_1, \dots, X_n \sim \text{Exp}(\beta)$ . Find the moment generating function of  $X_i$ . Prove that  $\sum_{i=1}^n X_i \sim \text{Gamma}(n, \beta)$ .

*Solution.* Note that the exponential distribution is a special case of the Gamma distribution. Thus, from our work on the previous problem we know that  $\psi_{X_i}(t) = \frac{1}{1 - \beta t}$ .

Now, we know that  $Y = \sum_{i=1}^n X_i$  satisfies  $\psi_Y(t) = \prod_{i=1}^n \psi_{X_i}(t) = \left(\frac{1}{1 - \beta t}\right)^n$ , which is the moment generating function of the  $\text{Gamma}(n, \beta)$  distribution. Therefore  $Y \sim \text{Gamma}(n, \beta)$ . □