

# Sampling Design for Spatial Cluster Randomized Trials Under Heterogeneous Incidence

Journal Title  
XX(X):1–7  
©The Author(s) 2026  
Reprints and permission:  
sagepub.co.uk/journalsPermissions.nav  
DOI: 10.1177/ToBeAssigned  
www.sagepub.com/

SAGE

Andrew Walther<sup>1</sup>, Ross Joseph Simpson, Jr.<sup>2</sup>, Ashkan Habib<sup>1</sup>, and Feng-Chang Lin<sup>1</sup>

## Abstract

**Background/Aims:** Spatial cluster randomized trials (CRTs) are vulnerable to bias from spatial spillover, in which treatment effects diffuse from treated to nearby control clusters, and from spatial dependence in the outcome itself. Existing spillover-mitigation methods are almost entirely analysis-stage corrections applied after data collection. We evaluated whether the choice of treatment assignment design, fixed before a trial begins, can reduce estimation error under spillover without requiring any post-hoc adjustment.

**Methods:** We simulated a 100-cluster spatial grid with outcomes generated from a spatial Durbin model incorporating a spillover term and spatially autocorrelated residual dependence. Baseline outcome incidence was generated under three heterogeneity mechanisms (independent, spatially clustered, and population-scaled Poisson) and used to inform incidence-aware designs. Six treatment assignment strategies, representing three conceptually distinct mechanisms of spillover control (a saturation approach, a buffering/isolation approach, and two incidence-stratified balancing approaches), plus a spatial-separation strategy and a spatial blocking strategy, were compared. The true direct treatment effect was swept across five magnitudes. Each design–parameter combination was replicated under 320 spatial and spillover parameter configurations, with maximum likelihood spatial regression as the estimator throughout. Formal non-parametric hypothesis testing (Friedman and Nemenyi tests) compared mean squared error (MSE) ranks across designs.

**Results:** An incidence-guided saturation design, which varies treatment saturation across spatial regions in proportion to historical incidence, achieved the lowest MSE (0.079) and near-nominal 95% confidence interval coverage (94%). A design that maximizes spatial separation between treated and control clusters (checkerboard-style alternation) performed worst by a wide margin (MSE 0.802) and its coverage collapsed to 53%, meaning its confidence intervals missed the true effect roughly half the time. The MSE ranking of designs was stable across all five treatment-effect magnitudes tested (Friedman  $\chi^2$  range 557.3–800.9, all  $p < 2.2 \times 10^{-16}$ ). A preliminary application-scale illustration, using synthetic placeholder incidence over North Carolina's Community College service-area network as candidate clusters for a planned sudden unexpected death (SUD) prevention pilot, showed a qualitatively consistent pattern favoring incidence-stratified and saturation-based designs.

**Conclusions:** Incidence-informed saturation designs can materially reduce estimation error and preserve valid inference under spatial spillover, while maximal spatial separation, a design intuition borrowed from non-spatial CRT practice, is actively harmful when spillover is present. These findings support incorporating spatial spillover structure into the design stage of cluster randomized trials, ahead of statistical modeling choices made after data collection. Application to a real SUD surveillance dataset, currently in preparation, is needed before a design recommendation for that setting can be finalized.

## Keywords

cluster randomized trial, spatial interference, spillover, treatment assignment, spatial autocorrelation, simulation study, trial design

<sup>1</sup>Department of Biostatistics, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

## 1 Introduction

2 Cluster randomized trials (CRTs) randomize intact groups,  
3 such as clinics, schools, or geographic areas, rather  
4 than individuals, and are the standard design when  
5 an intervention is delivered at the group level or  
6 when individual randomization would create unacceptable  
7 contamination.<sup>1,2</sup> A central assumption underlying standard  
8 CRT analysis is the stable unit treatment value assumption  
9 (SUTVA): a control cluster's outcome must not depend  
10 on which other clusters were treated.<sup>3</sup> This assumption  
11 is routinely violated when clusters are spatially arranged.  
12 Tobler's first law of geography, that near things are more  
13 related than distant things, implies that an intervention  
14 delivered in one location can plausibly influence outcomes  
15 in geographically adjacent locations through shared  
16 populations, service networks, information diffusion, or  
17 environmental pathways.<sup>4</sup> Documented examples of such  
18 spatial spillover span infectious disease control, where  
19 treating one household reduces transmission risk in  
20 neighboring households,<sup>5</sup> and deworming interventions,  
21 where treated schools reduce infection prevalence at  
22 nearby untreated schools.<sup>6</sup> When spillover is present but  
23 unmodeled, standard CRT estimators are biased, and the  
24 direction and magnitude of that bias depend on the spatial  
25 arrangement of treated and control clusters.

26 The methodological literature on spatial interference  
27 has concentrated on analysis-stage correction: detecting  
28 spillover using spatial autocorrelation statistics after  
29 outcomes are observed,<sup>7,8</sup> and fitting spatial regression  
30 models such as the spatial Durbin model to separate  
31 direct and spillover effects post hoc.<sup>9,10</sup> Design-stage  
32 mitigation, choosing a treatment assignment strategy  
33 before randomization that limits spillover's ability to bias  
34 estimation in the first place, has received comparatively  
35 little attention. Jarvis et al. reviewed the sparse literature  
36 on spatial CRT design and highlighted this gap explicitly.<sup>11</sup>  
37 Covariate-constrained randomization offers one partial  
38 design-stage tool, balancing measured covariates across  
39 arms to reduce confounding,<sup>12</sup> building on earlier work  
40 on balance in cluster randomized trials,<sup>13</sup> but it was not  
41 developed with spatial spillover specifically in mind. A  
42 cluster of recent work is now converging on this exact  
43 problem: randomized saturation designs deliberately vary  
44 treatment intensity across units to identify both direct and  
45 spillover effects,<sup>14</sup> fried-egg and buffer-zone designs used  
46 in malaria trials leave an untreated ring around treated  
47 clusters to limit contamination,<sup>15</sup> and a growing theoretical  
48 literature has derived rate-optimal cluster designs and

49 independent-set designs explicitly under network and spatial  
50 interference.<sup>16–19</sup> This body of work signals that the field  
51 is actively moving toward design-stage solutions, but it has  
52 not yet produced a systematic, simulation-based comparison  
53 of design strategies under realistic, heterogeneous spatial  
54 conditions.

55 This paper extends an earlier proof-of-concept study,  
56 which compared simple random assignment against a single  
57 block-stratified design on small, exhaustively enumerable  
58 grids (8–12 clusters) and estimated a structural spatial  
59 Durbin model with separate parameters for baseline, direct,  
60 spillover, and dependence effects. That prior work's own  
61 discussion explicitly anticipated the need for a larger  
62 design space and for heterogeneous, rather than uniform,  
63 baseline outcome risk. This paper answers that call: we  
64 scale to a 100-cluster grid, compare six conceptually  
65 distinct treatment assignment strategies rather than two,  
66 and generate baseline incidence under three heterogeneity  
67 mechanisms so that some designs can exploit prior  
68 knowledge of spatial risk while others cannot.

69 Our aims are threefold: (1) to rank the six designs  
70 by estimation accuracy (mean squared error, MSE) and  
71 inferential validity (95% confidence interval coverage)  
72 within each incidence-generating mechanism; (2) to test  
73 whether that ranking is stable across the spatial and  
74 spillover parameters a trial team cannot control in advance,  
75 including the magnitude of the true treatment effect; and  
76 (3) to translate the ranking into concrete design guidance,  
77 illustrated using a real, currently developing application:  
78 a planned sudden unexpected death (SUD) prevention  
79 pilot delivered through North Carolina's community college  
80 network. The remainder of this paper describes the  
81 simulation methods, reports the design comparison results,  
82 presents the SUD application context, and discusses  
83 implications and limitations.

## 84 Methods

### 85 *Spatial structure and outcome model*

86 We simulated a  $10 \times 10$  regular grid of 100 clusters, with  
87 adjacency defined by queen contiguity (clusters sharing an  
88 edge or corner are neighbors). A row-standardized spatial

2Division of Cardiology, University of North Carolina School of Medicine, Chapel Hill, NC, USA

#### Corresponding author:

Feng-Chang Lin, Department of Biostatistics, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA.

Email: flin33@email.unc.edu

weights matrix  $W$  was constructed from this adjacency structure. Cluster-level outcomes  $Y$  were generated from a spatial Durbin model,

$$Y = (I - \rho W)^{-1} [\tau Z + \gamma WZ + \beta X + \epsilon], \quad \epsilon \sim N(0, \sigma^2 I) \quad (1)$$

where  $Z$  is the binary treatment assignment vector,  $\tau$  is the direct treatment effect of interest,  $\gamma$  is the spillover coefficient governing how much a cluster's outcome is influenced by the treatment status of its neighbors ( $WZ$ ),  $\rho$  is the spatial autoregressive coefficient governing residual dependence in the outcome itself,  $X$  is a baseline incidence covariate with fixed coefficient  $\beta = 1$ , and  $\sigma = 1$ . Spillover was evaluated under two exposure regimes: one in which spillover accrues to control clusters only (as in classic contamination), and one in which it accrues to both treated and control clusters (as when neighboring effects reinforce treatment). All simulation code, exact parameter grids, and reproducibility details are provided in the online supplementary material.

### Heterogeneous baseline incidence

Unlike a setting with uniform baseline risk, real cluster-level health outcomes vary spatially in ways that a trial team may or may not be able to observe in advance. We generated baseline incidence  $X$  under three mechanisms: independent and identically distributed noise (no spatial structure), spatially autocorrelated incidence generated from the same spatial process as the outcome model at two autocorrelation strengths, and a population-scaled Poisson count process calibrated to a sudden-death base rate of 35 per 100,000 (motivated by our applied setting, described below). Incidence was generated once per configuration and held fixed across replications of a given design; only the first replication's "observed" incidence was made available to incidence-aware designs, mimicking how a trial team would use historical surveillance data rather than the true underlying risk surface.

### Treatment assignment designs

We compared six treatment assignment designs, each representing a conceptually distinct strategy for limiting spillover-induced bias (Table 1). Two designs are deterministic given the incidence surface (Checkerboard and High Incidence Focus) and were generated once and held fixed; the remaining four were re-randomized for each replication. Two additional designs evaluated in the underlying simulation study, plain saturation quadrants and a two-strata

(median-split) balancing design, were dropped from this comparison because they were statistically indistinguishable from a retained design that implements the same mechanism with an incidence-informed refinement (Nemenyi  $p = 1.00$  for saturation quadrants versus its incidence-guided counterpart; lower MSE and finer stratification for balanced quartiles versus the median-split variant). The complete eight-design comparison, including these two variants, is reported in the online supplementary material.

### Estimation

The direct treatment effect  $\tau$  was estimated using maximum likelihood spatial regression (`lagsarlm`) fit to a spatial Durbin specification matching the outcome-generating model, including the true spillover exposure term as a covariate. This oracle specification represents a best-case bound on estimation performance and is discussed as a limitation below.

### Simulation design and metrics

The parameter space combined five incidence configurations (independent, two spatial-incidence strengths, two Poisson strengths), two neighbor definitions, four levels of outcome spatial dependence  $\rho$ , four levels of spillover magnitude  $\gamma$ , two spillover exposure regimes, the six designs above, and five true treatment-effect magnitudes ( $\tau \in \{0.8, 1.0, 1.5, 2.0, 3.0\}$ ), totaling 9,600 scenarios for the six-design comparison (12,800 across all eight designs). Each scenario was replicated 25 times over treatment assignment and 10 times over outcome draws (250 total iterations), except the two deterministic designs, which were replicated only over outcome draws. We report bias, standard deviation, MSE, 95% confidence interval coverage, and statistical power to reject a null direct effect of zero. Exact resample counts, the full parameter grid, and metric formulas are given in the online supplementary material.

## Results

Incidence-Guided Saturation Quadrants achieved the lowest MSE (0.079) with coverage close to the nominal 95% level (94%), followed by Balanced Quartiles (MSE 0.102), High Incidence Focus (MSE 0.133), Isolation Buffer (MSE 0.148), and  $2 \times 2$  Blocking (MSE 0.228); all four maintained coverage between 93% and 97% (Table 2, Figure 1). Checkerboard, the design that maximizes spatial separation between treated and control clusters, performed worst by a wide margin (MSE 0.802) and its confidence interval coverage collapsed to 53% (Figure 3a),

**Table 1.** The six treatment assignment designs compared, ordered by mean squared error at the primary treatment-effect magnitude ( $\tau = 1.0$ ; see Table 2). “Deterministic” designs use the same assignment for every simulation replicate within a scenario.

| Design                                | Targeting logic                                                                                                                                | % treated | Deterministic |
|---------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------|-----------|---------------|
| Incidence-Guided Saturation Quadrants | Spatial quadrants receive a randomized saturation level (20–80% treated within quadrant), assigned in rank order of quadrant-average incidence | ~50       | No            |
| Balanced Quartiles                    | Clusters stratified into incidence quartiles; 1:1 randomization within each quartile                                                           | ~50       | No            |
| High Incidence Focus Isolation Buffer | Top 50% of clusters by observed incidence are treated                                                                                          | 50        | Yes           |
|                                       | Greedy assignment such that no two treated clusters are spatially adjacent                                                                     | 20–30     | No            |
| 2×2 Blocking                          | Grid partitioned into 2×2 spatial blocks; 1:1 randomization within each block                                                                  | 50        | No            |
| Checkerboard                          | Alternating assignment across the grid, maximizing spatial separation between treated and control clusters                                     | 50        | Yes           |

**Table 2.** Mean squared error and 95% confidence interval coverage by design at the primary treatment-effect magnitude ( $\tau = 1.0$ ), averaged over 320 spatial and spillover parameter configurations. Friedman rank-sum test:  $\chi^2 = 800.9$ ,  $df = 5$ ,  $p < 2.2 \times 10^{-16}$ . Nemenyi post-hoc comparisons found 13 of 15 pairwise design comparisons statistically significant at  $\alpha = 0.05$ .


| Design                                | MSE   | Coverage |
|---------------------------------------|-------|----------|
| Incidence-Guided Saturation Quadrants | 0.079 | 0.941    |
| Balanced Quartiles                    | 0.102 | 0.938    |
| High Incidence Focus                  | 0.133 | 0.966    |
| Isolation Buffer                      | 0.148 | 0.939    |
| 2×2 Blocking                          | 0.228 | 0.937    |
| Checkerboard                          | 0.802 | 0.527    |

meaning the nominal 95% interval failed to contain the true treatment effect roughly half the time. This is a design-induced inferential failure, not merely a loss of precision: maximizing distance between treated and control units, an intuition carried over from non-spatial CRT practice, concentrates spillover exposure asymmetrically across the grid in a way that a maximum likelihood spatial model estimating a single global  $\tau$  cannot fully correct. The complete eight-design comparison, full parameter-sensitivity breakdowns, and reproducibility details are provided in the online supplementary material.


Decomposing MSE into its squared-bias and variance components clarifies why the designs differ so sharply (Figure 2). Checkerboard’s error is almost entirely variance, not bias: its rigid alternating pattern creates near-perfect collinearity between the treatment vector  $Z$  and the average-neighbor-treatment vector  $WZ$ , so the spatial model cannot stably separate the direct effect from the spillover effect, inflating estimator variance without necessarily shifting its center. Incidence-Guided Saturation Quadrants and Balanced Quartiles, in contrast, achieve low bias and low variance simultaneously, because their heterogeneous, incidence-informed saturation levels preserve contrast between  $Z$  and  $WZ$  rather than collapsing it.

High Incidence Focus, while competitive in mean MSE, showed the largest cross-configuration variability of the top five designs (MSE ranging from 0.078 to 0.207 depending on the incidence-generating mechanism), because its performance is entirely contingent on how informative the observed incidence surface is; when historical incidence poorly reflects the true underlying risk surface, targeting the wrong half of the grid can approach the separation problem that degrades Checkerboard.


This ranking was stable across all five tested treatment-effect magnitudes. The omnibus Friedman test rejected equal design ranks at every level of  $\tau$  (Figure 3b;  $\chi^2 = 730.6, 800.9, 705.7, 658.5$ , and  $557.3$  for  $\tau = 0.8, 1.0, 1.5, 2.0, 3.0$  respectively; all  $p < 2.2 \times 10^{-16}$ ), indicating that the relative advantage of incidence-informed and buffering designs over Checkerboard does not depend on the size of the true effect being detected.



**Figure 1.** Mean squared error by design and spillover exposure regime, at the primary treatment-effect magnitude ( $\tau = 1.0$ ), faceted by outcome spatial dependence  $\rho$  and spillover exposure regime. Designs are ordered best to worst (left to right) by pooled mean MSE, matching Table 2.



**Figure 2.** Bias-squared and variance contributions to MSE for each design (Poisson incidence,  $\rho_X = 0.20$ ,  $\tau = 1.0$ ), ordered best to worst. Checkerboard's MSE is almost entirely variance; Incidence-Guided Saturation Quadrants and Balanced Quartiles achieve low bias and low variance simultaneously.




**Figure 3.** (a) 95% confidence interval coverage by design at  $\tau = 1.0$ . The dashed red line marks the nominal 95% level. (b) Mean squared error by design across the five tested treatment-effect magnitudes.

## Application: A Planned Sudden Unexpected Death Prevention Pilot in North Carolina

Sudden unexpected death (SUD) in adults is a substantial, incompletely understood cause of premature mortality; a validated death-certificate classification algorithm (SUDDEN) has been used to estimate a North Carolina base rate near 35 per 100,000, with productivity loss from SUD in working-age adults exceeding that of any individual cancer,<sup>20</sup> and documented geographic and socioeconomic gradients, including elevated incidence in rural counties<sup>21</sup> and an inverse relationship with household income in Wake County.<sup>22</sup>

A pilot intervention is being planned in collaboration with North Carolina's community college (CC) system, which offers a natural spatial and organizational unit for this design framework. North Carolina's 58 community colleges collectively serve all 100 counties, with a primary campus in one county and satellite campuses in neighboring counties (for example, Durham Technical Community College also operates a campus in Orange County); this network structure is a direct, real-world analog to the spillover mechanism modeled in this paper, because CCs communicate and coordinate with one another. The planned pilot site is Lenoir County Community College, with anticipated spillover to its Kinston and Jones County satellite sites. Because the teaching intervention under consideration is delivered county-wide rather than at finer spatial resolution, the natural treatment unit is the CC service area rather than the census tract. Real SUD incidence data for this application, classified from approximately 100,000 North Carolina death certificates (2018–2021) using the SUDDEN algorithm and pooled to the county level with Surveillance, Epidemiology, and End Results (SEER) population-at-risk denominators, is currently being finalized in collaboration with an epidemiology co-investigator and an active institutional review board protocol; it is not yet available for this manuscript.



**Figure 4.** Synthetic placeholder SUD incidence surface (Poisson SAR process) over the 58 North Carolina Community College service areas, illustrating the spatial heterogeneity this design framework is built to accommodate. **This is a simulated placeholder, not real surveillance data;** it will be replaced with the real SUDDEN-derived county incidence once finalized.

As a preliminary illustration only, we ran the same design-comparison pipeline over the actual 58-cluster CC service-area geography (in place of the abstract  $10 \times 10$  grid) using the synthetic, placeholder incidence surface shown in Figure 4, calibrated to the SUD base rate above. **These application-scale numeric results are a simulated placeholder and must not be interpreted as a finalized design recommendation;** they will be superseded once

the real SUDDEN-derived county dataset described above is finalized. With that caveat, the qualitative pattern was consistent with the primary grid-based results: incidence-stratified and saturation-based designs again achieved the lowest MSE, while designs relying on rigid spatial separation performed markedly worse on this irregular, real-world geography. Planned covariates for the eventual real-data application include county income, food-desert status, physician access, and distance to the nearest hospital. Because SUD rates are not expected to shift detectably on a pilot timescale, the eventual evaluation will likely rely on process measures, such as medication adherence and timely clinic visits, rather than the SUD rate itself; this measurement lag is discussed further below.

## Discussion

These results support treatment assignment design as an actionable, pre-registration-stage tool for mitigating spatial spillover in cluster randomized trials, rather than relying solely on post-hoc statistical correction. In particular, a design intuition imported directly from non-spatial CRT practice, maximizing physical separation between treatment arms, was actively harmful once spillover was present, both inflating MSE and severely undermining inferential validity. Trial teams working in spatially structured settings should not assume that greater separation is automatically safer.

Several limitations bound these conclusions. Estimation throughout used an oracle maximum likelihood specification with the true spillover exposure term supplied as a covariate; this represents a best-case bound; a non-oracle specification, in which the analyst must estimate spillover structure without knowing it in advance, would be expected to show more uniformly degraded performance and is a priority for future work. The outcome model held the incidence coefficient  $\beta$  fixed and assumed equal population size across clusters in the Poisson incidence mode; heterogeneous population sizes, as in the real North Carolina application, may alter the relative performance of incidence-informed designs. The primary comparison used a regular  $10 \times 10$  lattice, whereas the planned application uses an irregular, real-world community college network; the preliminary application-scale run reported above suggests the qualitative ranking is robust to this change, but a fixed grid size was not itself varied. Finally, the process-versus-incidence outcome measurement lag noted in our applied setting, where meaningful SUD rate changes are not expected on a pilot timescale, means that the eventual real-data evaluation of this design framework will need to rely

on intermediate process outcomes rather than the incidence outcome modeled here.

Future work includes relaxing the oracle estimation assumption, incorporating heterogeneous cluster population sizes, evaluating design performance directly on irregular real-world geographies at scale, and, most immediately, applying this design-selection framework to the real SUDDEN-derived North Carolina county dataset once it is finalized.

## Declaration of conflicting interests

The Author(s) declare that there is no conflict of interest.

## Funding

The Author(s) received no financial support for the research, authorship, and/or publication of this article.

## References

- Donner A and Klar N. *Design and analysis of cluster randomization trials in health research*. 1. publ ed. London: Arnold, 2000. ISBN 978-0-470-71100-2 978-0-340-69153-3.
- Hayes RJ and Moulton LH. *Cluster Randomised Trials, Second Edition*. Chapman and Hall/CRC, 2017. ISBN 978-1-315-37028-6. DOI:10.4324/9781315370286. URL <https://www.taylorfrancis.com/books/9781315370286>.
- Hudgens MG and Halloran ME. Toward Causal Inference With Interference. *Journal of the American Statistical Association* 2008; 103(482): 832–842. DOI:10.1198/016214508000000292. URL <https://www.tandfonline.com/doi/full/10.1198/016214508000000292>.
- Tobler WR. A Computer Movie Simulating Urban Growth in the Detroit Region. *Economic Geography* 1970; 46: 234. DOI:10.2307/143141. URL <https://www.jstor.org/stable/143141?origin=crossref>.
- Binka FN, Indome F and Smith T. Impact of spatial distribution of permethrin-impregnated bed nets on child mortality in rural northern Ghana. *The American Journal of Tropical Medicine and Hygiene* 1998; 59(1): 80–85. DOI: 10.4269/ajtmh.1998.59.80.
- Miguel E and Kremer M. Worms: Identifying Impacts on Education and Health in the Presence of Treatment Externalities. *Econometrica* 2004; 72(1): 159–217. DOI:10.1111/j.1468-0262.2004.00481.x. URL <http://doi.wiley.com/10.1111/j.1468-0262.2004.00481.x>.

- 353 7. Moran PAP. The Interpretation of Statistical Maps. 402 04026.  
354 *Journal of the Royal Statistical Society Series B: Statistical* 403  
355 *Methodology* 1948; 10(2): 243–251. DOI:10.1111/j. 404  
356 2517-6161.1948.tb00012.x. URL [https://academic. 405](https://academic.oup.com/jrsssb/article/10/2/243/7026536)  
357 [oup.com/jrsssb/article/10/2/243/7026536](https://academic.oup.com/jrsssb/article/10/2/243/7026536). 406
- 358 8. Geary RC. The Contiguity Ratio and Statistical Mapping. 407  
359 *The Incorporated Statistician* 1954; 5(3): 115. DOI:10.2307/ 408  
360 2986645. URL [https://www.jstor.org/stable/ 409](https://www.jstor.org/stable/2986645?origin=crossref)  
361 [2986645?origin=crossref](https://www.jstor.org/stable/2986645?origin=crossref). 410
- 362 9. Ord K. Estimation Methods for Models of Spatial 411  
363 Interaction. *Journal of the American Statistical Association* 412  
364 1975; 70(349): 120–126. DOI:10.1080/01621459.1975. 413  
365 10480272. URL [http://www.tandfonline.com/ 414](http://www.tandfonline.com/doi/abs/10.1080/01621459.1975.10480272)  
366 [doi/abs/10.1080/01621459.1975.10480272](http://www.tandfonline.com/doi/abs/10.1080/01621459.1975.10480272). 415
- 367 10. Bivand R. R Packages for Analyzing Spatial Data: A 416  
368 Comparative Case Study with Areal Data. *Geographical* 417  
369 *Analysis* 2022; 54(3): 488–518. DOI:10.1111/gean. 418  
370 12319. URL [https://onlinelibrary.wiley.com/ 419](https://onlinelibrary.wiley.com/doi/10.1111/gean.12319)  
371 [doi/10.1111/gean.12319](https://onlinelibrary.wiley.com/doi/10.1111/gean.12319). 420
- 372 11. Jarvis C, Di Tanna GL, Lewis D et al. Spatial anal- 421  
373 ysis of cluster randomised trials: a systematic review 422  
374 of analysis methods. *Emerging Themes in Epidemiol-* 423  
375 *ogy* 2017; 14(1): 12. DOI:10.1186/s12982-017-0066-2. 424  
376 URL [https://ete-online.biomedcentral.com/ 425](https://ete-online.biomedcentral.com/articles/10.1186/s12982-017-0066-2)  
377 [articles/10.1186/s12982-017-0066-2](https://ete-online.biomedcentral.com/articles/10.1186/s12982-017-0066-2). 426
- 378 12. Crisp AM, Halloran ME, Longini IM et al. Covariate- 427  
379 constrained randomization with cluster selection 428  
380 and substitution. *Clinical Trials* 2023; 20(3): 284– 429  
381 292. DOI:10.1177/17407745231160556. URL  
382 [https://journals.sagepub.com/doi/10.1177/ 427](https://journals.sagepub.com/doi/10.1177/17407745231160556)  
383 [17407745231160556](https://journals.sagepub.com/doi/10.1177/17407745231160556). 428
- 384 13. Raab GM and Butcher I. Balance in cluster randomized trials. 429  
385 *Statistics in Medicine* 2001; 20(3): 351–365. DOI:10.1002/ 430  
386 1097-0258(20010215)20:3<351::AID-SIM797>3.0.CO;2-C. 431
- 387 14. Baird S, Bohren JA, McIntosh C et al. Optimal design of 432  
388 experiments in the presence of interference. Technical report, 433  
389 The MIT Press, 2018. DOI:10.1596/29656. 434
- 390 15. McCann RS, van den Berg H, Takken W et al. Reducing 435  
391 contamination risk in cluster-randomized infectious disease- 436  
392 intervention trials. *International Journal of Epidemiol-* 437  
393 *ogy* 2018; 47(6): 2015–2024. DOI:10.1093/ije/dyy213. 438  
394 URL [https://academic.oup.com/ije/article/ 439](https://academic.oup.com/ije/article/47/6/2015/5079872)  
395 [47/6/2015/5079872](https://academic.oup.com/ije/article/47/6/2015/5079872). 440
- 396 16. Leung MP. Rate-optimal cluster-randomized designs for 441  
397 spatial interference. *The Annals of Statistics* 2022; 50(5). DOI: 442  
398 10.1214/22-AOS2224. 443
- 399 17. Cai C, Zhang X and Airoidi EM. Independent-set design 444  
400 of experiments for estimating treatment and spillover effects 445  
401 under network interference, 2023. ArXiv:2312.04026, 2312. 446
- 402 18. Leung MP. Cluster-randomized trials with cross-cluster 447  
403 interference. *Journal of the American Statistical Association* 448  
404 2025; : 1–11DOI:10.1080/01621459.2025.2566416. 449
- 405 19. Watson SI and Smith TA. Design and Analysis of Ran- 450  
406 domized Trials to Estimate Spatio-Temporally Heterogeneous 451  
407 Treatment Effects. *Journal of the American Statistical Asso-* 452  
408 *ciation* 2025; : 1–11DOI:10.1080/01621459.2025.2527424. 453  
409 URL [https://www.tandfonline.com/doi/full/ 454](https://www.tandfonline.com/doi/full/10.1080/01621459.2025.2527424)  
410 [10.1080/01621459.2025.2527424](https://www.tandfonline.com/doi/full/10.1080/01621459.2025.2527424). 455
- 411 20. Mirzaei M, Joodi G, Bogle B et al. Years of Life 456  
412 and Productivity Loss Because of Adult Sudden 457  
413 Unexpected Death in the United States. *Medical* 458  
414 *Care* 2019; 57(7): 498–502. DOI:10.1097/MLR. 459  
415 0000000000001129. URL [https://journals.lww. 460](https://journals.lww.com/00005650-201907000-00003)  
416 [com/00005650-201907000-00003](https://journals.lww.com/00005650-201907000-00003). 461
- 417 21. Gan ZSW, Choi W, Lin FC et al. Factors Underlying 462  
418 Increased Incidence of Sudden Unexpected Death in 463  
419 Rural Counties in North Carolina. *Journal of General* 464  
420 *Internal Medicine* 2019; 34(6): 815–817. DOI:10.1007/ 465  
421 s11606-018-4771-5. URL [http://link.springer. 466](http://link.springer.com/10.1007/s11606-018-4771-5)  
422 [com/10.1007/s11606-018-4771-5](http://link.springer.com/10.1007/s11606-018-4771-5). 467
- 423 22. Mounsey LA, Lin FC, Pursell I et al. Relation of Household 468  
424 Income to Incidence of Sudden Unexpected Death in Wake 469  
425 County, North Carolina. *The American Journal of Cardiology* 470  
426 2017; 119(7): 1030–1035. DOI:10.1016/j.amjcard.2016.11. 471  
427 061. URL [https://linkinghub.elsevier.com/ 472](https://linkinghub.elsevier.com/retrieve/pii/S0002914916320069)  
428 [retrieve/pii/S0002914916320069](https://linkinghub.elsevier.com/retrieve/pii/S0002914916320069). 473