

Research Paper

Factors Affecting the Geographic Distribution of West Nile Virus in Georgia, USA: 2002–2004

SAMANTHA E.J. GIBBS,¹ MICHAEL C. WIMBERLY,² MARGUERITE MADDEN,³
 JANNA MASOUR,³ MICHAEL J. YABSLEY,¹ and DAVID E. STALLKNECHT¹

ABSTRACT

The distribution of West Nile virus (WNV) is dependent on the occurrence of both susceptible avian reservoir hosts and competent mosquito vectors. Both factors can be influenced by geographic variables such as land use/landcover, elevation, human population density, physiographic region, and temperature. The current study uses geographic information systems (GIS) and logistic regression analyses to model the distribution of WNV in the state of Georgia based on a wild bird indicator system, and to identify human and environmental predictor variables that are important in the determination of WNV distribution. A database for Georgia was constructed that included (1) location points of all the avian samples tested for WNV, (2) local land use classifications, including temperature, physiographic divisions, land use/landcover, and elevation, (3) human demographic data from the U.S. Census, and (4) statistics summarizing land cover, elevation, and climate within a 1-km-radius landscape around each sample point. Logistic regression analysis was carried out using the serostatus of avian collection sites as the dependent variable. Temperature, housing density, urban/suburban land use, and mountain physiographic region were important variables in predicting the distribution of WNV in the state of Georgia. While weak, the positive correlation between WNV-antibody positive sites and the urban/suburban environment was consistent throughout the study period. The risks associated with WNV endemicity appear to be increased in urban/suburban areas and decreased in the mountainous region of the state. This information may be used in addressing regional public health needs and mosquito control programs. **Key Words:** West Nile virus—Avian—Risk—Land use—Physiographic region—Geographic information systems. *Vector-Borne Zoonotic Dis.* 6, 73–82.

INTRODUCTION

WEST NILE VIRUS (*Flaviridae*, *Flavivirus*; WNV) is a vector-borne pathogen of global importance. The geographic range of this virus has expanded since its discovery in Uganda in 1937 (Smithburn et al. 1940) and now includes Africa, Asia, Europe, North America, Central and South America, and the Caribbean (McIntosh et al. 1968, Hubalek and Halzouka 1999, Malkinson and Banet 2002, Steele et al. 2000,

OIE 2004, Cruz et al. 2005, Mattar et al. 2005, Quirin et al. 2004). The distribution of WNV is dependent on the occurrence of susceptible avian reservoir hosts and competent mosquito vectors, mosquito host preference, and availability of hosts. These factors can be influenced by geographic variables such as land use/landcover, elevation, human population density, physiographic region, and temperature.

The potential influence of environmental and social factors on WNV transmission has been

¹The Southeastern Cooperative Wildlife Disease Study, Department of Population Health, College of Veterinary Medicine, ²D.B. Warnell School of Forest Resources, and ³Center for Remote Sensing and Mapping Science, Department of Geography, University of Georgia, Athens, Georgia.

of great interest since the discovery of the virus. Taylor et al. (1956) noted differences in WNV seroprevalence between study zones along the Nile River Delta in human and hooded crow (*Corvus corone sardonius*) samples. While climatic and geologic features in these areas were not markedly different, there were significant differences in human population density and land use patterns. The increased mosquito breeding habitat and improved farmland created by irrigation in the Nile River Delta was also cited as one of the most important man-made modifications to the environment influencing the transmission of WNV (Hayes 1989).

In a more recent study, spatial analysis of WNV case distribution in the New York City area in 1999 revealed that vegetation abundance was significantly and positively associated with human WNV cases (Brownstein et al. 2002). This association was used to predict areas of greatest human risk for WNV infection; the model constructed in the study showed that the less populated suburban regions were at greatest risk. A study in the Chicago area found that risk factors associated with clusters of human cases and dead birds included vegetation, age, income, race, distance to reported WNV positive dead birds, age of housing, mosquito control activities, and geological factors (Ruiz et al. 2004). Mosquito abatement activities accounted for approximately 53% of the variation between clusters in that study. In Florida, spatial and temporal differences in periods of drought and rain were associated with variability in human WNV cases and infection of sentinel chickens (Shaman et al. 2005). The authors suggested that close proximity of birds and mosquito vectors during times of drought are responsible for increased virus transmission.

Spatial analyses have also have been applied to the surveillance data associated with dead bird submissions. Using geographic information systems (GIS), dead bird data was shown to be an effective indicator of WNV amplification and could be used to predict potential areas of high human risk at least 13 days prior to the onset of human illness in those areas (Theophilides et al. 2003). Another study applied a spatial scan statistic to detect small-area

clustering of dead birds (Mostashari et al. 2003). This information was used to predict areas of active virus transmission and served as a basis on which to target mosquito surveillance activities. In a retrospective study conducted on dead crow report data from Chicago in 2002, spatial analysis of the data showed that human cases were three times more likely to occur in areas of high early-season crow deaths (Watson et al. 2004).

Environmental conditions affecting both avian reservoir hosts and the mosquito vector populations may regulate WNV amplification. Identifying such factors will not only aid in understanding WNV epidemiology, but also will serve in predicting and possibly reducing the risk of WNV infection. In a recent study (Gibbs et al. 2006), wild birds were used as fine scale indicators of WNV transmission over the physiographic and land use variation present in Georgia. The current study builds on this work by examining these data using GIS and logistic regression analyses to predict the distribution of WNV in the state of Georgia based on a wild bird indicator system, and to identify human and environmental predictor variables that are important in the determination of WNV distribution.

METHODS

West Nile virus database construction

The serologic database used for the current study was derived from previous work in which avian serum samples ($n = 10,865$) from 70 species of birds captured throughout Georgia at approximately 200 sampling locations during the summers of 2002–2004 were tested by plaque reduction neutralization test (PRNT) for antibodies to WNV and St. Louis encephalitis virus (SLE) (Gibbs et al., 2006). Antibodies to WNV were detected in 850 (7.8%) of these samples.

A database for Georgia was constructed that included (1) location points of all the avian samples tested for WNV, (2) local land use classifications including temperature, physiographic divisions, land use/landcover, and el-

elevation, (3) human demographic data, and (4) statistics summarizing land cover, elevation, and climate within a 1-km-radius landscape around each sample point. The GIS software packages used to develop the database, extract summary statistics, and create map displays were the Environmental Systems Research Institute (ESRI) ArcView 3.3 and ArcGIS 9 (ESRI, Redlands, CA). Data layers for each environmental factor were created as ArcView shapefiles in the Universal Transverse Mercator (UTM) grid coordinate system tied to the North American Datum (NAD) of 1983. The database comprised 3 years (May–October of 2002–2004) of avian samples collected within the state of Georgia and tested for antibodies to WNV by PRNT.

Coordinates of the sample points were recorded in the field with hand-held Garmin Global Positioning System (GPS; Garmin International Inc., Olathe, KS) units in the UTM coordinate system. The sampled data points were arbitrarily distributed based on sampling opportunities and located in the two UTM zones (16 and 17) that cover the state of Georgia. Buffers of 1 km around each sampled data point were created with the ArcGIS proximity function in Arc Toolbox. Land use and demographic data falling within this 1-km buffer zone was then included in the database.

If one or more birds at a collection site were positive for WNV antibodies, that site was classified as positive. Site status was used in the analysis rather than WNV antibody status of individual birds in an effort to avoid bias introduced by species differences in prevalence and geographic distribution (Gibbs et al. 2006).

Logistic regression

Explanatory variables. Explanatory variables used in logistic regression modeling included land use/land cover, physiographic region, elevation, minimum and maximum temperatures for January and February, and 2000 census data for the state of Georgia, which included human population per acre and per hectare, and housing density per acre and per hectare.

A landscape-level land use/landcover of

Georgia dataset was created by the University of Georgia, Institute of Ecology, Natural Resources Spatial Analysis Laboratory and obtained from the Georgia GIS Clearinghouse (www.gis.state.ga.us). The Landsat Landcover (18 class) dataset, dated 1998, was produced from Landsat Thematic Mapper (TM) satellite imagery of 30-m spatial resolution. Based on the requirements of this project, we reduced the number of classes to five: urban-suburban, forest, agriculture, wetland, and other. This dataset was converted to an ArcView grid with a cell size of 30×30 m. A second dataset of local land use characteristics for each sampling site was created based on field observations recorded during avian sample collection; this provided a finer scale assessment of land use than was available using the Landsat Landcover dataset.

The physiographic divisions dataset was derived from the U.S. Geological Survey (USGS) "Physiographic Provinces" ESRI Export File (e00), dated 1992 and based on a 1:7,000,000-scale map. These data were obtained from the Georgia GIS Clearinghouse. Physiographic divisions are areas having similar topography, rock types, geology, and geomorphic history as defined by USGS. The original data in the Lambert Conic Conformal coordinate system were reprojected to the UTM coordinate system, NAD 83. We used four major physiographic divisions: mountains, piedmont, coastal plain, and coastal. These data were also converted to an ArcView grid with a cell size of 30×30 m.

The source for the elevation dataset was the USGS 1:24,000 scale National Elevation Dataset (NED), dated 1999. The resolution for this data set was 30×30 m. Data were obtained from the Georgia GIS Clearinghouse. The file format was ERDAS Imagine (.img) in a geographical projection (latitude and longitude). The data were converted to an ArcView grid file and reprojected to the UTM coordinate system, NAD83.

The source for temperature datasets of monthly mean maximum and minimum temperature for January and February was acquired from the National Climatic Data Center (NCDC; www.ncdc.noaa.gov/oa/ncdc.html). Data from the weather stations in nine Georgia

climatic divisions were combined with data from stations in 11 adjacent divisions of the neighboring states (totaling approximately 300 stations) in order to insure an accurate and complete spatial interpolation. A Kriging interpolation method was employed in ArcView to create a continuous surface of temperature data for the study area. The resulting triangulated irregular networks (TINs) of temperatures covering the state of Georgia were created for January and February of 2002, 2003, and 2004. The TINs were subsequently converted to GRID format for spatial analysis in ArcView. Grids for temperature were created with a cell size of 100×100 m. Four grids were created for each year—January mean minimum temperature, January mean maximum temperature, February mean minimum temperature, and February mean maximum temperature.

Housing density was computed using U.S. Census data from 2000. Block-level data for the state of Georgia was queried from the CensusCD 2000/Short Form Blocks dataset (GeoLytics, Inc., East Brunswick, NJ). Census blocks are the smallest spatial units at which census data is released, and their sizes vary depending on population density. They can be as small as a city block in urban areas, or as large as 400 acres in rural areas. The total count of housing units within each census block was normalized by the acreage of each block to generate housing density.

An overlay analysis function in ArcView, clip, was used to extract the grid data cells of environmental factors within each 1-km-radius buffer area surrounding the sample points. The ArcView Map Calculator function was then used to extract and summarize the average values of environmental factors within a 1-km buffer area surrounding each sample data point. Mean maximum and minimum temperature for January and February, physiographic divisions, land use/landcover, and elevation were thus summarized. Data for temperature and elevation were summarized as a weighted mean for each buffer area, while land use/landcover and physiographic divisions were calculated as the percent area covered by each class in the buffer area. Housing density was obtained from the census block within which

each sample point fell. Categorical variables, including physiographic region and local land use, were each coded as a set of (0, 1) dummy variables corresponding to each physiographic region or land use class.

Statistical analysis

Logistic regression analysis was carried out using S-Plus 6.1 (Insightful Corporation, Seattle, WA) using the serostatus of avian collection sites as the dependent variable. A forward stepwise procedure with a $p < 0.05$ was used to determine which environmental variables maximized the fit of the statistical model based on our data. Accuracy (percent of testing sites correctly classified), sensitivity (percent of positive testing sites correctly classified), and specificity (percent of negative testing sites correctly classified) were computed for the model. Also, the area under the receiver operating characteristic curve (AUC ROC) and the max rescaled R^2 were calculated as indices of the fit of the model (Fielding and Bell 1997, Nagelkerke 1991).

RESULTS

Avian samples were collected at sites within 151 counties of the state of Georgia (Fig. 1). The sampling sites were distributed throughout all land use types and physiographic regions present in the state. Background data for the avian samples on which the site data were based are shown in Table 1.

For the 2002 data, four variables were significantly related to WNV serostatus (Table 2). The probability of WNV being present in an area increased with field-observed urban-suburban land use and minimum January temperature. A unimodal response to the natural logarithm of housing density was observed, and the second-degree polynomial term was included in the final model. The probability of a site being WNV positive was highest at densities of approximately one housing unit per 10 acres, and was lower both in more rural areas (<1 housing unit per 10 acres) and in more heavily populated areas (>1 housing unit per 10 acres). The 2003 model included only the field observed urban-suburban land use vari-

WNV DISTRIBUTION IN GEORGIA

77

able, which was positively associated with WNV presence. In the 2004 model, the probability of WNV positive sites was lower in the mountains than in other physiographic regions. As with the 2002 model, the second-degree polynomial term for the natural logarithm of housing density indicated that the probability of a site containing WNV seropositive birds was highest at intermediate housing densities of approximately one housing unit per 10 acres.

The accuracy, sensitivity, specificity, maximum rescaled R^2 , and area under the receiver operating characteristic curve (AUC ROC) of the 2002 model were much greater than in the 2003 and 2004 models (Table 3).

The serologic status of collection sites that were sampled in more than one year are listed in Table 4. Approximately half of the sites which were initially negative changed status to

positive upon re-sampling. Few sites initially found positive changed to a negative status.

DISCUSSION

The results of this study illustrate the widespread distribution of WNV in the state of Georgia in just three years after introduction. Antibodies against WNV were found for each sample year in both adult and hatch year birds, indicating that the virus was able to over-winter and become endemic in the state (Gibbs et al. 2006). This finding was supported by dead bird and mosquito surveillance data (SCWDS, unpublished data). As demonstrated by the poor accuracy of the logistic regression models in the last two years of the study, environmental and demographic variables became less




FIG. 1. Distribution of West Nile virus antibody positive and negative sites in the state of Georgia for 2002–2004.

TABLE 1. WEST NILE VIRUS ANTIBODY PREVALENCE IN AVIAN SAMPLES BY SITE

	No. sites	Avg. number of birds sampled per site (range)	No. sites positive (% pos.)
2002			
Land use			
Ag	63	15 (1–71)	13 (20.6)
F	41	10 (1–56)	8 (19.5)
U/S	63	14 (1–165)	15 (23.8)
W	11	14 (1–37)	1 (9.0)
Phys region			
C	16	14 (1–37)	2 (12.5)
CP	74	18 (1–56)	24 (32.4)
M	32	9 (1–74)	3 (9.4)
P	56	17 (1–288)	8 (14.3)
2003			
Land use			
Ag	74	21 (2–72)	38 (51.4)
F	56	16 (1–65)	27 (48.2)
U/S	79	16 (1–137)	49 (62.0)
W	6	16 (4–28)	2 (33.3)
Phys region			
C	21	14 (3–32)	9 (42.9)
CP	83	20 (1–53)	49 (59.0)
M	43	13 (2–65)	19 (44.2)
P	68	18 (1–137)	39 (57.4)
2004			
Land use			
Ag	43	23 (2–47)	32 (74.4)
F	56	19 (2–83)	35 (62.5)
U/S	112	17 (1–139)	70 (62.5)
W	11	17 (5–41)	2 (18.2)
Phys region			
C	16	16 (4–30)	10 (62.5)
CP	91	18 (1–53)	64 (70.3)
M	29	16 (1–37)	13 (44.8)
P	86	21 (1–139)	51 (60.0)

Ag, agriculture; F, forest; U/S, urban/suburban; W, wetland; C, coastal; CP, coastal plain; M, mountain; P, piedmont; Phys region, physiographic region; No., number; Avg., average; pos., positive.

important in determining the distribution of the virus as time progressed. This is consistent with logistic regression findings based on data from individual avian samples in which the local land use variable was less important in the 2003 than 2002 model, and was not included at all in the 2004 model (Gibbs et al. 2006).

The thorough coverage of WNV across the Georgia landscape was most likely facilitated by the presence of several competent vectors. *Culex quinquefasciatus*, the primary vector in the state, as well as *Cx. nigripalpus*, *Cx. restauns*, and *Cx. salinarius*, each include Georgia in their distributions (Darsie and Ward 2005). The different behavioral characteristics of these mosquitoes, including host and habitat preference, allow for transmission in a diversity of envi-

ronments. *Cx. quinquefasciatus* may be found in abundance in human modified habitats such as residential areas (Reisen 1992). Such modifications include creation of mosquito habitat in flower pots, used tires, flooded basements, sewage treatment areas, and water-catchment basins in housing developments. There are few areas within the state of Georgia that are not heavily impacted by human activities.

Sites testing positive for WNV antibodies in 2002 may represent the areas to which WNV was first introduced in 2001. Serologic data from 2002 revealed two loose foci of positive sites, one in the coastal plain and one in the metro Atlanta area (Gibbs et al. 2006). A similar spatial distribution was also observed in dead bird surveillance data from 2001 (Gibbs et

WNV DISTRIBUTION IN GEORGIA

79

TABLE 2. LOGISTIC REGRESSION MODELS FOR PREDICTING THE DISTRIBUTION OF WEST NILE VIRUS IN THE STATE OF GEORGIA BASED ON A WILD BIRD INDICATOR SYSTEM

Year	Variable	Coefficient	Wald statistic	p
2002	Minimum January temperature	0.02	3.31	0.0009
	log(housing/hectare + 0.01)	-1.62	-3.28	0.0010
	[log(housing/hectare + 0.01)] ²	-0.30	-3.11	0.0019
	Urban-suburban land use based on field observation	1.74	3.25	0.0012
2003	Urban-suburban land use based on field observation	0.61	2.11	0.0350
2004	Mountain physiographic region	-0.01	-2.72	0.0066
	log(housing/hectare + 0.01)	-0.21	-1.56	0.1190
	[log(housing/hectare + 0.01)] ²	-0.09	-2.29	0.0223

al. 2006). In the following transmission seasons, the range of the virus might have then extended through local bird movements rather than being reintroduced by migratory species. The effects of the land use and physiographic region on site serologic status during 2002 may have been enhanced due to a more localized distribution associated with initial introduction. The early distribution may also reflect the importance of microhabitat rather than broader habitat patterns within the 1-km-radius area surrounding the capture site in determining avian WNV exposure. It may also be that mosquito breeding habitat is locally important and could, in future, be added as a local variable.

Most collection sites were chosen for the presence of a factor which concentrated bird populations, thus increasing sample size. Many of these included sites with bird feeders. The practice of capturing birds around feeders and

taking advantage of nuisance bird removal programs may have influenced distribution data.

The overall fit of the 2004 model was better than the 2003 model, but neither predicted the presence of WNV as accurately as the 2002 model. Housing density was a strong predictor of WNV presence in 2002, with higher numbers of positive sites situated in areas with intermediate housing densities. While not as strongly correlated with WNV positive sites as it was in 2002, the housing density was also included in the model for 2004. Together with the inclusion of urban-suburban land use data (based on field observation) in 2003, this information supports the contention that human activities in the urban/suburban landscape provide reservoir host and vector habitats suitable for efficient WNV transmission.

While weak, the positive correlation between WNV antibody positive sites and the urban/suburban environment was consistent throughout the study period. The human impact on disease ecology has been studied intensively as humans continue to expand and modify their environment. In an assessment of emerging pathogens of wildlife in North America between 1998 and 2000, the majority of outbreaks were linked to human activities (Dobson and Foufopoulos 2001). As demonstrated by a number of arboviruses, disease emergence is most often related to human activities that increase disease vector habitats or change the density of non-human vertebrates involved in virus amplification (birds in the case of WNV) (Shope 1997, Mackenzie et al. 2004, Anonymous 1994). Urbanization and deforestation have been linked to emergence of arboviruses such as Rift Valley fever, SLE, and dengue (Wilson 1994). De-

TABLE 3. COMPARISON OF THREE MODELS CONSTRUCTED FOR PREDICTING DISTRIBUTION OF WEST NILE VIRUS IN THE STATE OF GEORGIA BASED ON A WILD BIRD INDICATOR SYSTEM

Analysis ^a	2002	2003	2004
Accuracy (%)	83.1	55.6	64.9
Sensitivity (%)	76.9	64.6	65.8
Specificity (%)	83.6	50.4	58.6
Maximum rescaled R ²	0.239	0.028	0.072
AUC ROC	0.751	0.57	0.622

^aA sampling site was considered positive if ≥ 1 bird tested positive for West Nile Virus antibodies by plaque reduction neutralization test (PRNT).

AUC ROC, area under the receiver operating characteristic curve.

TABLE 4. WEST NILE VIRUS SEROLOGIC STATUS OF COLLECTION SITES WHICH WERE SAMPLED DURING MORE THAN ONE TRANSMISSION SEASON

Years	No. sites re-sampled	Site status				Variable
		Negative to negative	Positive to positive	Negative to positive	Positive to negative	
2002–2003	64	21	13	22	8	—
2003–2004	81	14	43	11	8	—
2002–2004	35	10	7	17	1	—
2002–2003–2004	27	3	6	12	1	5 ^a

^aVariable: negative to positive to negative, or positive to negative to positive.
No., number.

spite an abundance of review literature on the topic, minimal data are available to confirm the impact of human activities on disease epidemiology (Kuiken et al. 2003). The current study provides some of these data, as well as insight into the potential of a human altered environment, specifically the urban/suburban environment, to support vector borne disease. Human activities not only support mosquito populations, but also provide food, nesting, and roosting habitat for both native and introduced birds.

The importance of minimum January temperature in the 2002 model probably reflects the dependence of WNV transmission on mosquito vectors. In future studies, examination of year-round data might help to further explain maintenance of WNV in an area from year to year. A variable for surface water was not included in this analysis as the primary mosquito species involved in WNV transmission in Georgia tend not to use large bodies of water (rivers and lakes) for breeding. The ephemeral water sources which generally serve as breeding sites for these species would not be accurately represented by the 1998 landcover dataset used in this study. The inclusion of mountain physiographic region in the 2004 model also emphasizes the importance of elevation, temperature, and physiographic region in WNV epidemiology. These three variables are interrelated in Georgia, and in combination appear to limit the transmission of WNV. Reasons for this limitation probably relate to lower temperatures in mountainous regions, consequent decreases in mosquito abundance, and differences in avian species composition.

The changing serologic status of re-sampled sites in this study agrees with antibody prevalence data obtained from individual birds in this dataset (Gibbs et al. 2006); antibody prevalence for all species increased from 5.2% to 7.3% to 10.1% in 2002, 2003, and 2004, respectively. The consistent rise in seroprevalence between years reflects the increasing geographic distribution of WNV taking place during the study. Approximately 30% of the sites initially negative converted to positive sites, however, only 8.7% went from positive to negative. This is surprising considering the low prevalence and small sample size at some sites.

As demonstrated by this work, WNV is distributed throughout the state of Georgia. West Nile virus poses a health risk to humans, livestock, and wildlife in all physiographic regions and land use types; the challenge we experienced in developing a model with high sensitivity and specificity for the data set reflects this broad distribution. The data were grouped according to sampling site rather than by individual serum sample in an effort to decrease bias potentially introduced by differences in susceptibility and antibody formation in avian species. Sites where the results were based on only one species (such as the Canada goose collections) were removed from the spatial analysis to help address sampling bias.

Modeling based on site status (as used in this study) will also work in future studies; however, a targeted approach towards sites with good avian indicators (such as the northern cardinal) would be a better approach. Seroprevalence data from individual samples could then be included in the model, making it more ac-

curate in defining the influence environmental and demographic factors have on WNV distribution. The importance of these factors will most likely change as the environment becomes increasingly impacted by human population expansion.

The trends observed both in this study and work conducted on WNV antibody prevalence in avian species (Gibbs et al. 2006) suggest that the virus will continue to circulate in the environment at endemic levels. The risks associated with endemicity appear to be increased in urban/suburban areas and decreased in the mountainous region of the state. This information may be used in addressing regional public health needs and mosquito control programs; priority should be placed on campaigns aimed at decreasing man-made mosquito habitats in urban/suburban areas.

ACKNOWLEDGMENTS

We wish to thank Ben Wilcox, Douglas Hoffman, Darrell Kavanaugh, Page Luttrell, Lynn Lewis, Clay George, Chris Sewell, Lisa Balmert, Casey Sanders, Robbie Edalgo, Alecia Byrd, Daniel Brown, Andrea Howey, Blythe Layng, Katie Stumpf, John Leonard, Jo Williams, Tessa Smith, and Megan Parks for field expertise. This study was primarily supported by the Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention's "Applied Research in Emerging Infections—Investigations of West Nile Virus" grant program (contract U50/CCU420515-01). Additional support was provided by the Federal Aid to Wildlife Restoration Act (50 Stat. 917) and through sponsorship from fish and wildlife agencies in Alabama, Arkansas, Florida, Georgia, Kansas, Kentucky, Louisiana, Maryland, Mississippi, Missouri, North Carolina, Oklahoma, Puerto Rico, South Carolina, Tennessee, Virginia, and West Virginia.

REFERENCES

- Anonymous. Vector-borne emerging pathogens: introduction. In: Wilson, ME, Levins, R, Spielman, A, eds. *Disease in Evolution: Global Changes and Emergence of Infectious Diseases*. Ann NY Acad Sci 1994; 740:123–125.
- Brownstein, JS, Rosen, H, Purdy, D, et al. Spatial analysis of West Nile virus: rapid risk assessment of an introduced vector-borne zoonosis. *Vector Borne Zoonotic Dis* 2002; 2:157–164.
- Cruz, L, Cardenas, VM, Abarca, M, et al. Serological evidence of West Nile virus activity in El Salvador. *Am J Trop Med Hyg* 2005; 72:612–615.
- Darsie, RF, Ward, RA. *Identification and Geographical Distribution of Mosquitoes of North America, North of Mexico*. Gainesville, FL: University Press of Florida, 2005.
- Dobson, A, Foufopoulos, J. Emerging infectious pathogens of wildlife. *Phil Trans R Soc Lond* 2001; 356:1001–1012.
- Fielding, AH, Bell, JF. A review of methods for the assessment of prediction errors in conservation presence/absence models. *Environ Conserv* 1997; 24:38–49.
- Gibbs, SEJ, Allison, AB, Yabsley, MJ, Mead, DG, Wilcox, BR, Stallknecht, DE. West Nile virus antibodies in avian species of Georgia, USA: 2000–2004. *Vector Borne Zoonotic Dis* 2006; 1:57–72 (this issue).
- Hayes, CG. West Nile Fever. In: Monath, TP, ed. *The Arboviruses: Epidemiology and Ecology*, Vol. 5. Boca Raton, FL: CRC Press; 1989:59–88.
- Hubalek, Z, Halouzka, J. West Nile fever—a reemerging mosquito-borne viral disease in Europe. *Emerg Infect Dis* 1999; 5:643–650.
- Kuiken, T, Fouchier, R, Rimmelzwaan, G, et al. Emerging viral infections in a rapidly changing world. *Curr Opin Biotechnol* 2003; 14:641–646.
- MacKenzie, JS, Gubler, DJ, Petersen LR. Emerging flaviviruses: the spread and resurgence of Japanese encephalitis, West Nile and dengue viruses. *Nat Med* 2004; 10:S98–S109.
- Malkinson, M, Banet, C. The role of birds in the ecology of West Nile virus in Europe and Africa. *Curr Top Microbiol Immunol* 2002; 267:309–322.
- Mattar, S, Edwards, E, Laguado, J, et al. West Nile virus antibodies in Columbian horses. *Emerg Infect Dis* 2005; 11:1497–1498.
- McIntosh, BM, McGillivray, GM, Dickinson, DB, et al. Ecological studies on Sindbis and West Nile viruses in South Africa. IV: Infection in a wild avian population. *S Afr J Medical Sci* 1968; 33:105–112.
- Mostashari, F, Kulldorff, M, Hartman, JJ, et al. Dead bird clusters as an early warning system for West Nile virus activity. *Emerg Infect Dis* 2003; 9:641–646.
- Nagelkerke, NJD. A note on a general definition of the coefficient of determination. *Biometrika* 1991; 78:691–692.
- OIE, World Organization for Animal Health. West Nile fever in Belize: in October 2003. *Dis Inform* 2004; 17:10.
- Quirin, R, Salas, M, Zientara, S, et al. West Nile virus, Guadeloupe. *Emerg Infect Dis* 2004; 10:706–708.
- Reisen, WK, Milby, MM, Presser, SB, et al. Ecology of mosquitoes and St. Louis encephalitis virus in the Los Angeles Basin of California, 1987–1990. *J Med Entomol* 1992; 29:582–598.
- Ruiz, MO, Tedesco, C, McTighe, TJ, et al. Environmental

- and social determinants of human risk during a West Nile virus outbreak in the greater Chicago area, 2002. *Int J Health Geogr* 2004; 3:8.
- Shaman, J, Day, JF, Stieglitz, M. Drought-induced amplification and epidemic transmission of West Nile virus in southern Florida. *J Med Entomol* 2005; 42:134–141.
- Shope, RE. Emergence of arbovirus diseases following ecological modifications: epidemiological consequences. In: Sluzzo, JF, Dodet, B, eds. *Factors in the Emergence of Arbovirus Diseases*. Paris: Elsevier; 1997:19–22.
- Smithburn, KC, Hughs, TP, Burke, AW, et al. A neurotropic virus isolated from the blood of a native of Uganda. *Am J Trop Med Hyg* 1940; 20:471–92.
- Steele, KE, Linn, MJ, Schoepp, RJ, et al. Pathology of fatal West Nile virus infections in native and exotic birds during the 1999 outbreak in New York City, New York. *Vet Pathol* 2000; 37:208–224.
- Taylor, RM, Work, TH, Hurlbut, HS, et al. A study of the ecology of West Nile virus in Egypt. *Am J Trop Med Hyg* 1956; 5:579–620.
- Theophilides, CN, Ahearn, SC, Grady, S, et al. Identifying West Nile virus risk areas: the dynamic continuous-area space-time system. *Am J Epidemiol*. 2003; 157:843–854.
- Watson, JT, Jones, RC, Gibbs, K, et al. Dead crow reports and location of human West Nile virus cases, Chicago, 2002. *Emerg Infect Dis* 2004; 10:938–940.
- Wilson, ML. Rift Valley fever virus ecology and the epidemiology of disease emergence In: Wilson, ME, Levins, R, Spielman, A, eds. *Disease in Evolution: Global Changes and Emergence of Infectious Diseases*. Ann NY Acad Sci, 1994; 740:169–180.

Address reprint requests to:

Dr. Samantha E.J. Gibbs

Wildlife Health Building

Southeastern Cooperative Wildlife Disease Study

College of Veterinary Medicine

University of Georgia

Athens, GA 30602

E-mail: sgibbs@vet.uga.edu

This article has been cited by:

1. Justin K. Davis, Geoffrey P. Vincent, Michael B. Hildreth, Lon Kightlinger, Christopher Carlson, Michael C. Wimberly. 2018. Improving the prediction of arbovirus outbreaks: A comparison of climate-driven models for West Nile virus in an endemic region of the United States. *Acta Tropica* **185**, 242-250. [[Crossref](#)]
2. Meredith E Kernbach, Richard J Hall, Nathan D Burkett-Cadena, Thomas R Unnasch, Lynn B Martin. 2018. Dim light at night: physiological effects and ecological consequences for infectious disease. *Integrative and Comparative Biology* **16**. . [[Crossref](#)]
3. Minjung Seo, Ji Won Ha. 2018. Effective surface-enhanced Raman scattering of randomly branched gold nano-urchins with Rhodamine 6G as Raman reporters. *Microchemical Journal* **140**, 47-51. [[Crossref](#)]
4. Istifanus J Waziri, Junaidu Kabir, Jacob K P Kwaga, Patrick Nguku. 2018. Serosurvey of West Nile virus in household-reared pigeons in Bauchi metropolis, Nigeria. *Transactions of The Royal Society of Tropical Medicine and Hygiene* **112**:4, 169-174. [[Crossref](#)]
5. Annelise Tran, Grégory L'Ambert, Gilles Balança, Sophie Pradier, Vladimir Grosbois, Thomas Balenghien, Thierry Baldet, Sylvie Lecollinet, Agnès Leblond, Nicolas Gaidet-Drapier. 2017. An Integrative Eco-Epidemiological Analysis of West Nile Virus Transmission. *EcoHealth* **14**:3, 474-489. [[Crossref](#)]
6. Mohamed Sallam, Sarah Michaels, Claudia Riegel, Roberto Pereira, Wayne Zipperer, B. Lockaby, Philip Koehler. 2017. Spatio-Temporal Distribution of Vector-Host Contact (VHC) Ratios and Ecological Niche Modeling of the West Nile Virus Mosquito Vector, *Culex quinquefasciatus*, in the City of New Orleans, LA, USA. *International Journal of Environmental Research and Public Health* **14**:8, 892. [[Crossref](#)]
7. Alexander T Ciota. 2017. West Nile virus and its vectors. *Current Opinion in Insect Science* **22**, 28-36. [[Crossref](#)]
8. Anne Seltmann, Victor M. Corman, Andrea Rasche, Christian Drosten, G?bor ?. Czirj?k, Henry Bernard, Matthew J. Struebig, Christian C. Voigt. 2017. Seasonal Fluctuations of Astrovirus, But Not Coronavirus Shedding in Bats Inhabiting Human-Modified Tropical Forests. *EcoHealth* **14**:2, 272-284. [[Crossref](#)]
9. Tom Swan, Jon Harding. 2017. The distribution and occurrence of mosquito larvae (Diptera: Culicidae) in the Tongatapu Island Group, Kingdom of Tonga. *Austral Entomology* **56**:2, 160-168. [[Crossref](#)]
10. Clive McAlpine, Grant Brearley, Jonathan Rhodes, Adrian Bradley, Greg Baxter, Leonie Seabrook, Daniel Lunney, Yan Liu, Manuelle Cottin, Andrew G. Smith, Peter Timms. 2017. Time-delayed influence of urban landscape change on the susceptibility of koalas to chlamydiosis. *Landscape Ecology* **32**:3, 663-679. [[Crossref](#)]
11. Mohamed F. Sallam, Rui-De Xue, Roberto M. Pereira, Phillip G. Koehler. 2016. Ecological niche modeling of mosquito vectors of West Nile virus in St. John's County, Florida, USA. *Parasites & Vectors* **9**:1. . [[Crossref](#)]
12. Graeme Lockaby, Navideh Noori, Wayne Morse, Wayne Zipperer, Latif Kalin, Robin Governo, Rajesh Sawant, Matthew Ricker. 2016. Climatic, ecological, and socioeconomic factors associated with West Nile virus incidence in Atlanta, Georgia, U.S.A. *Journal of Vector Ecology* **41**:2, 232-243. [[Crossref](#)]
13. Nicholas Karam Skaff, Kendra Spence Cheruvilil. 2016. Fine-scale wetland features mediate vector and climate-dependent macroscale patterns in human West Nile virus incidence. *Landscape Ecology* **31**:7, 1615-1628. [[Crossref](#)]
14. L. Shand, W. M. Brown, L. F. Chaves, T. L. Goldberg, G. L. Hamer, L. Haramis, U. Kitron, E. D. Walker, M.O. Ruiz. 2016. Predicting West Nile Virus Infection Risk From the Synergistic Effects of Rainfall and Temperature. *Journal of Medical Entomology* **53**:4, 935-944. [[Crossref](#)]
15. Alexandra Gulachenski, Bruno Ghersi, Amy Lesen, Michael Blum. 2016. Abandonment, Ecological Assembly and Public Health Risks in Counter-Urbanizing Cities. *Sustainability* **8**:5, 491. [[Crossref](#)]
16. Benita Y. Tam, Leonard J. S. Tsuji. 2016. West Nile virus in American crows (*Corvus brachyrhynchos*) in Canada: projecting the influence of climate change. *GeoJournal* **81**:1, 89-101. [[Crossref](#)]
17. T. Luke George, Ryan J. Harrigan, Joseph A. LaManna, David F. DeSante, James F. Saracco, Thomas B. Smith. 2015. Persistent impacts of West Nile virus on North American bird populations. *Proceedings of the National Academy of Sciences* **112**:46, 14290-14294. [[Crossref](#)]
18. Jonathan F. Day, Walter J. Tabachnick, Chelsea T. Smartt. 2015. Factors That Influence the Transmission of West Nile Virus in Florida. *Journal of Medical Entomology* **52**:5, 743-754. [[Crossref](#)]
19. Benjamin Roche, Serge Morand, Eric Elguero, Thomas Balenghien, Jean-François Guégan, Nicolas Gaidet. 2015. Does host receptivity or host exposure drives dynamics of infectious diseases? The case of West Nile Virus in wild birds. *Infection, Genetics and Evolution* **33**, 11-19. [[Crossref](#)]
20. Matteo Marcantonio, Annapaola Rizzoli, Markus Metz, Roberto Rosà, Giovanni Marini, Elizabeth Chadwick, Markus Neteler. 2015. Identifying the Environmental Conditions Favours West Nile Virus Outbreaks in Europe. *PLOS ONE* **10**:3, e0121158. [[Crossref](#)]
21. S. Paz. 2015. Climate change impacts on West Nile virus transmission in a global context. *Philosophical Transactions of the Royal Society B: Biological Sciences* **370**:1665, 20130561-20130561. [[Crossref](#)]
22. Benita Y. Tam, Ian Martin, Leonard J. S. Tsuji. 2014. Geospatial analysis between the environment and past incidences of West Nile virus in bird specimens in Ontario, Canada. *GeoJournal* **79**:6, 805-817. [[Crossref](#)]
23. Andrew Curtis, Xinyue Ye, Elyse Heob, Joseph Targhetta, Vence Salvato, Martin Reyna, Rudy Bueno, Louisa Holmes. 2014. A comparison of three approaches to identify West Nile Virus mosquito space-time hotspots in the Houston Vicinity for the period 2002-2011. *Applied Geography* **51**, 58-64. [[Crossref](#)]
24. George Valiakos, Konstantinos Papaspyropoulos, Alexios Giannakopoulos, Periklis Birtsas, Sotirios Tsiodras, Michael R. Hutchings, Vassiliki Spyrou, Danai Pervanidou, Labrini V. Athanasiou, Nikolaos Papadopoulos, Constantina Tsokana, Agoritsa Baka, Katerina Manolakou, Dimitrios

Chatzopoulos, Marc Artois, Lisa Yon, Duncan Hannant, Liljana Petrovska, Christos Hadjichristodoulou, Charalambos Billinis. 2014. Use of Wild Bird Surveillance, Human Case Data and GIS Spatial Analysis for Predicting Spatial Distributions of West Nile Virus in Greece. *PLoS ONE* 9:5, e96935. [Crossref]

25. Sonnleitner Sissy Therese, Lundström Jan, Baumgartner Raphaela, Simeoni Josef, Schennach Harald, Zelger Roland, Prader Angelika, Schmutzhard Erich, Nowotny Norbert, Walder Gernot. 2014. Investigations on California Serogroup Orthobunyaviruses in the Tyrols: First Description of Tahyna Virus in The Alps. *Vector-Borne and Zoonotic Diseases* 14:4, 272-277. [Abstract] [Full Text] [PDF] [PDF Plus]
26. Qihao Weng, Bing Xu, Xuefei Hu, Hua Liu. 2014. Use of earth observation data for applications in public health. *Geocarto International* 29:1, 3-16. [Crossref]
27. Rebecca S. Levine, Daniel G. Mead, Uriel D. Kitron. 2013. Limited Spillover to Humans from West Nile Virus Viremic Birds in Atlanta, Georgia. *Vector-Borne and Zoonotic Diseases* 13:11, 812-817. [Abstract] [Full Text] [PDF] [PDF Plus]
28. Grant Brearley, Jonathan Rhodes, Adrian Bradley, Greg Baxter, Leonie Seabrook, Daniel Lunney, Yan Liu, Clive McAlpine. 2013. Wildlife disease prevalence in human-modified landscapes. *Biological Reviews* 88:2, 427-442. [Crossref]
29. Carolina Vergara Cid, Elizabet Lilia Estallo, Walter Ricardo Almirón, Marta Silvia Contigiani, Lorena Ivana Spinsanti. 2013. Landscape determinants of Saint Louis encephalitis human infections in Córdoba city, Argentina during 2010. *Acta Tropica* 125:3, 303-308. [Crossref]
30. Shlomit Paz, Dan Malkinson, Manfred S. Green, Gil Tsioni, Anna Papa, Kostas Danis, Anca Sirbu, Cornelia Ceianu, Krisztalovics Katalin, Emőke Ferenczi, Herve Zeller, Jan C. Semenza. 2013. Permissive Summer Temperatures of the 2010 European West Nile Fever Upsurge. *PLoS ONE* 8:2, e56398. [Crossref]
31. Shannon LaDeau, Paul Leisnham, Dawn Biehler, Danielle Bodner. 2013. Higher Mosquito Production in Low-Income Neighborhoods of Baltimore and Washington, DC: Understanding Ecological Drivers and Mosquito-Borne Disease Risk in Temperate Cities. *International Journal of Environmental Research and Public Health* 10:4, 1505. [Crossref]
32. Shlomit Paz, Jan Semenza. 2013. Environmental Drivers of West Nile Fever Epidemiology in Europe and Western Asia—A Review. *International Journal of Environmental Research and Public Health* 10:8, 3543. [Crossref]
33. Esra Ozdenerol, Gregory Taff, Cem Akkus. 2013. Exploring the Spatio-Temporal Dynamics of Reservoir Hosts, Vectors, and Human Hosts of West Nile Virus: A Review of the Recent Literature. *International Journal of Environmental Research and Public Health* 10:11, 5399. [Crossref]
34. Michael Wimberly, Paolla Giacomo, Lon Kightlinger, Michael Hildreth. 2013. Spatio-Temporal Epidemiology of Human West Nile Virus Disease in South Dakota. *International Journal of Environmental Research and Public Health* 10:11, 5584. [Crossref]
35. Theodore G. Andreadis. 2012. The Contribution of Culex pipiens Complex Mosquitoes to Transmission and Persistence of West Nile Virus in North America. *Journal of the American Mosquito Control Association* 28:4s, 137-151. [Crossref]
36. Alexander Carnes, Yelena Ogneva-Himmelberger. 2012. Temporal Variations in the Distribution of West Nile Virus Within the United States; 2000–2008. *Applied Spatial Analysis and Policy* 5:3, 211-229. [Crossref]
37. Michael C. Parrish, Jeffrey Hepinstall-Cymerman. 2012. Associations between multiscale landscape characteristics and breeding bird abundance and diversity across urban-rural gradients in Northeastern Georgia, USA. *Urban Ecosystems* 15:3, 559-580. [Crossref]
38. John P. DeGroot, Ramanathan Sugumaran. 2012. National and Regional Associations Between Human West Nile Virus Incidence and Demographic, Landscape, and Land Use Conditions in the Conterminous United States. *Vector-Borne and Zoonotic Diseases* 12:8, 657-665. [Abstract] [Full Text] [PDF] [PDF Plus]
39. Hua Liu, Qihao Weng. 2012. Environmental Factors and Risk Areas of West Nile Virus in Southern California, 2007–2009. *Environmental Modeling & Assessment* 17:4, 441-452. [Crossref]
40. Patrick T. Vander Kelen, Joni A. Downs, Nathan D. Burkett-Cadena, Christy L. Ottendorfer, Kevin Hill, Stephen Sickerman, José Hernandez, Joseph Jinright, Brenda Hunt, John Lusk, Victor Hoover, Keith Armstrong, Robert S. Unnasch, Lillian M. Stark, Thomas R. Unnasch. 2012. Habitat Associations of Eastern Equine Encephalitis Transmission in Walton County Florida. *Journal of Medical Entomology* 49:3, 746-756. [Crossref]
41. Christine W. Hockett, Lon Kightlinger, Ting-Wu Chuang, Michael C. Wimberly. 2012. Landscape-Level Spatial Patterns of West Nile Virus Risk in the Northern Great Plains. *The American Journal of Tropical Medicine and Hygiene* 86:4, 724-731. [Crossref]
42. Hua Liu, Qihao Weng. 2012. Enhancing temporal resolution of satellite imagery for public health studies: A case study of West Nile Virus outbreak in Los Angeles in 2007. *Remote Sensing of Environment* 117, 57-71. [Crossref]
43. S. T. Sonnleitner, J. Simeoni, E. Schmutzhard, M. Niedrig, F. Ploner, H. Schennach, M. P. Dierich, G. Walder. 2012. Absence of indigenous specific West Nile virus antibodies in Tyrolean blood donors. *European Journal of Clinical Microbiology & Infectious Diseases* 31:1, 77-81. [Crossref]
44. Eva A. Buckner, Mark S. Blackmore, Stephen W. Golladay, Alan P. Covich. 2011. Weather and landscape factors associated with adult mosquito abundance in southwestern Georgia, U.S.A. *Journal of Vector Ecology* 36:2, 269-278. [Crossref]
45. Andrew D. Haddow, Danae Bixler, Amy J. Schuh. 2011. The Demographic and Socioeconomic Factors Predictive for Populations at High-Risk for La Crosse Virus Infection in West Virginia. *PLoS ONE* 6:9, e25739. [Crossref]
46. Shannon L. LaDeau, Catherine A. Calder, Patrick J. Doran, Peter P. Marra. 2011. West Nile virus impacts in American crow populations are associated with human land use and climate. *Ecological Research* 26:5, 909-916. [Crossref]
47. Jayne M. Deichmeister, Aparna Telang. 2011. Abundance of West Nile virus mosquito vectors in relation to climate and landscape variables. *Journal of Vector Ecology* 36:1, 75-85. [Crossref]
48. Sarah E. Bowden, Krisztian Magori, John M. Drake. 2011. Regional Differences in the Association Between Land Cover and West Nile Virus Disease Incidence in Humans in the United States. *The American Journal of Tropical Medicine and Hygiene* 84:2, 234-238. [Crossref]

49. Debarchana Ghosh, Rajarshi Guha. 2011. Using a neural network for mining interpretable relationships of West Nile risk factors. *Social Science & Medicine* **72**:3, 418-429. [[Crossref](#)]
50. V. CHEVALIER, A. DUPRESSOIR, A. TRAN, O. M. DIOP, C. GOTTLAND, M. DIALLO, E. ETTER, M. NDIAYE, V. GROBOIS, M. DIA, N. GAIDET, A. A. SALL, V. SOTI, M. NIANG. 2010. Environmental risk factors of West Nile virus infection of horses in the Senegal River basin. *Epidemiology and Infection* **138**:11, 1601-1609. [[Crossref](#)]
51. A. Marm Kilpatrick, Gregory D. Ebel, Michael R. Reddy, Dina M. Fonseca, Laura D. Kramer. 2010. Spatial and Temporal Variation in Vector Competence of Culex pipiens and Cx. restuans Mosquitoes for West Nile Virus. *The American Journal of Tropical Medicine and Hygiene* **83**:3, 607-613. [[Crossref](#)]
52. Samantha E. J. Gibbs. 2010. Avian biology, the human influence on global avian influenza transmission, and performing surveillance in wild birds. *Animal Health Research Reviews* **11**:01, 35-41. [[Crossref](#)]
53. Debarchana Ghosh, Rajarshi Guha. 2010. Use of genetic algorithm and neural network approaches for risk factor selection: A case study of West Nile virus dynamics in an urban environment. *Computers, Environment and Urban Systems* **34**:3, 189-203. [[Crossref](#)]
54. Archie C.A. Clements, Dirk U. Pfeiffer. 2009. Emerging viral zoonoses: Frameworks for spatial and spatiotemporal risk assessment and resource planning. *The Veterinary Journal* **182**:1, 21-30. [[Crossref](#)]
55. V. Hongoh, L. Berrang-Ford, N. H. Ogden, R. Lindsay, M. E. Scott, H. Artsob. 2009. A review of environmental determinants and risk factors for avian-associated mosquito arboviruses in Canada. *Biodiversity* **10**:2-3, 83-91. [[Crossref](#)]
56. Karl L. Evans, Kevin J. Gaston, Stuart P. Sharp, Andrew McGowan, Michelle Simeoni, Ben J. Hatchwell. 2009. Effects of urbanisation on disease prevalence and age structure in blackbird Turdus merula populations. *Oikos* **118**:5, 774-782. [[Crossref](#)]
57. Michael C. Wimberly, Michael B. Hildreth, Stephen P. Boyte, Erik Lindquist, Lon Kightlinger. 2008. Ecological Niche of the 2003 West Nile Virus Epidemic in the Northern Great Plains of the United States. *PLoS ONE* **3**:12, e3744. [[Crossref](#)]
58. Shannon L. LaDeau, Peter P. Marra, A. Marm Kilpatrick, Catherine A. Calder. 2008. West Nile Virus Revisited: Consequences for North American Ecology. *BioScience* **58**:10, 937-946. [[Crossref](#)]
59. Catherine A. Bradley, Samantha E. J. Gibbs, Sonia Altizer. 2008. URBAN LAND USE PREDICTS WEST NILE VIRUS EXPOSURE IN SONGBIRDS. *Ecological Applications* **18**:5, 1083-1092. [[Crossref](#)]
60. John P. Swaddle, Stavros E. Calos. 2008. Increased Avian Diversity Is Associated with Lower Incidence of Human West Nile Infection: Observation of the Dilution Effect. *PLoS ONE* **3**:6, e2488. [[Crossref](#)]
61. Leilei Pan, Lixu Qin, Simon X. Yang, Jiangping Shuai. 2008. A Neural Network-Based Method for Risk Factor Analysis of West Nile Virus. *Risk Analysis* **28**:2, 487-496. [[Crossref](#)]
62. Shlomit Paz, Iris Albersheim. 2008. Influence of Warming Tendency on Culex pipiens Population Abundance and on the Probability of West Nile Fever Outbreaks (Israeli Case Study: 2001-2005). *EcoHealth* **5**:1, 40-48. [[Crossref](#)]
63. R. J. Eisen, L. Eisen. 2008. Spatial Modeling of Human Risk of Exposure to Vector-Borne Pathogens Based on Epidemiological Versus Arthropod Vector Data. *Journal of Medical Entomology* **45**:2, 181-192. [[Crossref](#)]
64. Shlomit Paz, Iris Albersheim. 2008. Influence of Warming Tendency on Culex pipiens Population Abundance and on the Probability of West Nile Fever Outbreaks (Israeli Case Study: 2001-2005). *EcoHealth* **5**:1, 40-48. [[Crossref](#)]
65. John P DeGroote, Ramanathan Sugumaran, Sarah M Brend, Brad J Tucker, Lyric C Bartholomay. 2008. Landscape, demographic, entomological, and climatic associations with human disease incidence of West Nile virus in the state of Iowa, USA. *International Journal of Health Geographics* **7**:1, 19. [[Crossref](#)]
66. John DeGroote, David R. Mercer, Jeffrey Fisher, Ramanathan Sugumaran. 2007. Spatiotemporal Investigation of Adult Mosquito (Diptera: Culicidae) Populations in an Eastern Iowa County, USA. *Journal of Medical Entomology* **44**:6, 1139-1150. [[Crossref](#)]
67. William J. Landesman, Brian F. Allan, R. Brian Langerhans, Tiffany M. Knight, Jonathan M. Chase. 2007. Inter-Annual Associations Between Precipitation and Human Incidence of West Nile Virus in the United States. *Vector-Borne and Zoonotic Diseases* **7**:3, 337-343. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
68. Vanessa O. Ezenwa, Lesley E. Milheim, Michelle F. Coffey, Marvin S. Godsey, Raymond J. King, Stephen C. Gupitill. 2007. Land Cover Variation and West Nile Virus Prevalence: Patterns, Processes, and Implications for Disease Control. *Vector-Borne and Zoonotic Diseases* **7**:2, 173-180. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
69. NIKOLAOS W. YIANNAKOULIAS, LAWRENCE W. SVENSON. 2007. West Nile Virus. *Annals of the New York Academy of Sciences* **1102**:1, 135-148. [[Crossref](#)]
70. Catherine A. Bradley, Sonia Altizer. 2007. Urbanization and the ecology of wildlife diseases. *Trends in Ecology & Evolution* **22**:2, 95-102. [[Crossref](#)]