

AndrewBowen_Data606_LAb1

Andrew Bowen

2022-08-26

```
## Loading required package: airports
## Loading required package: cherryblossom
## Loading required package: usdata
##
## Attaching package: 'dplyr'
## The following objects are masked from 'package:stats':
##
##   filter, lag
## The following objects are masked from 'package:base':
##
##   intersect, setdiff, setequal, union
data('arbuthnot', package='openintro')
head(arbuthnot, 10)
```

```
## # A tibble: 10 x 3
##   year  boys girls
##   <int> <int> <int>
## 1  1629  5218  4683
## 2  1630  4858  4457
## 3  1631  4422  4102
## 4  1632  4994  4590
## 5  1633  5158  4839
## 6  1634  5035  4820
## 7  1635  5106  4928
## 8  1636  4917  4605
## 9  1637  4703  4457
## 10 1638  5359  4952
```

Exercise 1


Just want to see the count of girls baptized

```
arbuthnot$girls

## [1] 4683 4457 4102 4590 4839 4820 4928 4605 4457 4952 4784 5332 5200 4910 4617
## [16] 3997 3919 3395 3536 3181 2746 2722 2840 2908 2959 3179 3349 3382 3289 3013
## [31] 2781 3247 4107 4803 4881 5681 4858 4319 5322 5560 5829 5719 6061 6120 5822
## [46] 5738 5717 5847 6203 6033 6041 6299 6533 6744 7158 7127 7246 7119 7214 7101
## [61] 7167 7302 7392 7316 7483 6647 6713 7229 7767 7626 7452 7061 7514 7656 7683
## [76] 5738 7779 7417 7687 7623 7380 7288
```

Plotting our DF

```
ggplot(data=arbuthnot, aes(x=year, y=boys)) + geom_point()
```




Same plot as above but in line format

Exercise 2

Plotting a line graph the # of girls baptized by year below. There is a general upward trend in baptisms. On a smaller timeframe, there was a dip in girl baptisms between 1640 and 1660.

```
ggplot(data=arbuthnot, aes(x=year, y=girls)) + geom_line()
```



```
arbuthnot$boys + arbuthnot$girls
```

```
## [1] 9901 9315 8524 9584 9997 9855 10034 9522 9160 10311 10150 10850
## [13] 10670 10370 9410 8104 7966 7163 7332 6544 5825 5612 6071 6128
## [25] 6155 6620 7004 7050 6685 6170 5990 6971 8855 10019 10292 11722
## [37] 9972 8997 10938 11633 12335 11997 12510 12563 11895 11851 11775 12399
## [49] 12626 12601 12288 12847 13355 13653 14735 14702 14730 14694 14951 14588
## [61] 14771 15211 15054 14918 15159 13632 13976 14861 15829 16052 15363 14639
## [73] 15616 15687 15448 11851 16145 15369 16066 15862 15220 14928
```

Adding total field to dataframe

```
arbuthnot <- arbuthnot %>% mutate(total = boys + girls)
```


```
head(arbuthnot, 10)
```

```
## # A tibble: 10 x 4
##   year boys girls total
##   <int> <int> <int> <int>
## 1 1629 5218 4683 9901
## 2 1630 4858 4457 9315
## 3 1631 4422 4102 8524
## 4 1632 4994 4590 9584
## 5 1633 5158 4839 9997
## 6 1634 5035 4820 9855
## 7 1635 5106 4928 10034
## 8 1636 4917 4605 9522
## 9 1637 4703 4457 9160
```

```
## 10 1638 5359 4952 10311
```

Plotting total over time

```
ggplot(data=arbuthnot, aes(x=year, y=total)) + geom_line()
```



Calculating boy-to-girl ratio

```
arbuthnot <- arbuthnot %>% mutate(boy_to_girl_ratio = boys / girls)
```

Exercise 3

Calculating the ratio of boys baptized to total.

```
arbuthnot <- arbuthnot %>% mutate(boy_ratio = boys / total)
```

Finding years where more boys were present than girls. Adding in a boolean flag column to represent

```
arbuthnot <- arbuthnot %>% mutate(more_boys = boys > girls)
```

Let's take a look at our DF with all our added columns!

```
head(arbuthnot, 10)
```

```
## # A tibble: 10 x 7
##   year boys girls total boy_to_girl_ratio boy_ratio more_boys
##   <int> <int> <int> <int>          <dbl>      <dbl> <lgl>
## 1 1629 5218 4683 9901          1.11      0.527 TRUE
## 2 1630 4858 4457 9315          1.09      0.522 TRUE
## 3 1631 4422 4102 8524          1.08      0.519 TRUE
## 4 1632 4994 4590 9584          1.09      0.521 TRUE
```

##	5	1633	5158	4839	9997		1.07	0.516	TRUE
##	6	1634	5035	4820	9855		1.04	0.511	TRUE
##	7	1635	5106	4928	10034		1.04	0.509	TRUE
##	8	1636	4917	4605	9522		1.07	0.516	TRUE
##	9	1637	4703	4457	9160		1.06	0.513	TRUE
##	10	1638	5359	4952	10311		1.08	0.520	TRUE

More Practice

```
data('present', package='openintro')
```

```
head(present, 10)
```

```
## # A tibble: 10 x 3
##   year    boys  girls
##   <dbl> <dbl> <dbl>
## 1  1940 1211684 1148715
## 2  1941 1289734 1223693
## 3  1942 1444365 1364631
## 4  1943 1508959 1427901
## 5  1944 1435301 1359499
## 6  1945 1404587 1330869
## 7  1946 1691220 1597452
## 8  1947 1899876 1800064
## 9  1948 1813852 1721216
## 10 1949 1826352 1733177
```

Exercise 4

Let's see our data range in the year column first (using `min` & `max`)

```
print(min(present$year))
```

```
## [1] 1940
```

```
print(max(present$year))
```

```
## [1] 2002
```

Finding out our data frame dimensions (nrows x ncols)

```
print(ncol(present))
```

```
## [1] 3
```

```
print(nrow(present))
```

```
## [1] 63
```

Getting out dataframe columns with the built-in `colnames` function

```
colnames(present)
```

```
## [1] "year" "boys" "girls"
```

Exercise 5

Going to use the median count for boys and girls from each data set (`present` vs `arbuthnot`) to compare magnitudes of counts in each data set

```
p_boys_med = median(present$boys)
p_girls_med = median(present$girls)

a_boys_med = median(arbuthnot$boys)
a_girls_med = median(arbuthnot$girls)

# Calculating ratio of present boy/girl counts
boys_ratio = p_boys_med / a_boys_med
girls_ratio = p_girls_med / a_girls_med
print(boys_ratio)
```

```
## [1] 316.955
```

```
print(girls_ratio)
```

```
## [1] 320.3356
```

It looks like the median present day birth counts are ~320 times higher than the birth counts listed in our `arbuthnot` dataset. We used median counts to summarize the data set, so it won't be exactly this ratio for the whole data set, but modern birth counts are significantly higher.

Exercise 6


Setting up our boy-girl ratio column

```
present <- present %>% mutate(total = boys + girls)
present <- present %>% mutate(boys_ratio = boys / total)
head(present, 10)
```

```
## # A tibble: 10 x 5
##   year    boys    girls    total boys_ratio
##   <dbl>  <dbl>  <dbl>  <dbl>    <dbl>
## 1  1940 1211684 1148715 2360399    0.513
## 2  1941 1289734 1223693 2513427    0.513
## 3  1942 1444365 1364631 2808996    0.514
## 4  1943 1508959 1427901 2936860    0.514
## 5  1944 1435301 1359499 2794800    0.514
## 6  1945 1404587 1330869 2735456    0.513
## 7  1946 1691220 1597452 3288672    0.514
## 8  1947 1899876 1800064 3699940    0.513
## 9  1948 1813852 1721216 3535068    0.513
## 10 1949 1826352 1733177 3559529    0.513
```

Let's plot the ratio of boys born over time in our `present` dataset:

```
ggplot(data=present, aes(x=year, y=boys_ratio)) + geom_line()
```



While the ratio of boys born has stayed over 50%, it is experiencing a downward trend over time since 1940, the beginning of our dataset. The observation of boys being born more than girls from the `arbuthnot` dataset does hold up, but has decreased since 1940 in the US.

Exercise 7

```
# Truncating output to 10 rows for readability
head(present %>% arrange(desc(total)), 10)
```

```
## # A tibble: 10 x 5
##   year    boys  girls  total boys_ratio
##   <dbl> <dbl> <dbl> <dbl>   <dbl>
## 1 1961 2186274 2082052 4268326    0.512
## 2 1960 2179708 2078142 4257850    0.512
## 3 1957 2179960 2074824 4254784    0.512
## 4 1959 2173638 2071158 4244796    0.512
## 5 1958 2152546 2051266 4203812    0.512
## 6 1962 2132466 2034896 4167362    0.512
## 7 1956 2133588 2029502 4163090    0.513
## 8 1990 2129495 2028717 4158212    0.512
## 9 1991 2101518 2009389 4110907    0.511
## 10 1963 2101632 1996388 4098020    0.513
```

We see the highest number of total births in the US come in 1961 with 4268326 total births (boys & girls). It's interesting to note that 8 of the top 10 years in terms of total births came during the baby boom years in the post-war era.