

Counterexamples to two conjectures on domination and minimum maximal matchings in regular graphs

Andrew Butz*

September 2026 (version 2)

Abstract

Baste, Fürst, Henning, Mohr and Rautenbach [1] conjectured that every regular graph G of positive degree satisfies $\gamma(G) \leq \gamma_e(G)$, where γ is the domination number and γ_e the edge domination number, the minimum size of a maximal matching; the conjecture is known for degree at least seven [4] and for every cubic graph on at most 48 vertices [4]. The stronger statement $i(G) \leq \gamma_e(G)$, with i the independent domination number, was proposed by the TxGraffiti system in 2020 and is Conjecture 3 of [6]. We exhibit six pairwise non-isomorphic connected cubic graphs on 50 vertices with $i(G) = 16 > 15 = \gamma_e(G)$, three of which also have $\gamma(G) = 16$. Thus both conjectures fail for cubic graphs, and by the 48-vertex bound of [4] the three graphs are counterexamples of minimum order to the first. Lexicographic products with edgeless graphs turn any of the six into a connected $3k$ -regular counterexample to the TxGraffiti conjecture for every $k \geq 1$. The graphs are the incidence graphs of minimally unsatisfiable 3-CNF formulas in which every literal occurs exactly twice, a construction that appears in [4]; the formulas are obtained by a gluing operation from a complete census of the eight such formulas of deficiency three on seven variables. A census of unsatisfiable formulas of this type on fifteen variables (Section 5) yields at least 239 pairwise non-isomorphic cubic graphs of order 50 with $\mu^* = 15$, of which 67 have $i = 16$ and 26 have $\gamma = 16$; 203 of the 239 are not gluings.

1 Statement

For a graph G , $\gamma(G)$ is the domination number, $i(G)$ the independent domination number (the minimum size of a maximal independent set), and $\gamma_e(G) = \mu^*(G)$ the edge domination number, the minimum size of a maximal matching. Since a maximal independent set is dominating, $\gamma(G) \leq i(G)$. Two conjectures compare these to γ_e .

- (A) [1]: if G is Δ -regular with $\Delta \geq 1$, then $\gamma(G) \leq \gamma_e(G)$. Known for $\Delta \leq 2$ trivially, for $\Delta \geq 7$ by the Lovász Local Lemma [4], for claw-free and fork-free graphs, and for every cubic graph on at most 48 vertices [4, Prop. 7]; the cubic case is described as open in [4].
- (B) [6, Conj. 3]: if G is r -regular with $r > 0$, then $i(G) \leq \mu^*(G)$, and this bound is sharp. Generated by TxGraffiti in 2020, reported open for every $r \geq 3$; (B) implies (A).

Theorem 1. *There are six pairwise non-isomorphic connected cubic graphs on 50 vertices with $i(G) = 16$ and $\mu^*(G) = 15$; three of them have $\gamma(G) = 16$. Hence (A) and (B) both fail for cubic graphs, and the three graphs are counterexamples to (A) of minimum order. Moreover, for every $r \equiv 0 \pmod{3}$ there is a connected r -regular graph with $i(G) > \mu^*(G)$.*

*andrew@cosmicbutz.com. The author gratefully acknowledges extensive AI assistance (Anthropic's Claude) in computation, proof development and verification.

The inequalities are attained with equality on many graphs ($K_{3,3}$, the Petersen graph, the prism over a triangle for (A), and the graph G_{50} of Remark 10 for both), so only the inequalities fail, not the sharpness clauses. Conjecture (B) had been verified exhaustively for cubic graphs on at most 20 vertices at minimum maximal matchings; the smallest cubic counterexample to (B) therefore has between 22 and 50 vertices. For (A) the minimum is exactly 50.

2 From formulas to graphs

A CNF formula is *minimally unsatisfiable* (MU) if it is unsatisfiable and every proper subset is satisfiable [8]. Let F be a 3-uniform CNF formula on variables $x_1, \dots, x_v$ with m clauses in which every literal occurs exactly twice (so $3m = 4v$). Define $G(F)$ as follows: for each variable x take two adjacent vertices x^+, x^- (a *pivot pair*); for each clause C take a vertex u_C adjacent to the endpoint named by each of its three literals ($x \mapsto x^+, \bar{x} \mapsto x^-$).

Lemma 2. *$G(F)$ is a simple 3-regular graph on $2v + m$ vertices; the v pivot edges form a maximal matching M ; and F is unsatisfiable if and only if no independent set of pivot endpoints dominates all clause vertices. If F is MU then $G(F)$ is connected.*

Proof. Each endpoint has one pivot edge and two clause edges; each clause vertex has three edges to endpoints of three distinct pairs. The clause vertices are pairwise non-adjacent, so M is maximal. An independent set of endpoints contains at most one endpoint per pair, i.e. it is a partial truth assignment, and it dominates u_C iff it makes a literal of C true. Connectedness follows since an MU formula has a connected clause–variable incidence graph. $\square$

Thus for unsatisfiable F , no independent dominating set of $G(F)$ can be built by extending the clause vertices, and $|M| = v$ competes with $i(G(F))$ with no further structure to help; $\mu^*(G(F)) \leq v$ always.

3 Gluing minimally unsatisfiable formulas

Lemma 3 (Gluing). *Let A, B be MU formulas on disjoint variable sets, $a_1 \neq a_2 \in A$, $b_1 \neq b_2 \in B$, and x a fresh variable. Then $K := (A \text{ with } x \text{ added to } a_1, a_2) \cup (B \text{ with } \bar{x} \text{ added to } b_1, b_2)$ is MU. Literal occurrences are unchanged except that x and $\bar{x}$ occur twice each, and $|K| - |\text{var } K| = (|A| - |\text{var } A|) + (|B| - |\text{var } B|) - 1$.*

Proof. $K[x=1] \supseteq B$ and $K[x=0] \supseteq A$, so K is unsatisfiable. Remove a clause of A 's side and set $x = 0$: the $\bar{x}$ -clauses are satisfied, A minus that clause is satisfiable, and $B \setminus \{b_1, b_2\}$ is satisfiable as a proper subset of an MU formula; the variable sets are disjoint. The other side is symmetric. $\square$

Let R_3 be the following formula on 7 variables (two 2-clauses and eight 3-clauses, every literal exactly twice, MU):

$$R_3 = (x_1 \vee x_2) (x_3 \vee x_4) (x_1 \vee \bar{x}_2 \vee x_5) (\bar{x}_1 \vee x_2 \vee x_6) (\bar{x}_1 \vee x_5 \vee x_7) \\ (\bar{x}_2 \vee x_6 \vee \bar{x}_7) (x_3 \vee \bar{x}_4 \vee \bar{x}_5) (\bar{x}_3 \vee x_4 \vee \bar{x}_6) (\bar{x}_3 \vee \bar{x}_5 \vee x_7) (\bar{x}_4 \vee \bar{x}_6 \vee \bar{x}_7)$$

It is one of exactly eight isomorphism classes of MU formulas with these parameters, found by a complete canonical enumeration (Appendix B). Gluing two copies of R_3 through a fresh variable

x_{15} placed in their four 2-clauses gives

$$\begin{aligned}
K_1 = & (x_1 \vee x_2 \vee x_{15}) \quad (x_3 \vee x_4 \vee x_{15}) \quad (x_1 \vee \bar{x}_2 \vee x_5) \quad (\bar{x}_1 \vee x_2 \vee x_6) \\
& (\bar{x}_1 \vee x_5 \vee x_7) \quad (\bar{x}_2 \vee x_6 \vee \bar{x}_7) \quad (x_3 \vee \bar{x}_4 \vee \bar{x}_5) \quad (\bar{x}_3 \vee x_4 \vee \bar{x}_6) \\
& (\bar{x}_3 \vee \bar{x}_5 \vee x_7) \quad (\bar{x}_4 \vee \bar{x}_6 \vee \bar{x}_7) \quad (x_8 \vee x_9 \vee \bar{x}_{15}) \quad (x_{10} \vee x_{11} \vee \bar{x}_{15}) \\
& (x_8 \vee \bar{x}_9 \vee x_{12}) \quad (\bar{x}_8 \vee x_9 \vee x_{13}) \quad (\bar{x}_8 \vee x_{12} \vee x_{14}) \quad (\bar{x}_9 \vee x_{13} \vee \bar{x}_{14}) \\
& (x_{10} \vee \bar{x}_{11} \vee \bar{x}_{12}) \quad (\bar{x}_{10} \vee x_{11} \vee \bar{x}_{13}) \quad (\bar{x}_{10} \vee \bar{x}_{12} \vee x_{14}) \quad (\bar{x}_{11} \vee \bar{x}_{13} \vee \bar{x}_{14})
\end{aligned}$$

a 3-uniform MU formula on 15 variables with 20 clauses in which every literal occurs exactly twice.

4 The graph G_1 and its verification

Let $G_1 = G(K_1)$: 50 vertices, 75 edges, connected, cubic. Its adjacency list is in Appendix A (vertices $2(j-1), 2(j-1)+1$ are $x_j^\pm$; vertices $30+t$ are the clause vertices in the order above).

Proposition 4. $i(G_1) = 16$, $\gamma(G_1) = 14$ and $\mu^*(G_1) = 15$.

Verification. The 15 pivot edges form a maximal matching (Lemma 2), so $\mu^*(G_1) \leq 15$. Conversely, in a cubic graph every edge of a maximal matching dominates at most five edges (itself and two at each endpoint), and a maximal matching dominates all edges, so any maximal matching of a cubic graph with 75 edges has at least $\lceil 75/5 \rceil = 15$ edges; hence $\mu^*(G_1) = 15$ (also confirmed by integer programming). The set $\{1, 3, 4, 7, 9, 11, 12, 15, 17, 19, 20, 23, 24, 27, 30, 40\}$ is independent and dominating, so $i(G_1) \leq 16$. That no independent dominating set of size 15 exists was established by four independent methods: two SAT solvers (CaDiCaL 1.5.3 with a sequential-counter cardinality encoding; Glucose 4 with a totalizer encoding), the HiGHS integer programming solver, and a solver-free exhaustive branch-and-bound over independent dominating sets (186,188 search nodes). The domination number 14 was computed by integer programming and SAT. $\square$

The same construction applied to the other pairs of blocks gives, among the 36 gluings of the eight blocks, exactly six graphs with $i = 16$ (all with $\mu^* = 15$), 25 with $i = \mu^* = 15$ and five with $i = 14$. The six are pairwise non-isomorphic; their domination numbers are listed as well. For the three graphs with $\gamma = 16$, the non-existence of a dominating set of size 15 was verified by the HiGHS integer programming solver, by Glucose 4 with a totalizer encoding, and by a solver-free exhaustive branch-and-bound over dominating sets (27,699,453, 24,899,125 and 25,088,473 search nodes respectively); dominating sets of size 16 were checked by definition. In all six graphs (and in G_{50}) the pivot matching is the unique maximal matching of size 15, so it is the unique minimum maximal matching, and it admits no dominating transversal in the sense of [4]. Every clause vertex is the centre of an induced claw and the graphs contain induced forks, so the claw-free and fork-free cases of (A), which are theorems, are not contradicted; the multiplicative bound $\gamma \leq (7/6 - 1/204)\gamma_e$ of [1] is respected, $16/15 < 1.16$.

graph	i	γ	μ^*	graph6
$R_3 + R_3$	16	14	15	q' ?G?C??G??@????_???@?????G?????C?????G?????@???????_???????@?????????
$R_3 + R_6$	16	15	15	q' ?G?C??G??@????_???@?????G?????C?????G?????@???????_???????@?????????
$R_3 + R_7$	16	15	15	q' ?G?C??G??@????_???@?????G?????C?????G?????@???????_???????@?????????
$R_6 + R_6$	16	16	15	q' ?G?C??G??@????_???@?????G?????C?????G?????@???????_???????@?????????
$R_6 + R_7$	16	16	15	q' ?G?C??G??@????_???@?????G?????C?????G?????@???????_???????@?????????
$R_7 + R_7$	16	16	15	q' ?G?C??G??@????_???@?????G?????C?????G?????@???????_???????@?????????

5 A census of tight formulas on fifteen variables

Call a 3-uniform CNF formula on 15 variables with 20 clauses in which every literal occurs exactly twice *tight*. Every unsatisfiable tight formula is MU: deleting a clause leaves a 3-uniform formula with 19 clauses in which every literal occurs at most twice, and such a formula is satisfiable because $\mu(3, 2, 2) = 20$ [5]. By Lemma 2 and the matching count in Section 4, every unsatisfiable tight formula F therefore gives a connected cubic graph $G(F)$ of order 50 with $\mu^*(G(F)) = 15$, and the only question is whether i or γ exceeds 15. The formulas $K_1 = R_3 \oplus R_3$ and its five companions are tight, but tightness does not require a cut variable, and most tight formulas have none.

Theorem 5. *There are at least 239 pairwise non-isomorphic connected cubic graphs of order 50 of the form $G(F)$ with F an unsatisfiable tight formula. Each has $\mu^* = 15$ and girth 5. Among them 67 satisfy $i = 16 > \mu^*$ and 26 satisfy $\gamma = 16 > \gamma_e$; the joint distribution is*

(γ, i)	(14, 14)	(14, 15)	(14, 16)	(15, 15)	(15, 16)	(16, 16)	total
no cut variable	34	50	0	58	38	23	203
cut variable (gluing)	5	15	1	10	2	3	36

Exactly 36 of the 239 formulas have a cut variable, i.e. a variable whose two literal vertices separate $G(F)$; these are precisely the 36 gluings $R_a \oplus R_b$ ($0 \leq a \leq b \leq 7$) of Section 3, and they are pairwise non-isomorphic. The remaining 203 formulas are not gluings of smaller MU formulas; 61 of them have $i = 16$ and 23 have $\gamma = 16$.

Verification. Formulas were found by simulated annealing over tight formulas: the state is an assignment of the 60 literal occurrences to 20 clauses, a move exchanges two occurrences between different clauses (which preserves the exact-twice property; moves creating a repeated variable in a clause are rejected), and the energy is the number of satisfying assignments, evaluated bit-parallel over all 2^{15} assignments. Energy zero is unsatisfiability, hence minimal unsatisfiability by the remark above. Every formula found was then expanded: all tight formulas reachable from it by one or two moves were tested, and new unsatisfiable ones were added and expanded in turn; the process closed with 239 isomorphism classes. Isomorphism of the graphs $G(F)$ was decided by canonical labelling with nauty [9]. For each class, γ and i were computed by the HiGHS integer programming solver, the optimal sets were checked by definition, the non-existence of a dominating set, resp. independent dominating set, of size one less than the computed value was confirmed independently for every class by CaDiCaL 1.5.3 with a sequential-counter cardinality encoding, and $\mu^* = 15$ was confirmed by integer programming over maximal matchings. Cut variables were detected by deleting the two literal vertices of each variable and testing connectivity. The 36 gluings were generated directly from the blocks $R_0, \dots, R_7$ and matched to the census by canonical form. All 239 formulas, graphs (graph6), optimal sets and certificates are in `census_tight_15.20.json` in the repository. $\square$

Remark 6. No single move takes an unsatisfiable tight formula to a non-isomorphic unsatisfiable one: in 110 of the 239 classes some move produces an isomorphic copy, in the other 129 every move produces a satisfiable formula; the list is closed under sequences of two moves. Independent annealing runs from fresh seeds, made after the closure, produced 92 unsatisfiable formulas in 47 isomorphism classes, all among the 239. For nine of the formulas—six without a cut variable, covering the five occupied cells of the first row of the table, and the gluings $R_6 \oplus R_7$, $R_3 \oplus R_3$ and $R_0 \oplus R_0$ —we also enumerated every formula reachable by three moves through valid intermediates: about 10^6 two-move endpoints each and, in total, 134,571 unsatisfiable three-move endpoints, all

of them among the 239 (a formula without a cut variable reaches 21–27 of the classes within three moves, a gluing 5–13). We do not know whether the list is complete. The automorphism groups of $G(F)$ are small: 96 of the 239 graphs are asymmetric, 97 have a group of order 2, 22 of order 4 and 14 of order 8; orders 16, 32 and 128 occur only for gluings, the last for $G_{50} = G(R_0 \oplus R_0)$. The census also shows that the frequency of $\gamma = 16$ among the gluings (3 of 36) is not representative: among the 203 formulas without a cut variable it is 23, and $i = 16$ occurs in 61.

Lemma 7. *Let F be an unsatisfiable 3-CNF formula with at most 20 clauses in which every literal occurs at most twice. Then*

- (a) *no two clauses of F contain the same two literals;*
- (b) *if a, b are literals of distinct variables such that some clause contains $\{a, b\}$ and some clause contains $\{\bar{a}, \bar{b}\}$, then some clause contains $\{a, \bar{b}\}$ or $\{\bar{a}, b\}$.*

In particular $G(F)$ has girth at least 5 for every unsatisfiable tight formula F .

Proof. Both parts use the substitution $b \mapsto \bar{a}$, $\bar{b} \mapsto a$, followed by the deletion of the clauses that have become tautologies; call the result F^* . A model of F^* extends to a model of F by setting $b = \bar{a}$, so F^* is unsatisfiable. We show that in each case F^* is a 3-CNF formula with at most 18 clauses in which every literal occurs at most twice, contradicting $\mu(3, 2, 2) > 18$ [5, Thm. 15].

(a) Let $C \neq C'$ both contain $\{a, b\}$. Since a and b occur at most twice, C and C' are the only clauses containing a and the only clauses containing b ; both become tautologies. A clause E containing $\bar{b}$ does not contain a , so it becomes $E - \bar{b} + a$, a clause on three distinct variables, unless it contains $\bar{a}$, in which case it is a tautology and is deleted. In F^* the literal a occurs at most twice (once for each former occurrence of $\bar{b}$), $\bar{a}$ at most twice, and every other literal at most as often as in F ; and F^* has at most 18 clauses.

(b) Let $K_1 \supseteq \{a, b\}$ and $K_2 \supseteq \{\bar{a}, \bar{b}\}$, and suppose that no clause contains $\{a, \bar{b}\}$ or $\{\bar{a}, b\}$. Then K_1 and K_2 become tautologies. The other clause containing a , if any, contains neither b (by (a)) nor $\bar{b}$ (by assumption), so it is unchanged; the other clause containing $\bar{b}$, if any, contains neither a (by assumption) nor $\bar{a}$ (by (a)), so it becomes a clause on three distinct variables with $\bar{b}$ replaced by a . Likewise the other clause containing $\bar{a}$, if any, is unchanged, and in the other clause containing b , if any, the literal b is replaced by $\bar{a}$. Hence F^* has at most 18 clauses, a and $\bar{a}$ occur at most twice in it, and no other literal occurs more often than in F .

For the last claim, $G(F)$ has no triangle: clause vertices are pairwise non-adjacent, no clause contains a variable twice, and x^+ , x^- have no common neighbour. A 4-cycle therefore uses no pivot edge and has the form $u_C \ell u_{C'} \ell' u_C$, so the clauses $C \neq C'$ both contain ℓ and ℓ' , which (a) excludes. $\square$

Remark 8. The *clause graph* of a tight formula F has a vertex for each clause and an edge for each literal, joining the two clauses that contain it; it is a cubic graph on 20 vertices, simple by Lemma 7(a), and F is recovered from it by specifying which pairs of edges are $\{x, \bar{x}\}$. The 239 formulas of Theorem 5 have only 15 distinct clause graphs: 9 connected ones, and the 6 disconnected ones formed by two of the three 10-vertex cubic graphs that carry the blocks $R_0, \dots, R_7$. Every one of the 15 contains at least four pairs of *twin* clauses, two clauses whose three literals continue into the same three other clauses (an induced $K_{2,3}$): 214 of the 239 formulas have exactly four such pairs, 18 have five and 7 have six, whereas among the 510,489 connected cubic graphs on 20 vertices only 217 have four or more, and among the 5,855 disconnected ones only 563. Each of the 15 clause graphs has at least 128 automorphisms, and none is 3-edge-connected. Whether every unsatisfiable tight formula has at least four twin pairs is open; a proof would reduce the exact census to these 780 cubic graphs and their edge pairings.

6 Higher regularity

For a graph G and $k \geq 1$ let $G[\overline{K}_k]$ be the lexicographic product with the edgeless graph on k vertices: each vertex v is replaced by k pairwise non-adjacent copies, and copies of adjacent vertices are completely joined. If G is connected and r -regular, $G[\overline{K}_k]$ is connected and rk -regular.

Lemma 9. $i(G[\overline{K}_k]) = k i(G)$ and $\mu^*(G[\overline{K}_k]) \leq k \mu^*(G)$.

Proof. Let I' be an independent dominating set of $G[\overline{K}_k]$ and I the set of vertices of G having a copy in I' . If a copy (v, j) lies in I' and another copy (v, j') does not, then (v, j') is dominated by some $(u, i) \in I'$ with $u \sim v$, but then $(u, i) \sim (v, j)$, contradicting independence; so $I' = I \times [k]$. Independence and domination of I' transfer to I , hence $|I'| \geq k i(G)$, and $I \times [k]$ for an optimal I shows equality. For the matching, take a maximal matching M of G and the k edges $(u, j)(v, j)$, $j \in [k]$, for each $uv \in M$; the uncovered vertices are the copies of the M -uncovered vertices, an independent set, so this matching of size $k|M|$ is maximal. $\square$

Hence $i(G_1[\overline{K}_k]) = 16k > 15k \geq \mu^*(G_1[\overline{K}_k])$ for every k , which proves the last claim of Theorem 1; for $k = 2$ we verified $i = 32$, $\mu^* = 30$ directly by integer programming. No such extension exists for γ : a dominating set of $G[\overline{K}_k]$ may use a single copy of each vertex of a total dominating set of G , so $\gamma(G[\overline{K}_k])$ stays bounded while γ_e grows, in accordance with the theorem of [4] that (A) holds for all $\Delta \geq 7$. Regularities $r \not\equiv 0 \pmod{3}$ remain open: the natural analogue for $r = 4$ is a 4-uniform MU formula in which every literal occurs exactly three times (then $4m = 6v$, i.e. $2m = 3v$), whose realization is a 4-regular graph with an obstructed maximal matching, and the smallest conceivable such formula has 12 variables and 18 clauses.

7 Remarks

Remark 10. Unsatisfiable 3-CNF formulas in which every literal occurs exactly twice are known to exist [2, 3], and the least number of clauses of such a formula is 20 [5]; it is this fact that yields the 48-vertex bound of [4, Prop. 7], and it shows that $n = 50$ is the least order at which the present construction can produce anything, since $3m = 4v$ and $m \geq 20$ force $v \geq 15$. Gupta [4, Thm. 2] builds the incidence graph of one such minimum formula and shows that its unique minimum maximal matching admits no dominating transversal while $\gamma = 14 < 15 = \gamma_e$. That graph is isomorphic to our $G_{50} = G(R_0 \oplus R_0)$, the gluing of two copies of R_0 (itself the gluing of two copies of F_3), for which also $i = 15 = \mu^*$; the counterexamples of Theorem 1 arise from the other blocks of the census. Gluing is, however, far from necessary: of the 239 unsatisfiable tight formulas of Theorem 5, only the 36 gluings have a cut variable.

Remark 11. In the language of minimally unsatisfiable formulas: every MU formula of deficiency one with literal degrees at most two has at least four more literal occurrences than clauses, every such formula of deficiency two has at least three, and no 3-uniform MU formula with literal degrees at most two has deficiency ≤ 3 (the last is a short descent argument using the pocket structure of restriction cores). The blocks $R_0, \dots, R_7$ are the complete list of MU formulas of deficiency three and defect two on seven variables; gluing produces 3-uniform members at every deficiency ≥ 5 , and by [5] none exists at deficiency four with fewer than 20 clauses.

Remark 12. The route to these graphs went through a structural theory of obstructed maximal matchings in cubic graphs (demand systems, forcing, the deficiency laws above); the counterexamples live in the one anatomy—all pivot pairs, all loads two, no cycles—that the all-load-one theory and Haxell's independent transversal theorem [7] exclude, which is why they were invisible

to searches guided by that theory. Regularities $r \not\equiv 0 \pmod{3}$ remain open for (B), and degrees $4 \leq \Delta \leq 6$ for (A); the natural analogue for $r = 4$ is a 4-uniform MU formula in which every literal occurs exactly three times.

A Adjacency list of G_1

0: 1, 30, 32
1: 0, 33, 34
2: 3, 30, 33
3: 2, 32, 35
4: 5, 31, 36
5: 4, 37, 38
6: 7, 31, 37
7: 6, 36, 39
8: 9, 32, 34
9: 8, 36, 38
10: 11, 33, 35
11: 10, 37, 39
12: 13, 34, 38
13: 12, 35, 39
14: 15, 40, 42
15: 14, 43, 44
16: 17, 40, 43
17: 16, 42, 45
18: 19, 41, 46
19: 18, 47, 48
20: 21, 41, 47
21: 20, 46, 49
22: 23, 42, 44
23: 22, 46, 48
24: 25, 43, 45
25: 24, 47, 49
26: 27, 44, 48
27: 26, 45, 49
28: 29, 30, 31
29: 28, 40, 41
30: 0, 2, 28
31: 4, 6, 28
32: 0, 3, 8
33: 1, 2, 10
34: 1, 8, 12
35: 3, 10, 13
36: 4, 7, 9
37: 5, 6, 11
38: 5, 9, 12
39: 7, 11, 13
40: 14, 16, 29

41: 18, 20, 29
 42: 14, 17, 22
 43: 15, 16, 24
 44: 15, 22, 26
 45: 17, 24, 27
 46: 18, 21, 23
 47: 19, 20, 25
 48: 19, 23, 26
 49: 21, 25, 27

B The eight blocks

Isomorphism classes of MU formulas on seven variables with two 2-clauses, eight 3-clauses and every literal occurring exactly twice (R_0 is the gluing of two copies of F_3 ; R_4 was found independently by a second engine):

$$R_0 \quad (x_1 \vee x_2) (x_3 \vee x_4) (x_1 \vee \bar{x}_2 \vee x_5) (\bar{x}_1 \vee x_2 \vee \bar{x}_5) (\bar{x}_1 \vee x_5 \vee x_6) (\bar{x}_2 \vee \bar{x}_5 \vee x_6) (x_3 \vee \bar{x}_4 \vee x_7) (\bar{x}_3 \vee x_4 \vee \bar{x}_7) (\bar{x}_3 \vee \bar{x}_6 \vee x_7) (\bar{x}_4 \vee \bar{x}_6 \vee \bar{x}_7)$$

$$R_1 \quad (x_1 \vee x_2) (x_3 \vee x_4) (x_1 \vee \bar{x}_2 \vee x_5) (\bar{x}_1 \vee x_2 \vee x_6) (\bar{x}_1 \vee \bar{x}_3 \vee x_5) (\bar{x}_2 \vee \bar{x}_4 \vee x_6) (\bar{x}_3 \vee \bar{x}_5 \vee x_7) (x_3 \vee \bar{x}_6 \vee x_7) (x_4 \vee \bar{x}_5 \vee \bar{x}_7) (\bar{x}_4 \vee \bar{x}_6 \vee \bar{x}_7)$$

$$R_2 \quad (x_1 \vee x_2) (x_3 \vee x_4) (x_1 \vee \bar{x}_2 \vee x_5) (\bar{x}_1 \vee x_2 \vee x_6) (\bar{x}_1 \vee \bar{x}_3 \vee x_5) (\bar{x}_2 \vee x_6 \vee x_7) (x_3 \vee \bar{x}_4 \vee \bar{x}_7) (\bar{x}_3 \vee \bar{x}_5 \vee \bar{x}_7) (x_4 \vee \bar{x}_5 \vee \bar{x}_6) (\bar{x}_4 \vee \bar{x}_6 \vee x_7)$$

$$R_3 \quad (x_1 \vee x_2) (x_3 \vee x_4) (x_1 \vee \bar{x}_2 \vee x_5) (\bar{x}_1 \vee x_2 \vee x_6) (\bar{x}_1 \vee x_5 \vee x_7) (\bar{x}_2 \vee x_6 \vee \bar{x}_7) (x_3 \vee \bar{x}_4 \vee \bar{x}_5) (\bar{x}_3 \vee x_4 \vee \bar{x}_6) (\bar{x}_3 \vee \bar{x}_5 \vee x_7) (\bar{x}_4 \vee \bar{x}_6 \vee \bar{x}_7)$$

$$R_4 \quad (x_1 \vee x_2) (\bar{x}_1 \vee x_3) (x_1 \vee \bar{x}_2 \vee x_3) (\bar{x}_1 \vee x_2 \vee x_4) (\bar{x}_2 \vee \bar{x}_3 \vee x_4) (\bar{x}_3 \vee x_5 \vee x_6) (\bar{x}_4 \vee \bar{x}_5 \vee x_7) (\bar{x}_4 \vee \bar{x}_6 \vee \bar{x}_7) (x_5 \vee \bar{x}_6 \vee x_7) (\bar{x}_5 \vee x_6 \vee \bar{x}_7)$$

$$R_5 \quad (x_1 \vee x_2) (\bar{x}_1 \vee x_3) (x_1 \vee \bar{x}_2 \vee x_3) (\bar{x}_1 \vee x_2 \vee x_4) (\bar{x}_2 \vee x_4 \vee x_5) (\bar{x}_3 \vee \bar{x}_5 \vee x_6) (\bar{x}_3 \vee \bar{x}_6 \vee x_7) (\bar{x}_4 \vee x_5 \vee \bar{x}_7) (\bar{x}_4 \vee x_6 \vee x_7) (\bar{x}_5 \vee \bar{x}_6 \vee \bar{x}_7)$$

$$R_6 \quad (x_1 \vee x_2) (\bar{x}_1 \vee x_3) (\bar{x}_1 \vee x_2 \vee x_4) (x_1 \vee x_3 \vee x_5) (\bar{x}_2 \vee x_4 \vee x_6) (\bar{x}_2 \vee \bar{x}_5 \vee \bar{x}_6) (\bar{x}_3 \vee x_5 \vee x_7) (\bar{x}_3 \vee \bar{x}_6 \vee \bar{x}_7) (\bar{x}_4 \vee \bar{x}_5 \vee x_7) (\bar{x}_4 \vee x_6 \vee \bar{x}_7)$$

$$R_7 \quad (x_1 \vee x_2) (\bar{x}_1 \vee x_3) (\bar{x}_1 \vee x_2 \vee x_4) (x_1 \vee x_3 \vee x_5) (\bar{x}_2 \vee x_4 \vee x_6) (\bar{x}_2 \vee \bar{x}_5 \vee x_7) (\bar{x}_3 \vee \bar{x}_4 \vee x_7) (\bar{x}_3 \vee x_5 \vee \bar{x}_6) (\bar{x}_4 \vee x_6 \vee \bar{x}_7) (\bar{x}_5 \vee \bar{x}_6 \vee \bar{x}_7)$$

References

- [1] J. Baste, M. Fürst, M. A. Henning, E. Mohr, D. Rautenbach, Domination versus edge domination, *Discrete Appl. Math.* 285 (2020) 343–349; arXiv:1906.10420.
- [2] P. Berman, M. Karpinski, A. D. Scott, Approximation hardness and satisfiability of bounded occurrence instances of SAT, ECCC report TR03-022, 2003.
- [3] P. Berman, M. Karpinski, A. D. Scott, Approximation hardness of short symmetric instances of MAX-3SAT, ECCC report TR03-049, 2003.

- [4] C. Gupta, Domination versus edge domination in regular graphs of degree at least seven, arXiv:2608.22498, 2026.
- [5] T. Zhang, T. Peitl, S. Szeider, Small unsatisfiable k -CNFs with bounded literal occurrence, in: *27th International Conference on Theory and Applications of Satisfiability Testing (SAT 2024)*, LIPIcs vol. 305, Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 2024, pp. 31:1–31:22.
- [6] R. Davila, B. Brimkov, R. Pepper, *In Reverie Together: Ten Years of Mathematical Discovery with a Machine Collaborator*, arXiv:2507.17780 [cs.DM], 2025.
- [7] P. E. Haxell, A note on vertex list colouring, *Combin. Probab. Comput.* 10 (2001) 345–347.
- [8] H. Kleine Büning, O. Kullmann, Minimal unsatisfiability and autarkies, in: *Handbook of Satisfiability*, IOS Press, 2009, ch. 11.
- [9] B. D. McKay, A. Piperno, Practical graph isomorphism, II, *J. Symbolic Comput.* 60 (2014) 94–112.