Follow-up Runs of the Water-Quality Benchmark on the Colorado River Network

Andrew Chincea
TDA Lab

November 2026

Abstract

This report continues `kron_vs_full_laplacian_report.pdf` (the "main report") and collects the benchmark runs made after it. Every run is a paired comparison against the leave-one-station-out (LOO) benchmark of the main report (its Table 1 values are repeated here), on identical test rows. So far it covers three runs. (1) Water-year indicators on the current rows change no target reliably except turbidity, where every linear model gains about 0.06 and the graph model 0.08; pH gets worse for Elastic Net and the Spline GAM. (2) With the longer 2021–2026 history, the water-year indicators remove the 1 °C cold offset found in the main report, but the history still does not beat the current rows for any target: the records cover different years at different stations, and for the linear models the indicators then also absorb which stations are present, so specific conductance and pH get much worse. (3) Gradient boosting chooses the neighbour reading for pH and turbidity only, and gains nothing reliable.

1 Setup

The data, graph, models, nested LOO and R^2 (equal weight per station) are those of the main report. Each run below changes one option of `water_quality_model_benchmark.py` and is compared with the same model of the main benchmark on identical test rows with `compare_runs.py`: ΔR^2 is the change, the 95% interval resamples stations with the predictions held fixed, and better/worse counts the stations whose mean squared error fell/rose. Intervals that exclude zero are in bold. With 17–24 stations, differences below about 0.03–0.05 are within noise.

Table 1: Reference: LOO R^2 of the main benchmark (main report, Table 14), which every run below is paired against.

Model	Water temp.	DO	Spec. cond.	pH	Turbidity
Graph-Laplacian	0.863	0.700	0.385	0.117	0.104
OLS	0.863	0.695	−0.099	−0.014	0.136
Ridge	0.864	0.701	−0.110	0.003	0.136
Elastic Net	0.863	0.705	−0.109	0.023	0.136
Spline GAM	0.838	0.708	0.022	0.012	0.091
Gradient Boosting	0.833	0.733	−0.094	−0.171	0.391

2 Water-year effect

Motivation. In the main report (Section 15), training on the longer 2021–2026 record made every model predict water temperature about 1 °C too cold: with no term for the year, the models learned the multi-year level, while 85% of the test rows are from 2025–26 and water year 2026 was 1.2 °C warmer than usual.

Option. `BENCHMARK_YEAR_EFFECT=wy` adds indicator columns for water years 2022–2026 (October–September; WY2021 is the reference, and the few hours after September 2026 count as WY2026) to every feature candidate of every model, so the inner loop cannot drop them. Their coefficients are estimated from the training stations, which are observed in the same years as the held-out station, so the held-out station's values never enter. Two runs were made with the five fast models and the graph model: on the current rows (a control, to separate the effect of the indicators from that of the extra data) and with the longer history.

2.1 On the current rows

Table 2: Water-year indicators on the current rows: paired ΔR^2 against the main benchmark.

Target	Model	Without	With	ΔR^2	Interval	Better/worse
Water temp.	Graph-Laplacian	0.863	0.859	−0.005	(−0.03, 0.01)	12/12
	OLS	0.863	0.862	−0.001	(−0.02, 0.01)	14/10
	Ridge	0.864	0.863	−0.001	(−0.02, 0.01)	13/11
	Elastic Net	0.863	0.866	+0.002	(−0.02, 0.02)	15/9
	Spline GAM	0.838	0.820	**−0.018**	(−0.03, −0.00)	7/17
DO	Graph-Laplacian	0.700	0.694	−0.005	(−0.03, 0.02)	6/11
	OLS	0.695	0.689	−0.006	(−0.03, 0.02)	9/8
	Ridge	0.701	0.701	+0.000	(−0.02, 0.03)	8/9
	Elastic Net	0.705	0.702	−0.003	(−0.02, 0.01)	8/9
	Spline GAM	0.708	0.695	**−0.013**	(−0.03, −0.00)	5/12
Spec. cond.	Graph-Laplacian	0.385	0.393	+0.009	(−0.08, 0.15)	12/12
	OLS	−0.099	−0.025	+0.074	(−0.14, 0.30)	10/14
	Ridge	−0.110	−0.073	+0.037	(−0.13, 0.16)	11/13
	Elastic Net	−0.109	−0.070	+0.039	(−0.12, 0.16)	11/13
	Spline GAM	0.022	−0.037	−0.059	(−0.26, 0.03)	9/15
pH	Graph-Laplacian	0.117	0.132	+0.015	(−0.00, 0.04)	10/9
	OLS	−0.014	−0.054	−0.040	(−0.09, 0.01)	8/11
	Ridge	0.003	−0.003	−0.006	(−0.04, 0.04)	8/11
	Elastic Net	0.023	−0.058	**−0.081**	(−0.12, −0.05)	4/15
	Spline GAM	0.012	−0.066	**−0.078**	(−0.15, −0.02)	6/13
Turbidity	Graph-Laplacian	0.104	0.186	+0.081	(−0.04, 0.47)	12/5
	OLS	0.136	0.194	**+0.058**	(0.01, 0.22)	13/4
	Ridge	0.136	0.196	**+0.060**	(0.01, 0.22)	13/4
	Elastic Net	0.136	0.193	**+0.056**	(0.01, 0.20)	13/4
	Spline GAM	0.091	0.135	+0.044	(−0.06, 0.14)	12/5

- **Water temperature, DO and specific conductance do not change** beyond noise for the graph and linear models. On the current rows the training and test stations cover the same

months, so there is little common offset to remove (mean signed station bias −0.05 before, +0.05 to +0.09 °C after). The Spline GAM loses a little on water temperature and DO.

- **Turbidity improves for every model**: the three linear models gain about 0.06 with intervals above zero (13 of 17 stations better), and the graph model gains 0.08 (0.104 → 0.186, interval crossing zero). Turbidity depends on runoff, which differs a lot from year to year, so a year level shared across stations is plausible. The best turbidity model is still gradient boosting (0.391), which was not part of this run.
- **pH gets worse** for Elastic Net and the Spline GAM (−0.08, intervals below zero) and does not change for the graph model (+0.015).
- The indicators were fixed in advance (water years, no tuning), so these are fair results. Because they help only turbidity, the option stays off by default.

2.2 With the longer history

Table 3: Water-year indicators with the 2021–2026 history: LOO R^2 of the main benchmark (Base), the history run of the main report (History) and the history run with water-year indicators (History + WY). Δ: History + WY against History, with the paired 95% interval.

Target	Model	Base	History	History + WY	Δ	Interval
Water temp.	Graph-Laplacian	0.863	0.816	0.813	−0.002	(−0.04, 0.03)
	OLS	0.863	0.840	0.856	+0.016	(−0.01, 0.04)
	Ridge	0.864	0.840	0.850	+0.009	(−0.02, 0.03)
	Elastic Net	0.863	0.841	0.851	+0.010	(−0.01, 0.03)
	Spline GAM	0.838	0.745	0.757	+0.012	(−0.05, 0.08)
DO	Graph-Laplacian	0.700	0.699	0.706	+0.007	(−0.02, 0.04)
	OLS	0.695	0.702	0.694	−0.008	(−0.03, 0.01)
	Ridge	0.701	0.708	0.698	−0.010	(−0.03, 0.01)
	Elastic Net	0.705	0.712	0.708	−0.004	(−0.02, 0.01)
	Spline GAM	0.708	0.719	0.715	−0.004	(−0.03, 0.01)
Spec. cond.	Graph-Laplacian	0.385	0.334	0.322	−0.013	(−0.15, 0.04)
	OLS	−0.099	−0.022	−0.727	−0.706	(−4.08, 0.08)
	Ridge	−0.110	−0.032	−0.227	−0.195	(−1.29, 0.08)
	Elastic Net	−0.109	−0.039	−0.124	−0.085	(−0.67, 0.07)
	Spline GAM	0.022	0.055	0.066	+0.011	(−0.12, 0.06)
pH	Graph-Laplacian	0.117	−0.037	−0.051	−0.013	(−0.08, 0.04)
	OLS	−0.014	−0.011	−0.273	**−0.262**	(−0.66, −0.03)
	Ridge	0.003	0.013	−0.173	**−0.186**	(−0.51, −0.02)
	Elastic Net	0.023	0.029	0.006	−0.023	(−0.08, 0.00)
	Spline GAM	0.012	0.009	0.006	−0.003	(−0.03, 0.03)
Turbidity	Graph-Laplacian	0.104	0.043	0.095	**+0.052**	(0.02, 0.13)
	OLS	0.136	0.032	0.100	**+0.068**	(0.01, 0.12)
	Ridge	0.136	0.031	0.095	+0.064	(−0.00, 0.10)
	Elastic Net	0.136	0.033	0.099	+0.067	(−0.01, 0.10)
	Spline GAM	0.091	0.085	0.128	+0.043	(−0.01, 0.09)

- **The cold offset is gone** (Table 4): the mean signed station bias of water temperature goes

Table 4: Water temperature: decomposition of the station-level error. Signed bias: mean over stations of the station-mean error (°C; without 09429490). Bias^2: mean squared station-mean error. Within: mean variance of the error about its station mean. R^2 without 09429490 in the last column.

Model	Run	Signed bias	Bias^2	Within	R^2 excl.
Graph-Laplacian	Base	−0.05	0.90	5.96	0.855
	History	−0.92	3.25	6.01	0.833
	History + WY	+0.04	3.09	6.28	0.840
Ridge	Base	−0.05	0.88	5.95	0.856
	History	−0.98	2.01	6.02	0.831
	History + WY	+0.01	1.27	6.29	0.841

from about −0.95 to +0.01–+0.04 °C, and the linear models gain about 0.01 (0.840 → 0.850–0.856).

- **But the history still does not beat the current rows.** Water temperature stays 0.01–0.05 below the main benchmark for every model, and the graph model does not change at all (0.816 → 0.813). Two things remain after the offset is removed: the station biases are still more spread out than in the main benchmark (bias^2 1.27 against 0.88 for Ridge; for the graph model most of the 3.09 is the 09429490 fold, whose bias is still about −6 °C), and the within-station error is larger (5.96 → 6.28). The seasonal coefficients fitted over six years describe the 2025–26 test period less well than those fitted on the same period.
- **Specific conductance and pH get much worse for OLS and Ridge.** OLS falls to −0.73 for specific conductance and −0.27 for pH, Ridge to −0.23 and −0.17 (pH intervals below zero). The specific-conductance loss comes from a few stations, mostly 09510000 (Verde below Bartlett, predicted 1,100–1,900 µS/cm too high) and the high-conductance station 96751415 (about 1,150 µS/cm too low). The graph model and the Spline GAM barely change.
- **Likely cause: the records cover different years at different stations.** In the history tables, some stations report only in the last one or two water years (09510000 from WY2024, 92116891 from WY2025, 75465016 only in WY2026) and others only in the first ones (133114453 and 134075067 stop in WY2022 for specific conductance; `results/history_coverage_by_wy.csv`). The linear models have no term for each station, so a water-year coefficient also absorbs which stations report in that year, and that mix changes with every held-out station. For specific conductance, whose level differs by a factor of five between stations, this swamps the year effect. The graph model has a level g_i for every station, so its year coefficients are estimated within stations, which is why it is unaffected. This explanation was not tested separately.
- **Turbidity improves against the history run** (+0.04 to +0.07, intervals above zero for the graph model and OLS), consistent with the current-row run, but stays at or below the main benchmark.
- **Conclusion.** With the year effect, the longer history no longer biases water temperature, but it still does not improve any target over the main benchmark, and with the uneven coverage it makes the linear models unstable. A fair use of the history would need the year effect estimated within stations (station-level offsets for every model, as the graph model has), or training restricted to stations that report in every year. Neither is run yet. The history option stays off by default.

3 Gradient boosting with the neighbour reading

The neighbour reading of the main report (Section 14: the mean same-hour anomaly of the target at the three nearest training stations) was run for gradient boosting, which was missing from that section. As before, every feature candidate is offered with and without it and the inner loop chooses.

Table 5: Gradient boosting with the neighbour reading: paired ΔR^2 against gradient boosting in the main benchmark. Folds: outer folds in which the inner loop chose a candidate with the reading.

Target	Without	With	ΔR^2	Interval	Better/worse	Folds
Water temp.	0.833	0.833	0.000	–	0/0	0/24
DO	0.733	0.729	−0.004	(−0.01, 0.00)	0/1	1/17
Spec. cond.	−0.094	−0.094	0.000	–	0/0	0/24
pH	−0.171	−0.123	+0.048	(−0.04, 0.17)	11/6	17/19
Turbidity	0.391	0.375	−0.016	(−0.10, 0.01)	8/8	16/17

- **Gradient boosting uses the reading only for pH and turbidity**, and never for water temperature or specific conductance, unlike the linear models, which chose it in every fold for every target but turbidity.
- **pH improves by** 0.05**, but stays below zero** (−0.123) and the interval crosses zero. The linear models gained 0.12–0.14 from the same reading and the graph model reached 0.178, so the reading does not make gradient boosting competitive for pH.
- **Turbidity is unchanged** (−0.016, within noise), so gradient boosting without the reading (0.391) remains the best turbidity model.

4 Conclusions so far

1. **A water-year effect helps turbidity only.** On the current rows it adds about 0.06 to every linear model (intervals above zero) and 0.08 to the graph model, changes nothing for water temperature, DO and specific conductance, and lowers pH for Elastic Net and the Spline GAM.
2. **The longer history is still not useful.** The year effect removes its 1 °C water-temperature offset, but no target beats the main benchmark, and because the stations' records cover different years the linear models become unstable for specific conductance and pH. The graph model, which has a level for every station, is unaffected by that problem but gains nothing either.
3. **The neighbour reading does not help gradient boosting** for any target beyond noise. Gradient boosting stays the best turbidity model without it.
4. The main benchmark (Table 1) remains the reference.

Next steps (proposed, not run). Turbidity with water-year indicators for gradient boosting and the Graph + GB hybrid; the history with station-level offsets for the linear models or restricted to stations that report in every year; clipping turbidity at the sensor maximum (3,800–4,000 FNU in the added years); a dam-tailwater indicator for the Lower Colorado.

Reproducibility. The runs come from `water_quality_model_benchmark.py` at commit `9f26e65` on the lab server, with `BENCHMARK_MODELS=Dummy,OLS,Ridge,Elastic Net,Spline GAM,Graph-Laplacian` and `BENCHMARK_YEAR_EFFECT=wy`, without and with `BENCHMARK_HISTORY=data/benchmark_data_history`, saved in `results/benchmark_results_wy/` and `results/benchmark_results_history_wy/`, and with `BENCHMARK_MODELS=Gradient Boosting` and `BENCHMARK_NEIGHBOUR=on`, saved in `results/benchmark_results_neighbour_gb/`. The server logs are `wy.log`, `hist_wy.log` and `nbr_gb.log`. The paired comparisons come from `compare_runs.py` against `results/benchmark_results_inner_fs_graph/` (graph model), `results/benchmark_results_inner_fs/` (other models) and the history run of the main report. They are saved as `results/wy_vs_base_*.csv`, `results/history_wy_vs_base_*.csv`, `results/history_wy_vs_history_*.csv` and `results/neighbour_vs_base_gradient_boosting.csv`. The error decomposition and the coverage table come from `year_effect_analysis.py` (seconds), saved in `results/year_effect_diagnostics.csv` and `results/history_coverage_by_wy.csv`.