Click here to access the guidelines/NICE algorithm N&WICS - TA389 - Pegylated liposomal doxorubicin hydrochloride (PLDH) monotherapy - Treating recurrent ovarian cancer Before providing patient identifiable data on this form, please confirm that the patient (or in the case of a minor or vulnerable adult with the parent/legal guardian/carer) has given appropriate explicit consent for sensitive personal information on this form to be passed to the CCG and/or CSU for processing this funding request and validating subsequent invoices. Consent given: If there is more than one NICE-approved treatment available, a discussion between the responsible clinician and the patient has taken place about the advantages and disadvantages of the treatments available. This has taken into consideration therapeutic need and whether or not the patient is likely to adhere to treatment. The most appropriate, least expensive, will be chosen (taking into account administration costs, dosage and price per dose) unless an order of preference is stated in the TAs. **Patient Practice** Trust: NHS No: Name: **Patient** Consultant **Practice** Hospital Making Postcode: No: Request: Patient's **Practice** Initials and Code: DoB: **Notification** Contact Email (@NHS.net account ONLY) name & Address: number: Provider: Start date ☐ Private ☐ NHS of Supplier: Sub-Type: ☐ Homecare ☐ Hospital requested treatment: (if applicable) By completing this form, you confirm that you intend to use the requested medicinal product described below as agreed in the local commissioning statement. Any requests which fall outside of this use will require an individual funding request (IFR). For support regarding IFRs, please contact: norfolkicd@nhs.net For support regarding the criteria listed below, please contact: norfolknontariff@nhs.net

Please indicate whether patient meets the following NICE criteria:	Please tick
1. Paclitaxel in combination with platinum or as monotherapy is recommended within its marketing authorisation as an option for treating recurrent ovarian cancer.	□Yes □No
2. Pegylated liposomal doxorubicin hydrochloride (PLDH) as monotherapy is recommended within its marketing authorisation as an option for treating recurrent ovarian cancer.	□Yes □No
3. PLDH in combination with platinum is recommended as an option for treating recurrent ovarian cancer.[1] [2]	□Yes □No
4. The following are not recommended within their marketing authorisations for treating the first recurrence of platinum-sensitive ovarian cancer:	
gemcitabine in combination with carboplatin	☐Yes ☐No
• trabectedin in combination with PLDH	
• topotecan. The appraisal committee was unable to make recommendations on the use of these technologies for treating platinum-sensitive ovarian cancer beyond the first recurrence.	
5. Topotecan is not recommended within its marketing authorisation for treating recurrent platinum-resistant or platinum-refractory ovarian cancer.	☐Yes ☐No
6. As per NICE guidance, treatment should normally be started with the least expensive drug (considering the dose required, price per dose and any additional administration costs).	
Please confirm that the choice of drug is considered the most cost-effective treatment for this individual patient, noting that alternative treatment options recommended by NICE may be lower in cost, including the use of biosimilar medications if applicable.	□Yes □No
Where an alternative lower cost treatment is available and has not been tried by the patient, please confirm that the clinical rationale for prescribing has been included in the patients' medical record for the purpose of auditing.	

7. The product is being used as described by local commission	ning position.	□Yes □No
I confirm that the patient meets the criteria for treatment		
Name of person completing:	Contact Details:	
Designation of person completing:	Date:	
Trust Authorising Pharmacist		
Name:		
Date:		