Click here to access the guidelines/NICE algorithm						
N&WICS - TA716 - Nivolumab with ipilimumab - Previously treated metastatic colorectal cancer with high microsatellite instability or mismatch repair deficiency						
Before providing patient identifiable parent/legal guardian/carer) has give CSU for processing this funding requirement of there is more than one NICE-approapout the advantages and disadvant patient is likely to adhere to treatment and price per dose) unless an order	data on this form, please configen appropriate explicit consent uest and validating subsequenced treatment available, a distages of the treatments availal ent. The most appropriate, least	irm that the patient (or in the for sensitive personal infont invoices. Consent given: scussion between the response. This has taken into corst expensive, will be chose	mation on this form  onsible clinician and asideration therapeur	to be passed to the the patient has tallic need and wheth	ken place her or not the	
Patient NHS No:	Trust:		Practice Name:			
Patient Hospital No:	Consultant Making Request:		Practice Postcode:			
Patient's Initials and DoB:			Practice Code:			
Notification Email Address:	(@NHS.ne	et account ONLY)	Contact name & number:			
Start date of requested treatment:	Committee	e □NHS care □Hospital	Sub-Type:	/A 🔽		
By completing this form, you confirm commissioning statement. Any requ For support regarding IFRs, please For support regarding the criteria lis	ests which fall outside of this contact: norfolkicd@nhs.net	use will require an individua			al	
Please indicate whether patient meets the following NICE criteria:			Please tick			
1. Nivolumab plus ipilimumab is recommended, within its marketing authorisation, as an option for treating metastatic colorectal cancer with high microsatellite instability (MSI) or mismatch repair (MMR) deficiency after fluoropyrimidine-based combination chemotherapy. It is recommended only if the company provides nivolumab and ipilimumab according to the commercial arrangement.						
2. As per NICE guidance, treatment should normally be started with the least expensive drug (considering the dose required, price per dose and any additional administration costs).  Please confirm that the choice of drug is considered the most cost-effective treatment for this individual patient, noting that alternative treatment options recommended by NICE may be lower in cost, including the use of biosimilar medications if applicable.  Where an alternative lower cost treatment is available and has not been tried by the patient, please confirm that the clinical rationale for prescribing has been included in the patients' medical record for the purpose of auditing.  □ Yes □ No						
3. The product is being used as described by local commissioning position.				□Yes □No		
I confirm that the patient meet	s the criteria for treatment					
Name of person completing:  Contact Details:						
Designation of person completing: Date:						

Trust Authorising Pharmacist

Name:

Date:	