## Click here to access the guidelines/NICE algorithm

	- Onek here to decess the galdelines/Hot algoridin				
N&WICS - TA264 - Alteplase - Ischaemic stroke (acute)					
Before providing patient identifiable parent/legal guardian/carer) has give CSU for processing this funding req	en appropriate explicit consent	for sensitive personal infor	mation on this form		
If there is more than one NICE-approabout the advantages and disadvant patient is likely to adhere to treatme and price per dose) unless an order	ages of the treatments availalent. The most appropriate, leas	ble. This has taken into con st expensive, will be chose	sideration therapeut	tic need and wheth	er or not the
Patient		1A3. [_]	Drestics		
NHS No:	Trust:		Practice Name:		
Patient Hospital No:	Consultant Making Request:		Practice Postcode:		
Patient's Initials and DoB:			Practice Code:		
Notification Email Address:	(@NHS.ne	et account ONLY)	Contact name & number:		
Start date of requested treatment:	0	e □NHS care □Hospital	Sub-Type:	/A 🔽	
By completing this form, you confirm commissioning statement. Any requ					ıl
		use will require air individue	arranamy request (ii	TK).	
For support regarding IFRs, please	contact: norfolkicd@nhs.net				
For support regarding the criteria lis	ted below, please contact: nor	folknontariff@nhs.net			
Please indicate whether patient meets the following NICE criteria:					
	it meets the following Nice	criteria:		Please tick	
Alteplase is recommended with if:			ic stroke in adults		
Alteplase is recommended with	hin its marketing authorisation	for treating acute ischaem		Please tick  ☐ Yes ☐ No	
Alteplase is recommended with if:	hin its marketing authorisation possible within 4.5 hours of c	n for treating acute ischaem			
Alteplase is recommended with if:     treatment is started as early as	possible within 4.5 hours of cen excluded by appropriate in ent should normally be started	n for treating acute ischaem onset of stroke symptoms, a maging techniques.	and		
Alteplase is recommended with if:     treatment is started as early as     intracranial haemorrhage has be     As per NICE guidance, treatment	possible within 4.5 hours of commendation possible within 4.5 hours of commendation and any additional administrative and any additional administrative drug is considered the most catment options recommended	n for treating acute ischaem onset of stroke symptoms, a maging techniques.  I with the least expensive di ation costs).  cost-effective treatment for	rug (considering		
1. Alteplase is recommended with if:     • treatment is started as early as     • intracranial haemorrhage has be     2. As per NICE guidance, treatmenthe dose required, price per dose  Please confirm that the choice of patient, noting that alternative treatments.	possible within 4.5 hours of commended and any additional administrative of the commended policable.	on for treating acute ischaem onset of stroke symptoms, a maging techniques.  I with the least expensive diation costs).  Cost-effective treatment for by NICE may be lower in control by the patier	rug (considering this individual cost, including the	□Yes □No	
1. Alteplase is recommended with if:  • treatment is started as early as  • intracranial haemorrhage has be  2. As per NICE guidance, treatmenthe dose required, price per dose  Please confirm that the choice of patient, noting that alternative treuse of biosimilar medications if all where an alternative lower cost to that the clinical rationale for presentations.	possible within 4.5 hours of commended and any additional administration drug is considered the most catment options recommended opticable.	onset of stroke symptoms, a maging techniques.  I with the least expensive diation costs).  cost-effective treatment for by NICE may be lower in cost been tried by the patier ne patients' medical record	rug (considering this individual cost, including the	□Yes □No	
1. Alteplase is recommended with if:  • treatment is started as early as  • intracranial haemorrhage has be  2. As per NICE guidance, treatmenthe dose required, price per dose  Please confirm that the choice of patient, noting that alternative treuse of biosimilar medications if all where an alternative lower cost to that the clinical rationale for presented.	possible within 4.5 hours of commendation and any additional administration and additional administration and additional administration and additional administration and administration administration and administration administration and a	onset of stroke symptoms, a maging techniques.  I with the least expensive diation costs).  cost-effective treatment for by NICE may be lower in cost been tried by the patier ne patients' medical record	rug (considering this individual cost, including the	☐Yes ☐No	
1. Alteplase is recommended with if:  • treatment is started as early as  • intracranial haemorrhage has be  2. As per NICE guidance, treatmenthe dose required, price per dose  Please confirm that the choice of patient, noting that alternative treuse of biosimilar medications if a Where an alternative lower cost to that the clinical rationale for presented auditing.  3. The product is being used as of the comments of the product of the	possible within 4.5 hours of commendation and any additional administration and additional administration and additional administration and additional administration and administration administration and administration administration and a	onset of stroke symptoms, a maging techniques.  I with the least expensive diation costs).  cost-effective treatment for by NICE may be lower in cost been tried by the patier ne patients' medical record	rug (considering this individual cost, including the	☐Yes ☐No	
1. Alteplase is recommended with if:  • treatment is started as early as  • intracranial haemorrhage has be  2. As per NICE guidance, treatmenthe dose required, price per dose  Please confirm that the choice of patient, noting that alternative treuse of biosimilar medications if all where an alternative lower cost to that the clinical rationale for presequential.  3. The product is being used as a light confirm that the patient meets.	possible within 4.5 hours of commendation and any additional administration and administration	n for treating acute ischaem onset of stroke symptoms, a maging techniques.  I with the least expensive diation costs).  cost-effective treatment for by NICE may be lower in ont been tried by the patience patients' medical record in hing position.	rug (considering this individual cost, including the	☐Yes ☐No	
1. Alteplase is recommended with if:  • treatment is started as early as  • intracranial haemorrhage has be  2. As per NICE guidance, treatmenthe dose required, price per dose  Please confirm that the choice of patient, noting that alternative treuse of biosimilar medications if all where an alternative lower cost to that the clinical rationale for present auditing.  3. The product is being used as of a confirm that the patient meets.  Name of person completing:	possible within 4.5 hours of commendation and any additional administration and administration	nonset of stroke symptoms, a maging techniques.  with the least expensive diation costs).  cost-effective treatment for by NICE may be lower in contract been tried by the patient patients' medical recording position.	rug (considering this individual cost, including the	☐Yes ☐No	

Date:	