## Click here to access the guidelines/NICE algorithm N&WICS - TA479 - Reslizumab, as an add-on therapy - Severe eosinophilic asthma Before providing patient identifiable data on this form, please confirm that the patient (or in the case of a minor or vulnerable adult with the parent/legal guardian/carer) has given appropriate explicit consent for sensitive personal information on this form to be passed to the CCG and/or CSU for processing this funding request and validating subsequent invoices. Consent given: If there is more than one NICE-approved treatment available, a discussion between the responsible clinician and the patient has taken place about the advantages and disadvantages of the treatments available. This has taken into consideration therapeutic need and whether or not the patient is likely to adhere to treatment. The most appropriate, least expensive, will be chosen (taking into account administration costs, dosage and price per dose) unless an order of preference is stated in the TAs. Patient **Practice** Trust: NHS No: Name: **Patient** Consultant **Practice** Hospital Making Postcode: No. Request: Patient's **Practice** Initials and Code: DoB: Notification Contact Email (@NHS.net account ONLY) name & Address: number: Provider: Start date ☐ Private ☐ NHS of Supplier: Sub-Type: ☐ Homecare ☐ Hospital requested treatment: (if applicable) By completing this form, you confirm that you intend to use the requested medicinal product described below as agreed in the local commissioning statement. Any requests which fall outside of this use will require an individual funding request (IFR). For support regarding IFRs, please contact: norfolkicd@nhs.net For support regarding the criteria listed below, please contact: norfolknontariff@nhs.net Please indicate whether patient meets the following NICE criteria: Please tick 1. Reslizumab, as an add-on therapy, is recommended as an option for the treatment of severe eosinophilic asthma that is inadequately controlled in adults despite maintenance therapy with high-dose inhaled corticosteroids plus another drug, only if: • the blood eosinophil count has been recorded as 400 cells per microlitre or more □Yes □No • the person has had 3 or more severe asthma exacerbations needing systemic corticosteroids in the past 12 months and • the company provides reslizumab with the discount agreed in the patient access scheme. 2. At 12 months: • stop reslizumab if the asthma has not responded adequately or • continue reslizumab if the asthma has responded adequately and assess response each year. An adequate response is defined as: □Yes □No • a clinically meaningful reduction in the number of severe exacerbations needing systemic corticosteroids • a clinically significant reduction in continuous oral corticosteroid use while maintaining or improving asthma control. 3. As per NICE guidance, treatment should normally be started with the least expensive drug (considering the dose required, price per dose and any additional administration costs). Please confirm that the choice of drug is considered the most cost-effective treatment for this individual

patient, noting that alternative treatment options recommended by NICE may be lower in cost, including the

use of biosimilar medications if applicable.

☐Yes ☐No

Where an alternative lower cost treatment is available and has that the clinical rationale for prescribing has been included in thauditing.		
4. The product is being used as described by local commissioning position.		□Yes □No
I confirm that the patient meets the criteria for treatment		
Name of person completing:	Contact Details:	
Designation of person completing:	Date:	
Trust Authorising Pharmacist	<u> </u>	
Name:		
Date:		