Click here to access the guidelines/NICE algorithm N&WICS - TA391 - Cabazitaxel in combination with prednisone or prednisolone - Treating hormone-relapsed metastatic prostate cancer treated with docetaxel Before providing patient identifiable data on this form, please confirm that the patient (or in the case of a minor or vulnerable adult with the parent/legal guardian/carer) has given appropriate explicit consent for sensitive personal information on this form to be passed to the CCG and/or CSU for processing this funding request and validating subsequent invoices. Consent given: If there is more than one NICE-approved treatment available, a discussion between the responsible clinician and the patient has taken place about the advantages and disadvantages of the treatments available. This has taken into consideration therapeutic need and whether or not the patient is likely to adhere to treatment. The most appropriate, least expensive, will be chosen (taking into account administration costs, dosage and price per dose) unless an order of preference is stated in the TAs. **Patient Practice** Trust: NHS No: Name: Consultant **Patient Practice** Hospital Making Postcode: No: Request: Patient's **Practice** Initials and Code: DoR: **Notification** Contact Email (@NHS.net account ONLY) name & Address: number: Provider: Start date ☐ Private ☐ NHS of Supplier: Sub-Type: ☐ Homecare ☐ Hospital requested treatment: (if applicable) By completing this form, you confirm that you intend to use the requested medicinal product described below as agreed in the local commissioning statement. Any requests which fall outside of this use will require an individual funding request (IFR). For support regarding IFRs, please contact: norfolkicd@nhs.net For support regarding the criteria listed below, please contact: norfolknontariff@nhs.net Please indicate whether patient meets the following NICE criteria: Please tick 1. Cabazitaxel in combination with prednisone or prednisolone is recommended as an option for treating metastatic hormone-relapsed prostate cancer in people whose disease has progressed during or after docetaxel chemotherapy, only if: • the person has an eastern cooperative oncology group (ECOG) performance status of 0 or 1 ☐ Yes ☐ No • the person has had 225 mg/m2 or more of docetaxel • treatment with cabazitaxel is stopped when the disease progresses or after a maximum of 10 cycles (whichever happens first) • the company provides cabazitaxel according to the commercial arrangement. 2. When using ECOG performance status, healthcare professionals should take into account any physical, sensory or learning disabilities, or communication difficulties that could affect ECOG performance status ☐ Yes ☐ No and make any adjustments they consider appropriate. 3. As per NICE guidance, treatment should normally be started with the least expensive drug (considering the dose required, price per dose and any additional administration costs). Please confirm that the choice of drug is considered the most cost-effective treatment for this individual patient, noting that alternative treatment options recommended by NICE may be lower in cost, including the ☐Yes ☐No

use of biosimilar medications if applicable.

auditing.

Where an alternative lower cost treatment is available and has not been tried by the patient, please confirm that the clinical rationale for prescribing has been included in the patients' medical record for the purpose of

4. The product is being used as described by local commissioning position.

□Yes □No

I confirm that the patient meets the criteria for treatment	
Name of person completing:	Contact Details:
Designation of person completing:	Date:
Trust Authorising Pharmacist	
Name: Date:	