Click here to access the guidelines/NICE algorithm N&WICS - TA339 - Omalizumab - Previously treated chronic spontaneous urticaria Before providing patient identifiable data on this form, please confirm that the patient (or in the case of a minor or vulnerable adult with the parent/legal guardian/carer) has given appropriate explicit consent for sensitive personal information on this form to be passed to the CCG and/or CSU for processing this funding request and validating subsequent invoices. Consent given: If there is more than one NICE-approved treatment available, a discussion between the responsible clinician and the patient has taken place about the advantages and disadvantages of the treatments available. This has taken into consideration therapeutic need and whether or not the patient is likely to adhere to treatment. The most appropriate, least expensive, will be chosen (taking into account administration costs, dosage and price per dose) unless an order of preference is stated in the TAs. **Patient Practice** Trust: NHS No: Name: **Patient** Consultant **Practice** Hospital Making Postcode: No: Request: Patient's **Practice** Initials and Code: DoB: **Notification** Contact Email (@NHS.net account ONLY) name & Address: number: Provider: Start date ☐ Private ☐ NHS of Supplier: Sub-Type: ☐ Homecare ☐ Hospital requested N/A treatment: (if applicable) By completing this form, you confirm that you intend to use the requested medicinal product described below as agreed in the local commissioning statement. Any requests which fall outside of this use will require an individual funding request (IFR). For support regarding IFRs, please contact: norfolkicd@nhs.net For support regarding the criteria listed below, please contact: norfolknontariff@nhs.net Please tick Please indicate whether patient meets the following NICE criteria:

. I sales maistre mistre parent mosts the renorming most entertain	i iodoo tiok
1. Omalizumab is recommended as an option as add-on therapy for treating severe chronic spontaneous urticaria in adults and young people aged 12 years and over only if:	
• the severity of the condition is assessed objectively using a weekly urticaria activity score of 28 or more for at least two weeks, over a 4 week period.	
• the person's condition has not responded to standard treatment with H1-antihistamines and leukotriene receptor antagonists	
• omalizumab is stopped at or before the fourth dose if the condition has not responded	□Yes □No
• omalizumab is stopped at the end of a course of treatment (6 doses) if the condition has responded, to establish whether the condition has gone into spontaneous remission, and is restarted only if the condition relapses	
• omalizumab is administered under the management of a secondary care specialist in dermatology, immunology or allergy	
• the company provides omalizumab with the discount agreed in the patient access scheme.	
2. As per NICE guidance, treatment should normally be started with the least expensive drug (considering the dose required, price per dose and any additional administration costs).	
Please confirm that the choice of drug is considered the most cost-effective treatment for this individual patient, noting that alternative treatment options recommended by NICE may be lower in cost, including the use of biosimilar medications if applicable.	□Yes □No
Where an alternative lower cost treatment is available and has not been tried by the patient, please confirm that the clinical rationale for prescribing has been included in the patients' medical record for the purpose of auditing.	

I confirm that the patient meets the criteria for treatment	
Name of person completing:	Contact Details:
Designation of person completing:	Date:
Trust Authorising Pharmacist	
Name: Date:	