Click here to access the guidelines/NICE algorithm							
N&WICS - TA335 - Rivaroxaban			n plus clopidogrel or as nt of acute coronary sy		nting adverse out	comes after	
Before providing patient identifiable of parent/legal guardian/carer) has give CSU for processing this funding requ	en appropriate explici	it consent	t for sensitive personal inf	formation on this form			
If there is more than one NICE-approabout the advantages and disadvant patient is likely to adhere to treatme and price per dose) unless an order	ages of the treatmer nt. The most approp	nts availal oriate, leas	ble. This has taken into c st expensive, will be chos	onsideration therapeu	tic need and wheth	ner or not the	
Patient NHS No:	Trust:	eu iii tile	IA5. Ц	Practice Name:			
Patient Hospital No:	Consultant Making Request:			Practice Postcode:			
Patient's Initials and DoB:				Practice Code:			
Notification Email Address:	(@NHS.ne	et account ONLY)	Contact name & number:			
Start date of requested treatment:	Provider: Supplier:		e □NHS care □Hospital	Sub-Type:	/A 🔽		
For support regarding IFRs, please For support regarding the criteria list	contact: norfolkicd@	nhs.net	·	lual funding request (I	FR).		
Please indicate whether patient meets the following NICE criteria: 1. Rivaroxaban is recommended as an option within its marketing authorisation, in combination with aspirin plus clopidogrel or aspirin alone, for preventing atherothrombotic events in people who have had an acute coronary syndrome with elevated cardiac biomarkers.					Please tick		
2. Clinicians should carefully assess the person's risk of bleeding before treatment with rivaroxaban is started. The decision to start treatment should be made after an informed discussion between the clinician and the patient about the benefits and risks of rivaroxaban in combination with aspirin plus clopidogrel or with aspirin alone, compared with aspirin plus clopidogrel or aspirin alone.					□Yes □No		
3. A decision on continuation of treatment should be taken no later than 12 months after starting treatment. Clinicians should regularly reassess the relative benefits and risks of continuing treatment with rivaroxaban and discuss them with the patient.					☐Yes ☐No		
As per NICE guidance, treatment the dose required, price per dose				drug (considering			
Please confirm that the choice of drug is considered the most cost-effective treatment for this individual patient, noting that alternative treatment options recommended by NICE may be lower in cost, including the use of biosimilar medications if applicable.					□Yes □No		
Where an alternative lower cost to that the clinical rationale for presonauditing.							
I confirm that the patient meets	s the criteria for tre	eatment				l	
Name of person completing:			Contact Details:			1	
Designation of person completing:					l		

Trust Authorising Pharmacist	
Name:	
Date:	