

[Click here to access the guidelines/NICE algorithm](#)

N&WICS - TA195 - Etanercept - Rheumatoid Arthritis

Before providing patient identifiable data on this form, please confirm that the patient (or in the case of a minor or vulnerable adult with the parent/legal guardian/carer) has given appropriate explicit consent for sensitive personal information on this form to be passed to the CCG and/or CSU for processing this funding request and validating subsequent invoices. Consent given: ☐

If there is more than one NICE-approved treatment available, a discussion between the responsible clinician and the patient has taken place about the advantages and disadvantages of the treatments available. This has taken into consideration therapeutic need and whether or not the patient is likely to adhere to treatment. The most appropriate, least expensive, will be chosen (taking into account administration costs, dosage and price per dose) unless an order of preference is stated in the TAs. ☐

Patient NHS No:	Trust:	Practice Name:
Patient Hospital No: <input type="text"/>	Consultant Making Request: <input type="text"/>	Practice Postcode:
Patient's Initials and DoB:		Practice Code:
Notification Email Address: <input type="text"/> (@NHS.net account ONLY)		Contact name & number: <input type="text"/>
Start date of requested treatment: <input type="text"/>	Provider: <input type="checkbox"/> Private <input type="checkbox"/> NHS Supplier: <input type="checkbox"/> Homecare <input type="checkbox"/> Hospital	Sub-Type: <input type="text"/> N/A <input type="text"/> <input type="text"/> (if applicable)

By completing this form, you confirm that you intend to use the requested medicinal product described below as agreed in the local commissioning statement. Any requests which fall outside of this use will require an individual funding request (IFR).

For support regarding IFRs, please contact: norfolkicd@nhs.net

For support regarding the criteria listed below, please contact: norfolknontariff@nhs.net

Please indicate whether patient meets the following NICE criteria:	Please tick
1. Rituximab in combination with methotrexate is recommended as an option for the treatment of adults with severe active rheumatoid arthritis who have had an inadequate response to, or are intolerant of, other disease-modifying anti-rheumatic drugs (DMARDs), including at least one tumour necrosis factor (TNF) inhibitor. Treatment with rituximab should be given no more frequently than every 6 months.	<input type="checkbox"/> Yes <input type="checkbox"/> No
2. Treatment with rituximab in combination with methotrexate should be continued only if there is an adequate response following initiation of therapy and if an adequate response is maintained following retreatment with a dosing interval of at least 6 months. An adequate response is defined as an improvement in disease activity score (DAS28) of 1.2 points or more.	<input type="checkbox"/> Yes <input type="checkbox"/> No
3. Adalimumab, etanercept, infliximab and abatacept, each in combination with methotrexate, are recommended as treatment options only for adults with severe active rheumatoid arthritis who have had an inadequate response to, or have an intolerance of, other DMARDs, including at least one TNF inhibitor, and who cannot receive rituximab therapy because they have a contraindication to rituximab, or when rituximab is withdrawn because of an adverse event.	<input type="checkbox"/> Yes <input type="checkbox"/> No
4. Adalimumab monotherapy and etanercept monotherapy are recommended as treatment options for adults with severe active rheumatoid arthritis who have had an inadequate response to, or have an intolerance of, other DMARDs, including at least one TNF inhibitor, and who cannot receive rituximab therapy because they have a contraindication to methotrexate, or when methotrexate is withdrawn because of an adverse event.	<input type="checkbox"/> Yes <input type="checkbox"/> No
5. Treatment with adalimumab, etanercept, infliximab and abatacept should be continued only if there is an adequate response (as defined in 1.2) 6 months after initiation of therapy. Treatment should be monitored, with assessment of DAS28, at least every 6 months and continued only if an adequate response is maintained.	<input type="checkbox"/> Yes <input type="checkbox"/> No
6. When using DAS28, healthcare professionals should take into account any physical, sensory or learning disabilities, communication difficulties, or disease characteristics that could adversely affect patient assessment and make any adjustments they consider appropriate.	<input type="checkbox"/> Yes <input type="checkbox"/> No
7. A team experienced in the diagnosis and treatment of rheumatoid arthritis and working under the supervision of a rheumatologist should initiate, supervise and assess response to treatment with rituximab,	<input type="checkbox"/> Yes <input type="checkbox"/> No

adalimumab, etanercept, infliximab or abatacept.

I confirm that the patient meets the criteria for treatment

Name of person completing:

Contact Details:

Designation of person completing:

Date:

Trust Authorising Pharmacist

Name:

Date: