Click here to access the guidelines/NICE algorithm

NAWICS - TA757 - Cabotegravir with rilniviring - Treating HIV-1					
N&WICS - TA757 - Cabotegravir with rilpivirine - Treating HIV-1 Before providing patient identifiable data on this form, please confirm that the patient (or in the case of a minor or vulnerable adult with the parent/legal guardian/carer) has given appropriate explicit consent for sensitive personal information on this form to be passed to the CCG and/or CSU for processing this funding request and validating subsequent invoices. Consent given:					
If there is more than one NICE-approved treatment available, a discussion between the responsible clinician and the patient has taken place about the advantages and disadvantages of the treatments available. This has taken into consideration therapeutic need and whether or not the patient is likely to adhere to treatment. The most appropriate, least expensive, will be chosen (taking into account administration costs, dosage and price per dose) unless an order of preference is stated in the TAs.					
Patient	Trust:		actice		
NHS No:	Consultant		Name:		
Hospital No:	Making Request:		actice tcode:		
Patient's Initials and DoB:		Pr	Practice Code:		
Notification Email Address:	(@NHS.net acco	ount ONLY) na	ontact ame & imber:		
Start date	Provider:	NHS			
of requested	Supplier: Homecare	☐ Hospital Sub	-Type:	/A 🔽	
treatment:		(if app	olicable)	/A 🛂	
By completing this form, you confirm commissioning statement. Any requ For support regarding IFRs, please	ests which fall outside of this use v contact: norfolkicd@nhs.net				41
For support regarding the criteria its	ted below, please contact: norfolkn	ontariff@nhs.net			
	ted below, please contact: norfolkn			Please tick	
Please indicate whether patier	·	eria:	treating	Please tick	
Please indicate whether patier 1. Cabotegravir with rilpivirine is r HIV-1 infection in adults:	nt meets the following NICE crit	eria: authorisation, as an option for	-		
Please indicate whether patier 1. Cabotegravir with rilpivirine is r HIV-1 infection in adults: • with virological suppression (HIV • without any evidence of viral res	nt meets the following NICE crit ecommended, within its marketing /-1 RNA fewer than 50 copies/ml) consistance to, and no previous virologic integrase inhibitors. It is recomme	eria: authorisation, as an option for on a stable antiretroviral regime cal failure with, any non-nucleo	en and	Please tick ☐ Yes ☐ No	
Please indicate whether patier 1. Cabotegravir with rilpivirine is r HIV-1 infection in adults: • with virological suppression (HIV • without any evidence of viral res reverse transcriptase inhibitors or according to the commercial arrai	nt meets the following NICE crit ecommended, within its marketing /-1 RNA fewer than 50 copies/ml) consistance to, and no previous virologic integrase inhibitors. It is recomme	authorisation, as an option for on a stable antiretroviral regime cal failure with, any non-nucleous nded only if the company prove the least expensive drug (con	en and oside vides it		
Please indicate whether patier 1. Cabotegravir with rilpivirine is r HIV-1 infection in adults: • with virological suppression (HIV • without any evidence of viral res reverse transcriptase inhibitors or according to the commercial arrai 2. As per NICE guidance, treatme the dose required, price per dose Please confirm that the choice of	nt meets the following NICE crit recommended, within its marketing /-1 RNA fewer than 50 copies/ml) of the copies integrase inhibitors. It is recommended in the should normally be started with and any additional administration of the copies considered the most cost-atment options recommended by N	authorisation, as an option for on a stable antiretroviral regime cal failure with, any non-nucleous nded only if the company prove the least expensive drug (concosts).	en and poside vides it sidering		
Please indicate whether patier 1. Cabotegravir with rilpivirine is r HIV-1 infection in adults: • with virological suppression (HIV) • without any evidence of viral res reverse transcriptase inhibitors or according to the commercial arrail 2. As per NICE guidance, treatmenthe dose required, price per dose Please confirm that the choice of patient, noting that alternative treatments of biosimilar medications if all where an alternative lower cost to	nt meets the following NICE crit recommended, within its marketing /-1 RNA fewer than 50 copies/ml) of the copies integrase inhibitors. It is recommended in the should normally be started with and any additional administration of the copies considered the most cost-atment options recommended by N	authorisation, as an option for on a stable antiretroviral regime cal failure with, any non-nucleous nded only if the company prove the least expensive drug (concosts). Effective treatment for this individed may be lower in cost, income tried by the patient, pleas	en and poside rides it sidering ridual luding the e confirm	□Yes □No	
Please indicate whether patier 1. Cabotegravir with rilpivirine is r HIV-1 infection in adults: • with virological suppression (HIV) • without any evidence of viral res reverse transcriptase inhibitors or according to the commercial arrail 2. As per NICE guidance, treatmenthe dose required, price per dose Please confirm that the choice of patient, noting that alternative tresuse of biosimilar medications if all where an alternative lower cost to that the clinical rationale for presequiditing.	nt meets the following NICE crit recommended, within its marketing /-1 RNA fewer than 50 copies/ml) of the copies integrase inhibitors. It is recommended in the same and any additional administration drug is considered the most cost-catment options recommended by Nopplicable.	authorisation, as an option for on a stable antiretroviral regime cal failure with, any non-nucleous nded only if the company prove the least expensive drug (concosts). Effective treatment for this individed may be lower in cost, income tried by the patient, pleas tients' medical record for the p	en and poside rides it sidering ridual luding the e confirm	□Yes □No	
Please indicate whether patier 1. Cabotegravir with rilpivirine is r HIV-1 infection in adults: • with virological suppression (HIV) • without any evidence of viral res reverse transcriptase inhibitors or according to the commercial arrail 2. As per NICE guidance, treatmenthe dose required, price per dose Please confirm that the choice of patient, noting that alternative tresuse of biosimilar medications if all where an alternative lower cost to that the clinical rationale for presequiditing.	nt meets the following NICE crit recommended, within its marketing /-1 RNA fewer than 50 copies/ml) of sistance to, and no previous virologic integrase inhibitors. It is recommengement. rent should normally be started with and any additional administration drug is considered the most cost-atment options recommended by Nopplicable. reatment is available and has not be cribing has been included in the particles.	authorisation, as an option for on a stable antiretroviral regime cal failure with, any non-nucleous nded only if the company prove the least expensive drug (concosts). Effective treatment for this individed may be lower in cost, income tried by the patient, pleas tients' medical record for the p	en and poside rides it sidering ridual luding the e confirm	☐Yes ☐No	
Please indicate whether patier 1. Cabotegravir with rilpivirine is r HIV-1 infection in adults: • with virological suppression (HIV) • without any evidence of viral res reverse transcriptase inhibitors or according to the commercial arrail 2. As per NICE guidance, treatmenthe dose required, price per dose Please confirm that the choice of patient, noting that alternative treatments of biosimilar medications if all where an alternative lower cost to that the clinical rationale for presentations. 3. The product is being used as of the service of the product of the product of the patients.	nt meets the following NICE criteria to the commended, within its marketing of the commended, within its marketing of the commended, within its marketing of the commended to the commended in the commended by North and any additional administration of the commended by North and considered the most cost-catment options recommended by North and commended by North and	authorisation, as an option for on a stable antiretroviral regime cal failure with, any non-nucleous nded only if the company prove the least expensive drug (concosts). Effective treatment for this individed may be lower in cost, income tried by the patient, pleas tients' medical record for the p	en and poside rides it sidering ridual luding the e confirm	☐Yes ☐No	
Please indicate whether patier 1. Cabotegravir with rilpivirine is r HIV-1 infection in adults: • with virological suppression (HIV) • without any evidence of viral res reverse transcriptase inhibitors or according to the commercial arrance. 2. As per NICE guidance, treatmenthe dose required, price per dose Please confirm that the choice of patient, noting that alternative treatments of biosimilar medications if all Where an alternative lower cost to that the clinical rationale for present auditing. 3. The product is being used as confirm that the patient meets.	nt meets the following NICE critical commended, within its marketing of 1/21 RNA fewer than 50 copies/ml) of the sistance to, and no previous virologic integrase inhibitors. It is recommended in the sand any additional administration of drug is considered the most costatment options recommended by Nopplicable. The same of the s	authorisation, as an option for on a stable antiretroviral regime cal failure with, any non-nucleunded only if the company provethe least expensive drug (concosts). Effective treatment for this individed may be lower in cost, income tried by the patient, pleas tients' medical record for the proposition.	en and poside rides it sidering ridual luding the e confirm	☐Yes ☐No	

Name:	
Date:	