Click here to access the guidelines/NICE algorithm						
	N&WICS - TA	134 - Infliximab (2nd line) - F	Plaque psoriasis			
Before providing patient identifiable	data on this form, pl	ease confirm that the patient (c	or in the case of a minor or	vulnerable adult w	rith the	
parent/legal guardian/carer) has give				to be passed to the	e CCG and/or	
CSU for processing this funding req	uest and validating s	subsequent invoices. Consent of	given: 🔲			
If there is more than one NICE-appro	oved treatment avail	able, a discussion between the	responsible clinician and	the patient has tak	en place	
about the advantages and disadvant	tages of the treatme	nts available. This has taken in	nto consideration therapeut	ic need and wheth	er or not the	
patient is likely to adhere to treatme			chosen (taking into accou	nt administration c	osts, dosage	
and price per dose) unless an order	of preference is stat	ted in the TAs				
Patient NHS No:	Trust:		Practice Name:			
Patient	Consultant		Practice			
Hospital	Making		Postcode:			
No:	Request:					
Patient's Initials and			Practice			
DoB:			Code:			
Notification			Contact			
Email	(@NHS.net account ONLY)		name &			
Address:	T		number:			
Start date	Provider:	☐ Private ☐ NHS				
of	Supplier:		Sub-Type:			
requested treatment:	0.00	☐ Homecare ☐ Hospital	N/	Ά 🔽		
treatment.			(if applicable)			
For support regarding IFRs, please For support regarding the criteria lis			et			
Please indicate whether patier	nt meets the follow	ving NICE criteria:		Please tick		
Infliximab, within its licensed in	dications is recomm	monded as a treatment entire	for adults with plague			
psoriasis only when the following		nended as a treatment option	ioi addits with plaque			
• The disease is very severe as defined by a total Psoriasis Area Severity Index (PASI) of 20 or more and a						
Dermatology Life Quality Index (DLQI) of more than 18.						
The psoriasis has failed to respond to standard systemic therapies such as ciclosporin, methotrexate or						
PUVA (psoralen and long-wave u these treatments.	Itraviolet radiation),	or the person is intolerant to or	has a contraindication to			
2. Infliximab treatment should be						
adequate response to treatment v	within to weeks. An	adequate response is defined	as either.			
• a 75% reduction in the PASI score from when treatment started (PASI 75) or						
• a 50% reduction in the PASI scored.	ore (PASI 50) and a	five-point reduction in the DLQ	I from when treatment			
3. When using the DLQI healthcat patient's disabilities (such as phy	•		•			
reaching conclusions on the seve	erity of plaque psoria	asis. In such cases healthcare	professionals should	☐Yes ☐No		
ensure that their use of the DLQI should apply in the context of a d						
section 1.2.	COSION ADOUL WHELH	ici to continue the use of the u	rug in accordance with			
	ant aboutd same	he storted with the least correct	univo drug (noncidarine			
4. As per NICE guidance, treatment the dose required, price per dose			isive arug (considering			
	•	,				
Please confirm that the choice of patient, noting that alternative treations.	-					
- Lugueni, nomo mai anemanye ne						

Where an alternative lower cost treatment is available and has not been tried by the patient, please confirm that the clinical rationale for prescribing has been included in the patients' medical record for the purpose of auditing.					
I confirm that the patient meets the criteria for treatment					
Name of person completing:	Contact Details:				
Designation of person completing:	Date:				
Trust Authorising Pharmacist					
Name: Date:					