## Click here to access the guidelines/NICE algorithm N&WICS - TA315 - Canagliflozin in combination with metformin and a sulfonylurea/thiazolidinedione (triple therapy) - Type 2 diabetes Before providing patient identifiable data on this form, please confirm that the patient (or in the case of a minor or vulnerable adult with the parent/legal guardian/carer) has given appropriate explicit consent for sensitive personal information on this form to be passed to the CCG and/or CSU for processing this funding request and validating subsequent invoices. Consent given: If there is more than one NICE-approved treatment available, a discussion between the responsible clinician and the patient has taken place about the advantages and disadvantages of the treatments available. This has taken into consideration therapeutic need and whether or not the patient is likely to adhere to treatment. The most appropriate, least expensive, will be chosen (taking into account administration costs, dosage and price per dose) unless an order of preference is stated in the TAs. **Patient Practice** Trust: NHS No: Name: Consultant **Patient Practice** Hospital Making Postcode: No: Request: Patient's **Practice** Initials and Code: DoR: Notification Contact Email (@NHS.net account ONLY) name & Address: number: Provider: Start date ☐ Private ☐ NHS of Supplier: Sub-Type: ☐ Homecare ☐ Hospital requested treatment: (if applicable) By completing this form, you confirm that you intend to use the requested medicinal product described below as agreed in the local commissioning statement. Any requests which fall outside of this use will require an individual funding request (IFR). For support regarding IFRs, please contact: norfolkicd@nhs.net For support regarding the criteria listed below, please contact: norfolknontariff@nhs.net Please indicate whether patient meets the following NICE criteria: Please tick 1. Canagliflozin in a dual therapy regimen in combination with metformin is recommended as an option for treating type 2 diabetes, only if: ☐ Yes ☐ No • a sulfonylurea is contraindicated or not tolerated or • the person is at significant risk of hypoglycaemia or its consequences. 2. Canagliflozin in a triple therapy regimen is recommended as an option for treating type 2 diabetes in combination with: □Yes □No • metformin and a sulfonylurea or • metformin and a thiazolidinedione. 3. Canagliflozin in combination with insulin with or without other antidiabetic drugs is recommended as an ☐ Yes ☐ No option for treating type 2 diabetes. 4. People currently receiving treatment initiated within the NHS with canadiffozin that is not recommended for them by NICE in this guidance should be able to continue treatment until they and their NHS clinician ☐ Yes ☐ No consider it appropriate to stop. I confirm that the patient meets the criteria for treatment Contact Details: Name of person completing: Designation of person completing: Date:

Trust Authorising Pharmacist

Name:			
Date:			