N&WICS - TA336 - Empagliflozin in a triple therapy regimen in combination with metformin and either a sulfonylurea or a

pug	thiazolidinedione -	Type 2 diabetes		
parent/legal guardian/carer) has give	data on this form, please confirm that the appropriate explicit consent for sensions and validating subsequent invoices	tive personal information on this fo		
about the advantages and disadvant patient is likely to adhere to treatme	oved treatment available, a discussion be tages of the treatments available. This hent. The most appropriate, least expens of preference is stated in the TAs.	nas taken into consideration therap	peutic need and whether or	not the
Patient NHS No:	Trust:	Practice Name:		
Patient Hospital No:	Consultant Making Request:	Practice Postcode:		
Patient's Initials and DoB:		Practice Code:		
Notification Email Address:	(@NHS.net account	ONLY) Contact name & number:		
Start date of requested treatment:	Provider: ☐ Private ☐ NHS Supplier: ☐ Homecare ☐ H		N/A 🔽	
For support regarding IFRs, please	ests which fall outside of this use will re contact: norfolkicd@nhs.net	equire an individual funding reques	st (IFR).	
For support regarding the criteria lis	ted below, please contact: norfolknonta	riff@nhs.net		
	ted below, please contact: norfolknonta		Please tick	
Please indicate whether paties	nt meets the following NICE criteria	:		
Please indicate whether patien 1. Empagliflozin in a dual therapy treating type 2 diabetes, only if: • a sulfonylurea is contraindicated	nt meets the following NICE criteria	: n is recommended as an option for		
Please indicate whether patien 1. Empagliflozin in a dual therapy treating type 2 diabetes, only if: • a sulfonylurea is contraindicated • the person is at significant risk	nt meets the following NICE criterian regimen in combination with metforming	: n is recommended as an option for		
Please indicate whether patien 1. Empagliflozin in a dual therapy treating type 2 diabetes, only if: • a sulfonylurea is contraindicated • the person is at significant risk 2. Empagliflozin in a triple therapy	nt meets the following NICE criterian regimen in combination with metforming d or not tolerated, or of hypoglycaemia or its consequences. y regimen is recommended as an option	: n is recommended as an option for		
Please indicate whether patien 1. Empagliflozin in a dual therapy treating type 2 diabetes, only if: • a sulfonylurea is contraindicated • the person is at significant risk 2. Empagliflozin in a triple therapy combination with:	nt meets the following NICE criterian regimen in combination with metforming d or not tolerated, or of hypoglycaemia or its consequences. y regimen is recommended as an option	: n is recommended as an option for	☐Yes ☐No	
Please indicate whether patien 1. Empagliflozin in a dual therapy treating type 2 diabetes, only if: • a sulfonylurea is contraindicated. • the person is at significant risk 2. Empagliflozin in a triple therapy combination with: • metformin and a sulfonylurea or • metformin and a thiazolidinedion	nt meets the following NICE criterian regimen in combination with metforming d or not tolerated, or of hypoglycaemia or its consequences. regimen is recommended as an option one.	: n is recommended as an option for n for treating type 2 diabetes in	☐Yes ☐No	
Please indicate whether patien 1. Empagliflozin in a dual therapy treating type 2 diabetes, only if: • a sulfonylurea is contraindicated. • the person is at significant risk 2. Empagliflozin in a triple therapy combination with: • metformin and a sulfonylurea or • metformin and a thiazolidinedion 3. Empagliflozin in combination we option for treating type 2 diabetes 4. People currently receiving treating type and the patients of the p	nt meets the following NICE criterian regimen in combination with metforming d or not tolerated, or of hypoglycaemia or its consequences. regimen is recommended as an option one.	is recommended as an option for n for treating type 2 diabetes in the petic drugs is recommended as an eagliflozin that is not recommended.	Yes No	
Please indicate whether patien 1. Empagliflozin in a dual therapy treating type 2 diabetes, only if: • a sulfonylurea is contraindicated • the person is at significant risk 2. Empagliflozin in a triple therapy combination with: • metformin and a sulfonylurea or • metformin and a thiazolidinedion 3. Empagliflozin in combination wo option for treating type 2 diabetes 4. People currently receiving treat for them by NICE in this guidance	nt meets the following NICE criterian regimen in combination with metforming of or not tolerated, or of hypoglycaemia or its consequences. The regimen is recommended as an option of the combination of th	is recommended as an option for n for treating type 2 diabetes in the petic drugs is recommended as an eagliflozin that is not recommended.	Yes No	
Please indicate whether patien 1. Empagliflozin in a dual therapy treating type 2 diabetes, only if: • a sulfonylurea is contraindicated. • the person is at significant risk 2. Empagliflozin in a triple therapy combination with: • metformin and a sulfonylurea or • metformin and a thiazolidinedion 3. Empagliflozin in combination we option for treating type 2 diabetes 4. People currently receiving treation for them by NICE in this guidance consider it appropriate to stop.	nt meets the following NICE criteria regimen in combination with metforming d or not tolerated, or of hypoglycaemia or its consequences. regimen is recommended as an option one. rith insulin with or without other antidiates. treet initiated within the NHS with emple should be able to continue treatment sthe criteria for treatment	is recommended as an option for n for treating type 2 diabetes in the petic drugs is recommended as an eagliflozin that is not recommended.	Yes No	
Please indicate whether patien 1. Empagliflozin in a dual therapy treating type 2 diabetes, only if: • a sulfonylurea is contraindicated. • the person is at significant risk 2. Empagliflozin in a triple therapy combination with: • metformin and a sulfonylurea or • metformin and a thiazolidinedion 3. Empagliflozin in combination we option for treating type 2 diabetes 4. People currently receiving treation them by NICE in this guidance consider it appropriate to stop. I confirm that the patient meet	nt meets the following NICE criteria. It regimen in combination with metforming of or not tolerated, or of hypoglycaemia or its consequences. It regimen is recommended as an option of hypoglycaemia or its consequences. It is recommended as an option of hypoglycaemia or its consequences. It is recommended as an option of hypoglycaemia or its consequences. It is regimen is recommended as an option of hypoglycaemia or its consequences. It is the criteria for treatment of hypoglycaemia or its consequences. It is the criteria for treatment of hypoglycaemia or its consequences.	n is recommended as an option for treating type 2 diabetes in the petic drugs is recommended as an analysis and their NHS clinician.	Yes No	

Name:			
Date:			