Click here to access the guidelines/NICE algorithm

NOWICE TACO2 Petinemen Humaniclesmia					
N&WICS - TA623 - Patiromer - Hyperkalaemia Before providing patient identifiable data on this form, please confirm that the patient (or in the case of a minor or vulnerable adult with the parent/legal guardian/carer) has given appropriate explicit consent for sensitive personal information on this form to be passed to the CCG and/or CSU for processing this funding request and validating subsequent invoices. Consent given:					
If there is more than one NICE-approved treatment available, a discussion between the responsible clinician and the patient has taken place about the advantages and disadvantages of the treatments available. This has taken into consideration therapeutic need and whether or not the patient is likely to adhere to treatment. The most appropriate, least expensive, will be chosen (taking into account administration costs, dosage					
and price per dose) unless an order	of preference is stated in the	As.			
Patient NHS No:	Trust:		Practice Name:		
Patient Hospital No:	Consultant Making Request:		Practice Postcode:		
Patient's Initials and DoB: Practice Code:					
Notification Email Address:	(@NHS.net	account ONLY)	Contact name & number:		
Start date	Provider:				
of requested treatment:	0	care Hospital	Sub-Type:	/A 🔽	
By completing this form, you confirm					al
commissioning statement. Any requests which fall outside of this use will require an individual funding request (IFR).					
For support regarding IFRs, please contact: norfolkicd@nhs.net					
For support regarding the criteria listed below, please contact: norfolknontariff@nhs.net					
For support regarding the criteria list	ted below, please contact: norf	olknontariff@nhs.net			
For support regarding the criteria list Please indicate whether patien	·			Please tick	
	nt meets the following NICE	criteria:	d:	Please tick	
Please indicate whether patien	an option for treating hyperkal	criteria: aemia in adults only if used	d:	Please tick	
Please indicate whether patier 1. Patiromer is recommended as	an option for treating hyperkal	criteria: aemia in adults only if used		Please tick	
Please indicate whether patier 1. Patiromer is recommended as • in emergency care for acute life-	an option for treating hyperkal threatening hyperkalaemia alo	criteria: aemia in adults only if used ingside standard care or hronic kidney disease or he		Please tick ☐ Yes ☐ No	
Please indicate whether patien 1. Patiromer is recommended as • in emergency care for acute life- • for people with persistent hyperl	an option for treating hyperkal threatening hyperkalaemia alo kalaemia and stages 3b to 5 cl um level of at least 6.0 mmol/l	criteria: aemia in adults only if used ingside standard care or thronic kidney disease or he itre and	eart failure, if they:		
Please indicate whether patien 1. Patiromer is recommended as • in emergency care for acute life- • for people with persistent hyperl • have a confirmed serum potassi • are not taking, or are taking a re	an option for treating hyperkal threatening hyperkalaemia alo kalaemia and stages 3b to 5 cl um level of at least 6.0 mmol/l	criteria: aemia in adults only if used ingside standard care or thronic kidney disease or he itre and	eart failure, if they:		
Please indicate whether patien 1. Patiromer is recommended as • in emergency care for acute life- • for people with persistent hyperl • have a confirmed serum potassi • are not taking, or are taking a rebecause of hyperkalaemia and	an option for treating hyperkal threatening hyperkalaemia alo kalaemia and stages 3b to 5 cl um level of at least 6.0 mmol/leduced dosage of, a renin-angio	criteria: aemia in adults only if used ingside standard care or thronic kidney disease or he itre and	eart failure, if they:		
Please indicate whether patien 1. Patiromer is recommended as • in emergency care for acute life- • for people with persistent hyperl • have a confirmed serum potassi • are not taking, or are taking a rebecause of hyperkalaemia and • are not on dialysis.	an option for treating hyperkal an option for treating hyperkal athreatening hyperkalaemia alockalaemia and stages 3b to 5 clum level of at least 6.0 mmol/leduced dosage of, a renin-angions are no longer suitable.	criteria: aemia in adults only if used ingside standard care or inronic kidney disease or he itre and otensin-aldosterone system	eart failure, if they:	□Yes □No	
Please indicate whether patien 1. Patiromer is recommended as • in emergency care for acute life- • for people with persistent hyperl • have a confirmed serum potassi • are not taking, or are taking a rebecause of hyperkalaemia and • are not on dialysis. 2. Stop patiromer if RAAS inhibitor 3. As per NICE guidance, treatments	an option for treating hyperkal an option for treating hyperkal athreatening hyperkalaemia alockalaemia and stages 3b to 5 clum level of at least 6.0 mmol/leduced dosage of, a renin-angions are no longer suitable. The standard normally be started and any additional administration options recommended	criteria: aemia in adults only if used ingside standard care or inronic kidney disease or he itre and otensin-aldosterone system with the least expensive drawing tion costs).	eart failure, if they: n (RAAS) inhibitor rug (considering	□Yes □No	
Please indicate whether patien 1. Patiromer is recommended as in emergency care for acute life- for people with persistent hyperl have a confirmed serum potassi are not taking, or are taking a rebecause of hyperkalaemia and are not on dialysis. 2. Stop patiromer if RAAS inhibited 3. As per NICE guidance, treatmethe dose required, price per dose Please confirm that the choice of patient, noting that alternative treatmethe	an option for treating hyperkal an option for treating hyperkal athreatening hyperkalaemia alockalaemia and stages 3b to 5 clum level of at least 6.0 mmol/leduced dosage of, a renin-angions are no longer suitable. The standard normally be started and any additional administration of the standard and any additional administration options recommended applicable.	criteria: aemia in adults only if used ingside standard care or inconic kidney disease or he itre and itre and otensin-aldosterone system with the least expensive dration costs). ost-effective treatment for the inconic kidney disease or he itre and itre	eart failure, if they: n (RAAS) inhibitor rug (considering this individual cost, including the	☐Yes ☐No	
Please indicate whether patien 1. Patiromer is recommended as • in emergency care for acute life- • for people with persistent hyperl • have a confirmed serum potassi • are not taking, or are taking a re because of hyperkalaemia and • are not on dialysis. 2. Stop patiromer if RAAS inhibito 3. As per NICE guidance, treatmenthe dose required, price per dose Please confirm that the choice of patient, noting that alternative treatuse of biosimilar medications if approximately approximately and the clinical rationale for presonal recommendation.	an option for treating hyperkal an option for treating hyperkal athreatening hyperkalaemia alockalaemia and stages 3b to 5 clum level of at least 6.0 mmol/leduced dosage of, a renin-angination and any additional administration of the complete commended and any additional administration of the commended and the commended an	criteria: aemia in adults only if used ingside standard care or inconic kidney disease or he itre and itre and otensin-aldosterone system with the least expensive dration costs). ost-effective treatment for the inconic kidney disease or he itre and itre	eart failure, if they: n (RAAS) inhibitor rug (considering this individual cost, including the	☐Yes ☐No	

Designation of person completing:	Date:
Trust Authorising Pharmacist	
Name: Date:	