## Click here to access the guidelines/NICE algorithm N&WICS - TA681 - Baricitinib for treating moderate to severe atopic dermatitis - Moderate to severe atopic dermatitis Before providing patient identifiable data on this form, please confirm that the patient (or in the case of a minor or vulnerable adult with the parent/legal guardian/carer) has given appropriate explicit consent for sensitive personal information on this form to be passed to the CCG and/or CSU for processing this funding request and validating subsequent invoices. Consent given: If there is more than one NICE-approved treatment available, a discussion between the responsible clinician and the patient has taken place about the advantages and disadvantages of the treatments available. This has taken into consideration therapeutic need and whether or not the patient is likely to adhere to treatment. The most appropriate, least expensive, will be chosen (taking into account administration costs, dosage and price per dose) unless an order of preference is stated in the TAs. Patient **Practice** Trust: NHS No: Name: **Patient** Consultant **Practice** Hospital Making Postcode: No. Request: Patient's **Practice** Initials and Code: DoB: Notification Contact Email (@NHS.net account ONLY) name & Address: number: Provider: Start date ☐ Private ☐ NHS of Supplier: Sub-Type: ☐ Homecare ☐ Hospital requested treatment: (if applicable) By completing this form, you confirm that you intend to use the requested medicinal product described below as agreed in the local commissioning statement. Any requests which fall outside of this use will require an individual funding request (IFR). For support regarding IFRs, please contact: norfolkicd@nhs.net For support regarding the criteria listed below, please contact: norfolknontariff@nhs.net Please indicate whether patient meets the following NICE criteria: Please tick 1. Baricitinib is recommended as an option for treating moderate to severe atopic dermatitis in adults, only • the disease has not responded to at least 1 systemic immunosuppressant, such as ciclosporin, ☐Yes ☐No methotrexate, azathioprine and mycophenolate mofetil, or these are not suitable, and • the company provides it according to the commercial arrangement. 2. Assess response from 8 weeks and stop baricitinib if there has not been an adequate response at 16 weeks, defined as a reduction of at least: □Yes □No • 50% in the Eczema Area and Severity Index score (EASI 50) from when treatment started and • 4 points in the Dermatology Life Quality Index (DLQI) from when treatment started. 3. When using the EASI, take into account skin colour and how this could affect the EASI score, and make ☐ Yes ☐ No appropriate clinical adjustments. 4. When using the DLQI, take into account any physical, psychological, sensory or learning disabilities, or communication difficulties that could affect the responses to the DLQI, and make any appropriate ☐ Yes ☐ No adjustments. 5. As per NICE guidance, treatment should normally be started with the least expensive drug (considering the dose required, price per dose and any additional administration costs). Please confirm that the choice of drug is considered the most cost-effective treatment for this individual patient, noting that alternative treatment options recommended by NICE may be lower in cost, including the ☐Yes ☐No use of biosimilar medications if applicable. Where an alternative lower cost treatment is available and has not been tried by the patient, please confirm

that the clinical rationale for prescribing has been included in the patients' medical record for the purpose of

auditing.

Contact Details:
Date: