Click here to access the guidelines/NICE algorithm

	ANOL algorithm				
	N&WICS - TA	A383 - Golimumab - Anky	losing spondylitis		
Before providing patient identifiable parent/legal guardian/carer) has give CSU for processing this funding requ	n appropriate explici	it consent for sensitive pers	onal information on this form		
If there is more than one NICE-approabout the advantages and disadvant patient is likely to adhere to treatme	ages of the treatmen	nts available. This has taker	n into consideration theraped	utic need and wheth	ner or not the
and price per dose) unless an order	of preference is state	ed in the TAs.			
Patient NHS No:	Trust:		Practice Name:		
Patient Hospital No:	Consultant Making Request:		Practice Postcode:		
Patient's Initials and DoB:			Practice Code:		
Notification Email Address:	(@NHS.net account ONLY)	Contact name & number:		
Start date	Provider:	Drivete DAULO			
of requested treatment:	Supplier:	☐ Private ☐ NHS ☐ Homecare ☐ Hospital	Sub-Type:	J/A 🔽	
commissioning statement. Any required For support regarding IFRs, please For support regarding the criteria list	contact: norfolkicd@	nhs.net		ifk).	
Please indicate whether patier	nt meets the follow	ing NICE criteria:		Please tick	
Golimumab is recommended, vankylosing spondylitis in adults wasteroidal anti-inflammatory drugs.	☐Yes ☐No				
The choice of treatment should advantages and disadvantages of conditions such as extra-articular (taking into account administration).	☐Yes ☐No				
The response should be asses continued if there is clear evidence.			ment should only be		
• a reduction in the Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) score to 50% of the pre- treatment value or by 2 or more units and					
a reduction in the spinal pain vis	sual analogue scale ((VAS) by 2 cm or more.			
5. Treatment with another tumour necrosis factor (TNF) -alpha inhibitor is recommended for people who cannot tolerate, or whose disease has not responded to, treatment with the first TNF-alpha inhibitor, or whose disease has stopped responding after an initial response.					
6. When using BASDAI and spinal pain VAS scores, healthcare professionals should take into account any physical, sensory or learning disabilities, or communication difficulties that could affect the responses to the questionnaires, and make any adjustments they consider appropriate.					
7. As per NICE guidance, treatment should normally be started with the least expensive drug (considering the dose required, price per dose and any additional administration costs).					
Please confirm that the choice of drug is considered the most cost-effective treatment for this individual patient, noting that alternative treatment options recommended by NICE may be lower in cost, including the use of biosimilar medications if applicable.					

Where an alternative lower cost treatment is available and has not been tried by the patient, please confirm that the clinical rationale for prescribing has been included in the patients' medical record for the purpose of auditing.				
8. The product is being used as described by local commissioning position.				
I confirm that the patient meets the criteria for treatment				
Name of person completing:	Contact Details:			
Designation of person completing:	Date:			
Trust Authorising Pharmacist				
Name: Date:				