Click here to access the guidelin	es/NICE algorithm				
	N&WICS - TA814 -	Abrocitinib - Moderate to severe	atopic dermatitis		
		lease confirm that the patient (or in t icit consent for sensitive personal info			
		subsequent invoices. Consent given			
about the advantages and disadva	antages of the treatme	ilable, a discussion between the respents available. This has taken into copriate, least expensive, will be chos	nsideration theraped	itic need and whethe	er or not the
and price per dose) unless an ord	er of preference is sta	ated in the TAs.			
Patient NHS No:	Trust:		Practice Name:		
Patient Hospital No:	Consultant Making Request:		Practice Postcode:		
Patient's Initials and DoB:			Practice Code:		
Notification Email Address:		(@NHS.net account ONLY)	Contact name & number:		
Start date of requested treatment:	Provider: Supplier:	☐ Private ☐ NHS ☐ Homecare ☐ Hospital	Sub-Type:	I/A 🔽	
	quests which fall outs				ı
Please indicate whether pati	ent meets the follo	wing NICE criteria:		Please tick	
Abrocitinib and upadacitinib are recommended as options for treating moderate to severe atopic dermatitis that is suitable for systemic treatment in adults and young people 12 years and over, only if:     the disease has not responded to at least 1 systemic immunosuppressant, or these are not suitable     the companies provide abrocitinib and upadacitinib according to the commercial arrangement.				□Yes □No	
	ed as an option for tr	eating moderate to severe atopic del			
the disease has not responded to at least 1 systemic immunosuppressant, or these are not suitable					
the company provides traloking	numab according to the	ne commercial arrangement.			
3. Stop abrocitinib, upadacitinib or tralokinumab at 16 weeks if the atopic dermatitis has not responded adequately. An adequate response is:					
at least a 50% reduction in the Eczema Area and Severity Index score (EASI 50) from when treatment started and					
• at least a 4-point reduction in	the Dermatology Life	Quality Index (DLQI) from when trea	atment started.		
4. Take into account how skin colour could affect the EASI score, and make any appropriate adjustments.					
		ensory or learning disabilities, or con QL and make any appropriate adjust		☐Yes ☐No	

6. As per NICE guidance, treatment should normally be started with the least expensive drug (considering the dose required, price per dose and any additional administration costs).

Please confirm that the choice of drug is considered the most cost-effective treatment for this individual

patient, noting that alternative treatment options recommended by NICE may be lower in cost, including the use of biosimilar medications if applicable.					
Where an alternative lower cost treatment is available and has not been tried by the patient, please confirm that the clinical rationale for prescribing has been included in the patients' medical record for the purpose of auditing.					
I confirm that the patient meets the criteria for treatment					
Name of person completing:	Contact Details:				
Designation of person completing:	Date:				
Trust Authorising Pharmacist					
Name: Date:					