Click here to access the guidelines/NICE algorithm								
		N&WICS - TA21	7 - Memantine - Alzheimer's disea	ise (severe)				
p	arent/legal guardian/carer) has give	en appropriate explici	ease confirm that the patient (or in the it consent for sensitive personal informubsequent invoices. Consent given:	mation on this form				
a	bout the advantages and disadvant	ages of the treatmen	able, a discussion between the responts available. This has taken into contrate, least expensive, will be choser	sideration therapeu	tic need and wheth	ner or not the		
a	nd price per dose) unless an order	of preference is state	ed in the TAs.					
	Patient NHS No:	Trust:		Practice Name:				
	Patient Hospital No:	Consultant Making Request:		Practice Postcode:				
I	Patient's nitials and DoB:			Practice Code:				
N	lotification Email Address:	(@NHS.net account ONLY)	Contact name & number:				
	Start date of	Cummlian	☐ Private ☐ NHS	Sub-Type:				
	requested treatment:	оприст.	☐ Homecare ☐ Hospital	(if applicable)	/A 🔽			
			use the requested medicinal product le of this use will require an individua			al		
F	or support regarding IFRs, please	contact: norfolkicd@	nhs.net					
	for support regarding the criteria lis							
	Please indicate whether patier	nt meets the follow	ing NICE criteria:		Please tick			
	monotherapies are recommended	l as options for mana	onepezil, galantamine and rivastigmin aging mild to moderate Alzheimer's di 1.5.5 of the NICE guideline on demo	isease under all of	☐Yes ☐No			
	2. Memantine monotherapy is rec	commended as an op	otion for managing Alzheimer's diseas	se for people with:				
	moderate Alzheimer's disease w	who are intolerant of	or have a contraindication to AChE ir	hhibitors or	□Yes □No			
	severe Alzheimer's disease. Tre in the NICE guideline on dementiant	ere Alzheimer's disease. Treatment should be under the conditions specified in recommendation 1.5.5 NICE guideline on dementia.						
	3. This recommendation has been dementia.	n updated and replac	eed by recommendation 1.5.5 in the N	NICE guideline on	☐Yes ☐No			
	started with the drug with the low per dose once shared care has st considered appropriate when taking	cribing an AChE inhibitor (donepezil, galantamine or rivastigmine), treatment should normally be with the drug with the lowest acquisition cost (taking into account required daily dose and the price once shared care has started). However, an alternative AChE inhibitor could be prescribed if it is ead appropriate when taking into account adverse event profile, expectations about adherence, comorbidity, possibility of drug interactions and dosing profiles.						
	5. When using assessment scales to determine the severity of Alzheimer's disease, healthcare professionals should take into account any physical, sensory or learning disabilities, or communication difficulties that could affect the results and make any adjustments they consider appropriate. Healthcare professionals should also be mindful of the need to secure equality of access to treatment for patients from different ethnic groups, in particular those from different cultural backgrounds.							
			e and the need for treatment, healthd ances in which it would be inappropri					

• if the cognition score is not, or is not by itself, a clinically appropriate tool for assessing the severity of that

patient's dementia because of the patient's learning difficulties or other disabilities (for example, sensory impairments), linguistic or other communication difficulties or level of education or					
if it is not possible to apply the tool in a language in which the patient is sufficiently fluent for it to be appropriate for assessing the severity of dementia or					
• if there are other similar reasons why using a cognition score, or the score alone, would be inappropriate for assessing the severity of dementia. In such cases healthcare professionals should determine the need for initiation or continuation of treatment by using another appropriate method of assessment.					
7. As per NICE guidance, treatment should normally be started with the least expensive drug (considering the dose required, price per dose and any additional administration costs).					
Please confirm that the choice of drug is considered the most cost-effective treatment for this individual patient, noting that alternative treatment options recommended by NICE may be lower in cost, including the use of biosimilar medications if applicable.					
Where an alternative lower cost treatment is available and has not been tried by the patient, please confirm that the clinical rationale for prescribing has been included in the patients' medical record for the purpose of auditing.					
8. The product is being used as described by local commissioning position.					
I confirm that the patient meets the criteria for treatment					
Name of person completing: Contact Details:					
Designation of person completing: Date:					
Trust Authorising Pharmacist					
Name: Date:					