## Click here to access the guidelines/NICE algorithm N&WICS - TA715 - Adalimumab monotherapy - Moderate rheumatoid arthritis after conventional DMARDs have failed Before providing patient identifiable data on this form, please confirm that the patient (or in the case of a minor or vulnerable adult with the parent/legal guardian/carer) has given appropriate explicit consent for sensitive personal information on this form to be passed to the CCG and/or CSU for processing this funding request and validating subsequent invoices. Consent given: If there is more than one NICE-approved treatment available, a discussion between the responsible clinician and the patient has taken place about the advantages and disadvantages of the treatments available. This has taken into consideration therapeutic need and whether or not the patient is likely to adhere to treatment. The most appropriate, least expensive, will be chosen (taking into account administration costs, dosage and price per dose) unless an order of preference is stated in the TAs. Patient **Practice** Trust: NHS No: Name: **Patient** Consultant **Practice** Hospital Making Postcode: No. Request: Patient's **Practice** Initials and Code: DoB: Notification Contact Email (@NHS.net account ONLY) name & Address: number: Provider: Start date ☐ Private ☐ NHS of Sub-Type: Supplier: ☐ Homecare ☐ Hospital requested treatment: (if applicable) By completing this form, you confirm that you intend to use the requested medicinal product described below as agreed in the local commissioning statement. Any requests which fall outside of this use will require an individual funding request (IFR). For support regarding IFRs, please contact: norfolkicd@nhs.net For support regarding the criteria listed below, please contact: norfolknontariff@nhs.net Please indicate whether patient meets the following NICE criteria: Please tick 1. Adalimumab, etanercept and infliximab, all with methotrexate, are recommended as options for treating active rheumatoid arthritis in adults, only if: • intensive therapy with 2 or more conventional disease-modifying antirheumatic drugs (DMARDs) has not controlled the disease well enough and □Yes □No • disease is moderate (a disease activity score [DAS28] of 3.2 to 5.1) and • the companies provide adalimumab, etanercept and infliximab at the same or lower prices than those agreed with the Commercial Medicines Unit. 2. Adalimumab and etanercept can be used as monotherapy when methotrexate is contraindicated or not □Yes □No tolerated, when the criteria in 1.1 are met. 3. Continue treatment only if there is a moderate response measured using European League Against □Yes □No Rheumatism (EULAR) criteria at 6 months after starting therapy. If this initial response is not maintained, stop treatment. 4. If more than one treatment is suitable, start treatment with the least expensive drug (taking into account administration costs, dose needed and product price per dose). This may vary because of differences in how ☐ Yes ☐ No the drugs are used and treatment schedules. 5. Take into account any physical, psychological, sensory or learning disabilities, or communication ☐Yes ☐No difficulties that could affect the responses to the DAS28 and make any appropriate adjustments. 6. Abatacept with methotrexate is not recommended, within its marketing authorisation, for treating moderate active rheumatoid arthritis in adults when 1 or more DMARDs has not controlled the disease well ☐ Yes ☐ No enough.

Contact Details:

I confirm that the patient meets the criteria for treatment

Name of person completing:

Trust Authorising Pharmacist
Name: Date: