Click here to access the guidelines/NICE algorithm

Click here to access the guidelines	MICE algorithm			
	N&WICS - TA383 - Infliximab - Ankylosing	spondylitis		
parent/legal guardian/carer) has give	data on this form, please confirm that the patient (or in appropriate explicit consent for sensitive personal in uest and validating subsequent invoices. Consent give	formation on this form		
about the advantages and disadvant	oved treatment available, a discussion between the resages of the treatments available. This has taken into out. The most appropriate, least expensive, will be cho	consideration therapeu	tic need and wheth	ner or not the
and price per dose) unless an order	of preference is stated in the TAs.			
Patient NHS No:	Trust:	Practice Name:		
Patient Hospital No:	Consultant Making Request:	Practice Postcode:		
Patient's Initials and DoB:		Practice Code:		
Notification Email Address:	(@NHS.net account ONLY)	Contact name & number:		
Start date of requested treatment:	Provider: ☐ Private ☐ NHS Supplier: ☐ Homecare ☐ Hospital	Sub-Type:	/A 🔽	
For support regarding IFRs, please For support regarding the criteria list	contact: norfolkicd@nhs.net ed below, please contact: norfolknontariff@nhs.net			
Please indicate whether patier	t meets the following NICE criteria:		Please tick	
	hin it's marketing authorisations, as an option for treat hose disease has responded inadequately to, or who		Пусь Пы	
	if treatment is started with the least expensive inflixing mab should be able to continue treatment with the sar	·	☐ Yes ☐ No	
until they and their NHS clinician				
advantages and disadvantages of conditions such as extra-articular	be made after discussion between the clinician and the treatments available. This may include considerin manifestations. If more than 1 treatment is suitable, to costs and patient access schemes) should be chost	g associated he least expensive	☐Yes ☐No	
3. The response should be asses continued if there is clear evidence	sed 12 weeks after the start of treatment. Treatment se of response, defined as:	should only be		
a reduction in the Bath Ankylosi treatment value or by 2 or more u	ng Spondylitis Disease Activity Index (BASDAI) score nits and	to 50% of the pre-	□Yes □No	
a reduction in the spinal pain vis	sual analogue scale (VAS) by 2 cm or more.			
	necrosis factor (TNF) -alpha inhibitor is recommended has not responded to, treatment with the first TNF-al onding after an initial response.		☐Yes ☐No	
physical, sensory or learning disa	al pain VAS scores, healthcare professionals should t bilities, or communication difficulties that could affect justments they consider appropriate.		□Yes □No	
	ent should normally be started with the least expensive and any additional administration costs).	e drug (considering		

Please confirm that the choice of drug is considered the most cost-effective treatment for this individual patient, noting that alternative treatment options recommended by NICE may be lower in cost, including the use of biosimilar medications if applicable. Where an alternative lower cost treatment is available and has not been tried by the patient, please confirm that the clinical rationale for prescribing has been included in the patients' medical record for the purpose of auditing.			
8. The product is being used as described by local commissioning position.			
I confirm that the patient meets the criteria for treatment			
Name of person completing:	Contact Details:		
Designation of person completing: Date:			
Trust Authorising Pharmacist			
Name: Date:			