

Derivation reduction of metarules in meta-interpretive learning

Andrew Cropper & Sophie Tournet

Input	Output
Examples Background knowledge Bias	Logic program

Biases

- Mode declarations (Progol, ILASP, Aleph, XHAIL, ...)
- **Metarules** (Metagol, MIL-Hex, ∂ ILP, ProPPR, Clint, MOBAL ...)

Metarules

$$\frac{\exists P Q \forall A B \quad P(A, B) \leftarrow Q(A, B)}{\text{Metarule 1}}$$

$$\frac{\exists P Q R \forall A B \quad P(A, B) \leftarrow Q(A), R(A, B)}{\text{Metarule 2}}$$

$$\frac{\exists P Q R \forall A B C \quad P(A, B) \leftarrow Q(A, C), R(C, B)}{\text{Metarule 3}}$$

Metarules

$$P(A, B) \leftarrow Q(A, B)$$

$$P(A, B) \leftarrow Q(A), R(A, B)$$

$$P(A, B) \leftarrow Q(A, C), R(C, B)$$

P, Q, R are **existentially** quantified **second-order** variables

A, B, C are **universally** quantified **first-order** variables

Input	Output
<pre>% background parent(ann,amy)← parent(amy,amelia)← % example grandparent(ann,amelia)← % metarule P(A,B)←Q(A,C),R(C,B)</pre>	

Input	Output
<pre>% background parent(ann,amy)← parent(amy,amelia)←</pre>	<pre>grandparent(A,B)← parent(A,C), parent(C,B)</pre>
<pre>% example grandparent(ann,amelia)←</pre>	<pre>{ P\granparent, Q\parent,</pre>
<pre>% metarule P(A,B)←Q(A,C),R(C,B)</pre>	<pre> R\parent }</pre>

Why?

Completeness

cannot learn grandparent/2 with only $P(X) \leftarrow Q(X)$

Efficiency

more metarules = larger hypothesis space

Remove redundant metarules [ILP14]

The Horn clause C is **entailment redundant** in the Horn theory $T \cup \{C\}$ when $T \models C$

Entailment redundancy

$$C1 = h(A, B) \leftarrow s(A, B)$$

$$C2 = h(A, B) \leftarrow s(A, B), u(B)$$

$$C3 = h(A, B) \leftarrow s(A, B), u(A, B)$$

$$C4 = h(A, B) \leftarrow s(A, B), u(A, B), v(A, B)$$

Entailment redundancy

$C1 = h(A, B) \leftarrow s(A, B)$

~~$C2 = h(A, B) \leftarrow s(A, B), u(B)$~~

~~$C3 = h(A, B) \leftarrow s(A, B), u(A, B)$~~

~~$C4 = h(A, B) \leftarrow s(A, B), u(A, B), v(A, B)$~~

$\{C1\} \models \{C2, C3, C4\}$

Entailment reduction of metarules [ILP14]

$P(A, B) \leftarrow Q(A, B)$

$P(A, B) \leftarrow Q(B, A)$

$P(A, B) \leftarrow Q(A, C), R(B, C)$

$P(A, B) \leftarrow Q(A, C), R(C, B)$

$P(A, B) \leftarrow Q(B, A), R(A, B)$

$P(A, B) \leftarrow Q(B, A), R(B, A)$

$P(A, B) \leftarrow Q(B, C), R(A, C)$

$P(A, B) \leftarrow Q(B, C), R(C, A)$

$P(A, B) \leftarrow Q(C, A), R(B, C)$

$P(A, B) \leftarrow Q(C, A), R(C, B)$

$P(A, B) \leftarrow Q(C, B), R(A, C)$

$P(A, B) \leftarrow Q(C, B), R(C, A)$



?

Entailment reduction of metarules [ILP14]

$P(A, B) \leftarrow Q(A, B)$
 $P(A, B) \leftarrow Q(B, A)$
 $P(A, B) \leftarrow Q(A, C), R(B, C)$
 $P(A, B) \leftarrow Q(A, C), R(C, B)$
 $P(A, B) \leftarrow Q(B, A), R(A, B)$
 $P(A, B) \leftarrow Q(B, A), R(B, A)$
 $P(A, B) \leftarrow Q(B, C), R(A, C)$
 $P(A, B) \leftarrow Q(B, C), R(C, A)$
 $P(A, B) \leftarrow Q(C, A), R(B, C)$
 $P(A, B) \leftarrow Q(C, A), R(C, B)$
 $P(A, B) \leftarrow Q(C, B), R(A, C)$
 $P(A, B) \leftarrow Q(C, B), R(C, A)$



$P(A, B) \leftarrow Q(B, A)$

$P(A, B) \leftarrow Q(A, C), R(C, B)$

Entailment redundancy

C1 = $P(A, B) \leftarrow Q(A, B)$

C2 = $P(A, B) \leftarrow Q(A, B), R(A)$

C3 = $P(A, B) \leftarrow Q(A, B), R(A, B)$

C4 = $P(A, B) \leftarrow Q(A, B), R(A, B), S(A, B)$

Entailment redundancy

$C1 = P(A,B) \leftarrow Q(A,B)$

~~$C2 = P(A,B) \leftarrow Q(A,B), R(A)$~~

~~$C3 = P(A,B) \leftarrow Q(A,B), R(A,B)$~~

~~$C4 = P(A,B) \leftarrow Q(A,B), R(A,B), S(A,B)$~~

$\{C1\} \models \{C2, C3, C4\}$

Entailment redundancy

$C1 = P(A,B) \leftarrow Q(A,B)$

~~$C2 = P(A,B) \leftarrow Q(A,B), R(A)$~~

~~$C3 = P(A,B) \leftarrow Q(A,B), R(A,B)$~~

~~$C4 = P(A,B) \leftarrow Q(A,B), R(A,B), S(A,B)$~~

$\{C1\} \models \{C2, C3, C4\}$

$\text{father}(A,B) \leftarrow \text{parent}(A,B), \text{male}(A)$ ✖

Derivation redundancy

The Horn clause C is **derivationally redundant** in the Horn theory $T \cup \{C\}$ when $T \vdash C$

SLD-resolution



Derivation redundancy

C1 = $P(A, B) \leftarrow Q(A, B)$

C2 = $P(A, B) \leftarrow Q(A, B), R(A)$

C3 = $P(A, B) \leftarrow Q(A, B), R(A, B)$

C4 = $P(A, B) \leftarrow Q(A, B), R(A, B), S(A, B)$

Derivation redundancy

$C1 = P(A,B) \leftarrow Q(A,B)$

$C2 = P(A,B) \leftarrow Q(A,B), R(A)$

$C3 = P(A,B) \leftarrow Q(A,B), R(A,B)$

~~$C4 = P(A,B) \leftarrow Q(A,B), R(A,B), S(A,B)$~~

$\{C1, C2, C3\} \vdash \{C4\}$

$\text{father}(A,B) \leftarrow \text{parent}(A,B), \text{male}(A) \quad \checkmark$

Derivation redundancy

While there is a clause in T such that $T - \{C\} \vdash_k C$:

Set $T = T - \{C\}$

Connected clauses

body literals are connected to the head literal

$P(A) \leftarrow Q(A)$ ✓

$P(A, B) \leftarrow Q(A, C)$ ✓

$P(A, B) \leftarrow Q(A, B), R(B, D), S(D, B)$ ✓

$P(A) \leftarrow \mathbf{Q(B)}$ ✗

$P(A) \leftarrow Q(A), \mathbf{R(B, C)}$ ✗

$P(A, B) \leftarrow Q(A, B), \mathbf{S(C)}$ ✗

H^2_m

restriction on literal arity

$P(A, B) \leftarrow Q(A, B) \quad \checkmark$

$P(A) \leftarrow Q(A, B), R(B) \quad \checkmark$

$P(A, B, C) \leftarrow \mathbf{Q(A, B, C)} \quad \times$

$P(A) \leftarrow \mathbf{Q(A, B, C)}, R(B, C) \quad \times$

$$H^2_m$$

$$P(A, B) \leftarrow Q(A, B) \quad \checkmark$$

$$P(A, B) \leftarrow Q(A, C), R(C, B) \quad \checkmark$$

$$P(A) \leftarrow \mathbf{Q(A)} \quad \times$$

$$P(A, B) \leftarrow Q(A, B), \mathbf{R(B)} \quad \times$$

H^a_2

restriction on number of body literals

$P(A, B) \leftarrow Q(A, B) \quad \checkmark$

$P(A) \leftarrow Q(A, B, C), R(B, C) \quad \checkmark$

$P(A) \leftarrow Q(A), R(A), S(A) \quad \times$

$P(A, B) \leftarrow Q(A), R(B), S(A, B) \quad \times$

$H^a_2 =$

$P(A) \leftarrow Q(A), R(A) \quad \checkmark$

$P(A, B) \leftarrow Q(A, B), R(A, B) \quad \checkmark$

$P(A) \leftarrow \mathbf{Q(A)} \quad \times$

$P(A, B) \leftarrow \mathbf{Q(A, B), R(B)} \quad \times$

Exactly-two connected

each first-order variable appears exactly twice

$P(A) \leftarrow Q(A)$ ✓

$P(A, B) \leftarrow Q(A, B)$ ✓

$P(A, B) \leftarrow Q(A, C), R(C, B)$ ✓

$P(A, \mathbf{B}) \leftarrow Q(A)$ ✗

$P(A) \leftarrow Q(A, \mathbf{B})$ ✗

$P(\mathbf{A}) \leftarrow Q(A), R(A)$ ✗

$$E^2=5$$

$$E^2 =_5$$

E-reduction	D-reduction
$P(A, B) \leftarrow Q(B, A)$ $P(A, B) \leftarrow Q(A, C), R(C, B)$	$P(A, B) \leftarrow Q(B, A)$ $P(A, B) \leftarrow Q(A, C), R(C, B)$

$$E^2=5$$

E-reduction	D-reduction
$P(A, B) \leftarrow Q(B, A)$	$P(A, B) \leftarrow Q(B, A)$
$P(A, B) \leftarrow Q(A, C), R(C, B)$	$P(A, B) \leftarrow Q(A, C), R(C, B)$

$$E^2=2 \vdash E^2=\infty \checkmark$$

$$E^2_5$$

E-reduction	D-reduction
$P(A) \leftarrow Q(A)$	$P(A) \leftarrow Q(A)$
$P(A) \leftarrow Q(A, B), R(B)$	$P(A) \leftarrow Q(A, B), R(B)$
$P(A, B) \leftarrow Q(B, A)$	$P(A, B) \leftarrow Q(B, A)$
$P(A, B) \leftarrow Q(A), R(B)$	$P(A, B) \leftarrow Q(A), R(B)$
$P(A, B) \leftarrow Q(A, C), R(C, B)$	$P(A, B) \leftarrow Q(A, C), R(C, B)$

E^2_5

E-reduction	D-reduction
$P(A) \leftarrow Q(A)$	$P(A) \leftarrow Q(A)$
$P(A) \leftarrow Q(A, B), R(B)$	$P(A) \leftarrow Q(A, B), R(B)$
$P(A, B) \leftarrow Q(B, A)$	$P(A, B) \leftarrow Q(B, A)$
$P(A, B) \leftarrow Q(A), R(B)$	$P(A, B) \leftarrow Q(A), R(B)$
$P(A, B) \leftarrow Q(A, C), R(C, B)$	$P(A, B) \leftarrow Q(A, C), R(C, B)$

$E^2_2 \vdash E^2_\infty \checkmark$

Two connected

each first-order variable appears at least twice
(i.e. forbids singleton variables)

$P(A) \leftarrow Q(A)$ ✓

$P(A) \leftarrow Q(A), R(A)$ ✓

$P(A, B) \leftarrow Q(A, B), R(B)$ ✓

$P(A, B) \leftarrow Q(A, C), R(C, B)$ ✓

$P(A, \mathbf{B}) \leftarrow Q(A)$ ✗

$P(A) \leftarrow Q(A, \mathbf{B})$ ✗

$P(A) \leftarrow Q(A), R(A, \mathbf{B})$ ✗

$K^2=5$ two connected

$K^2=5$ two connected

E-reduction	D-reduction
$P(A,B) \leftarrow Q(B,A)$	$P(A,B) \leftarrow Q(B,A)$
$P(A,B) \leftarrow Q(A,C), R(C,B)$	$P(A,B) \leftarrow Q(A,C), R(C,B)$
	$P(A,B) \leftarrow Q(A,B), R(A,B)$
	$P(A,B) \leftarrow Q(A,B), R(A,C), S(C,D), T(C,D)$
	$P(A,B) \leftarrow Q(A,C), R(A,C), S(B,D), T(B,D)$
	$P(A,B) \leftarrow Q(A,C), R(A,D), S(B,C), T(B,D), U(C,D)$

$K^2=5$ two connected

E-reduction	D-reduction
$P(A,B) \leftarrow Q(B,A)$	$P(A,B) \leftarrow Q(B,A)$
$P(A,B) \leftarrow Q(A,C), R(C,B)$	$P(A,B) \leftarrow Q(A,C), R(C,B)$
	$P(A,B) \leftarrow Q(A,B), R(A,B)$
	$P(A,B) \leftarrow Q(A,B), R(A,C), S(C,D), T(C,D)$
	$P(A,B) \leftarrow Q(A,C), R(A,C), S(B,D), T(B,D)$
	$P(A,B) \leftarrow Q(A,C), R(A,D), S(B,C), T(B,D), U(C,D)$

$K^2=2 \not\models K^2=\infty$ ✖

K^2_5 two connected

K^2_5 two connected

E-reduction	D-reduction
$P(A) \leftarrow Q(A)$	$P(A) \leftarrow Q(A)$
$P(A) \leftarrow R(A, B), Q(A, B)$	$P(A) \leftarrow R(A, B), Q(A, B)$
	$P(A) \leftarrow Q(A), R(A)$
	$P(A) \leftarrow Q(B), R(A, B)$
$P(A, B) \leftarrow Q(B, A)$	$P(A, B) \leftarrow Q(B, A)$
$P(A, B) \leftarrow Q(A), R(B)$	$P(A, B) \leftarrow Q(A), R(B)$
$P(A, B) \leftarrow Q(A, C), R(C, B)$	$P(A, B) \leftarrow Q(A, C), R(C, B)$
	$P(A, B) \leftarrow Q(A, B), R(A, B)$
	$P(A, B) \leftarrow Q(A), R(A, B)$
	$P(A, B) \leftarrow Q(A, C), R(A, D), S(C, B), T(B, D), U(C, D)$

K^2_5 two connected


E-reduction	D-reduction
$P(A) \leftarrow Q(A)$	$P(A) \leftarrow Q(A)$
$P(A) \leftarrow R(A, B), Q(A, B)$	$P(A) \leftarrow R(A, B), Q(A, B)$
	$P(A) \leftarrow Q(A), R(A)$
	$P(A) \leftarrow Q(B), R(A, B)$
$P(A, B) \leftarrow Q(B, A)$	$P(A, B) \leftarrow Q(B, A)$
$P(A, B) \leftarrow Q(A), R(B)$	$P(A, B) \leftarrow Q(A), R(B)$
$P(A, B) \leftarrow Q(A, C), R(C, B)$	$P(A, B) \leftarrow Q(A, C), R(C, B)$
	$P(A, B) \leftarrow Q(A, B), R(A, B)$
	$P(A, B) \leftarrow Q(A), R(A, B)$
	$P(A, B) \leftarrow Q(A, C), R(A, D), S(C, B), T(B, D), U(C, D)$

$K^2=2 \not\models K^2=5 \times$

K^2_5 two connected

E-reduction	D-reduction
$P(A) \leftarrow Q(A)$	$P(A) \leftarrow Q(A)$
$P(A) \leftarrow R(A, B), Q(A, B)$	$P(A) \leftarrow R(A, B), Q(A, B)$
	$P(A) \leftarrow Q(A), R(A)$
	$P(A) \leftarrow Q(B), R(A, B)$
$P(A, B) \leftarrow Q(B, A)$	$P(A, B) \leftarrow Q(B, A)$
$P(A, B) \leftarrow Q(A), R(B)$	$P(A, B) \leftarrow Q(A), R(B)$
$P(A, B) \leftarrow Q(A, C), R(C, B)$	$P(A, B) \leftarrow Q(A, C), R(C, B)$
	$P(A, B) \leftarrow Q(A, B), R(A, B)$
	$P(A, B) \leftarrow Q(A), R(A, B)$
	$P(A, B) \leftarrow Q(A, C), R(A, D), S(C, B), T(B, D), U(C, D)$


K^2_∞ cannot be reduced ✕




$$P_0(x_1, x_2) \leftarrow P_1(x_1, x_3), P_2(x_1, x_4), P_3(x_2, x_3), P_4(x_2, x_4), P_5(x_3, x_4)$$

Does it matter?

1. TRAINS GOING EAST



2. TRAINS GOING WEST



Accuracies

Task	E-reduction	D-reduction	D*-reduction
T1	95 \pm 1	100 \pm 0	100 \pm 0
T2	99 \pm 1	100 \pm 0	100 \pm 0
T3	56 \pm 3	96 \pm 2	96 \pm 2
T4	69 \pm 4	96 \pm 2	96 \pm 2
T5	59 \pm 3	93 \pm 3	93 \pm 3
T6	50 \pm 1	96 \pm 3	96 \pm 3
T7	68 \pm 4	95 \pm 2	95 \pm 2
T8	54 \pm 3	60 \pm 3	90 \pm 3

Learning times

Task	E-reduction	D-reduction	D*-reduction
T1	0.01 ± 0	0 ± 0	0 ± 0
T2	0.01 ± 0	0 ± 0	0 ± 0
T3	431 ± 59	0.01 ± 0	0.01 ± 0
T4	300 ± 68	0 ± 0	0.01 ± 0
T5	427 ± 60	1 ± 0.3	1 ± 0.41
T6	600 ± 0	1 ± 0.41	1 ± 0.42
T7	917 ± 535	1 ± 0.27	1 ± 0.36
T8	487 ± 51	360 ± 67	26 ± 5

```
% target program  
f(X):-has_car(X,C1),  
      long(C1),  
      two_wheels(C1),  
      has_car(X,C2),  
      long(C2),  
      three_wheels(C2).
```

% E-reduction

$f(A) :- \text{has_car}(A, B), f1(A, B).$

$f1(A, B) :- \text{has_car}(A, C), f2(C, B).$

$f2(A, B) :- \text{long}(A), \text{three_wheels}(B).$

% D-reduction

$f(A) :- \neg f_1(A), f_2(A).$

$f_1(A) :- \neg \text{has_car}(A, B), \text{three_wheels}(B).$

$f_2(A) :- \neg \text{has_car}(A, B), \text{roof_open}(B).$

% D*-reduction

$f(A) : \neg f_1(A), f_2(A).$

$f_1(A) : \neg \text{has_car}(A, B), \text{three_wheels}(B).$

$f_2(A) : \neg \text{has_car}(A, B), f_3(B).$

$f_3(A) : \neg \text{long}(A), \text{two_wheels}(A).$

% target program

f(X):-

 has_car(X,C1),
 long(C1),
 two_wheels(C1),
 has_car(X,C2),
 long(C2),
 three_wheels(C2).

% D*-reduction

f(A):-

 has_car(A,B),
 three_wheels(B),
 has_car(A,C),
 long(C),
 two_wheels(C).

Todo

- Study derivation reduction problem
- Other fragments
 - Triadics
 - Connected
- Unconstrained resolution